

1990–2021

New Zealand's Greenhouse Gas Inventory: Volume 2, annexes

Fulfilling reporting requirements under the United Nations
Framework Convention on Climate Change and the Kyoto Protocol



Ministry for the
Environment
Manatū Mō Te Taiao



Te Kāwanatanga o Aotearoa
New Zealand Government

Table A3.2.15 (p. 106) of this publication was updated on 19 January 2024.

Contents

Tables	v
Figures	xii
Annex 1: Key categories	1
A1.1 Methodology used for identifying key categories	1
A1.2 Disaggregation	2
A1.3 Tables 4.2 to 4.3 of the 2006 IPCC Guidelines (General Guidance and Reporting)	2
Annex 1: References	21
Annex 2: Uncertainty analysis	22
A2.1 Approach 1 uncertainty calculation	22
Annex 2: References	34
Annex 3: Detailed methodological information for other sectors	35
A3.1 Supplementary information for the Agriculture sector	35
A3.2 Supplementary information for the LULUCF sector	45
Annex 3: References	132
Annex 4: Methodology and data collection for estimating emissions from fossil fuel combustion	141
A4.1 Emissions from liquid fuels	144
A4.2 Emissions from solid fuels	147
A4.3 Emissions from gaseous fuels	148
A4.4 Energy balance	148
A4.5 Carbon dioxide reference approach for the Energy sector	149
Annex 4: References	151
Annex 5: Supplementary information for the KP-LULUCF sector	152
A5.1 Technical corrections to the FMRL	152
A5.2 Natural disturbance	163
A5.3 Carbon equivalent forests	165
Annex 5: References	168
Annex 6: Additional information on the inventory system and inventory completeness	169
A6.1 Quality assurance and quality control processes	169
A6.2 General assessment of completeness	172
Annex 6: References	180

Annex 7: Tokelau	181
A7.1 Emissions estimate data and relevant supporting information by category for Tokelau	181
Annex 8: Agricultural emissions from fertilisers and by livestock type	250
A8.1 Agricultural emissions from fertilisers	250
A8.2 Agricultural emissions by livestock type	252

See volume 1, chapters 1–15, for the National Inventory Report, which is presented separately.

Tables

Table A1.3.1(a)	Results of the key category level analysis for 99 per cent of the net emissions and removals for New Zealand in 2021	2
Table A1.3.1(b)	Results of the key category level analysis for 99 per cent of the gross emissions and removals for New Zealand in 2021	5
Table A1.3.2(a)	Results of the level analysis for 99 per cent of the net emissions and removals for New Zealand in 1990 included for reference only	7
Table A1.3.2(b)	Results of the level analysis for 99 per cent of the gross emissions and removals for New Zealand in 1990 included for reference only	10
Table A1.3.3(a)	Results of the key category trend analysis for 99 per cent of the net emissions and removals for New Zealand in 1990 to 2021	12
Table A1.3.3(b)	Results of the key category trend analysis for 99 per cent of the gross emissions and removals for New Zealand in 1990–2021	16
Table A2.1.1	Uncertainty calculation (including LULUCF) for New Zealand’s Greenhouse Gas Inventory 1990–2021 (IPCC, 2006, Approach 1)	24
Table A2.1.2	Uncertainty calculation (excluding LULUCF) for New Zealand’s Greenhouse Gas Inventory 1990–2021 (IPCC, 2006, Approach 1)	29
Table A3.1.1	Imputation levels and sampling errors for recent Agricultural Production Surveys	35
Table A3.1.2	Characterisation of major livestock subcategories (dairy cattle, beef cattle, sheep and deer) in New Zealand’s Tier 2 livestock modelling	36
Table A3.1.3	Proportion of annual milk yield each month for major livestock categories	37
Table A3.1.4	Methane emission factors for Tier 1 enteric fermentation livestock and manure management	38
Table A3.1.5	Emission factors for New Zealand’s agricultural nitrous oxide emissions	38
Table A3.1.6	Direct nitrous oxide emission factors for urine deposited by cattle, sheep and deer, by livestock type and slope	39
Table A3.1.7	Parameter values for New Zealand’s agriculture nitrous oxide emissions	39
Table A3.1.8	Parameter values for New Zealand’s cropping emissions	40
Table A3.1.9	Allocation of urine deposition to low slope (0–12 degrees) and steep slope (more than 24 degrees), split by the percentage of low slope and steep slope land available	41
Table A3.1.10	Share of livestock population, and amount of urine nitrogen (N) deposition in 2021, by Beef + Lamb New Zealand Ltd farm class	43
Table A3.1.11	Proportion of total sheep, beef and deer land on different hill slopes, by Beef + Lamb New Zealand Ltd farm class, for 2020/21	43
Table A3.1.12	Proportion of total sheep, beef and deer urine nitrogen deposited on different hill slopes, by Beef + Lamb New Zealand Ltd farm class, for 2021	44
Table A3.2.1	Satellite imagery used for land use mapping in 1990, 2008, 2012 and 2016	45
Table A3.2.2	Ancillary mosaicked imagery data sets used in land use mapping	50
Table A3.2.3	Methods used to estimate Afforestation total area and trends between 2008 and 2016	56
Table A3.2.4	Annual land-use changes (units in 000s hectares)	58
Table A3.2.5	Land use effect coefficients with standard errors, <i>t</i> -values, and corresponding <i>p</i> -value significance estimates, extracted from full model results	76

Table A3.2.6	Steady state soil organic carbon stocks, with 95 per cent confidence intervals, calculated from Soil CMS model	76
Table A3.2.7	New Zealand emission factors for organic soils	81
Table A3.2.8	Proportion of total destocking area by age across all planted forest, 1990–2021	85
Table A3.2.9	Summary of methods used to calculate New Zealand’s natural forest biomass carbon stock from plot data	88
Table A3.2.10	Summary of root:shoot ratios applied to the different taxonomic groups in pre-1990 natural forest	89
Table A3.2.11	Influence of individual site and management factors on predicted wood density for New Zealand planted forest	92
Table A3.2.12	Comparison of the average biomass carbon stock per hectare for each forest type, calculated as the area weighted average of plot measurements and yield table values	100
Table A3.2.13	Pre-1990 natural forest – tall forest carbon stocks by year (tonnes C ha ⁻¹) ¹	104
Table A3.2.14	Pre-1990 natural forest – regenerating forest carbon stocks by year (tonnes C ha ⁻¹) ¹	105
Table A3.2.15	Post-1989 natural forest yield table (tonnes C ha ⁻¹) ¹	106
Table A3.2.16	Post-1989 natural forest yield table (tonnes C ha ⁻¹) ¹ (for transitions from grassland with woody biomass)	106
Table A3.2.17	Pre-1990 ‘planted before 1990’ planted forest yield table (tonnes C ha ⁻¹) ¹	107
Table A3.2.18	Pre-1990 ‘planted 1990 onwards’ planted forest yield table (tonnes C ha ⁻¹) ¹	109
Table A3.2.19	Post-1989 planted forest yield table (tonnes C ha ⁻¹) ¹	110
Table A3.2.20	Harvested wood product type, waste and fuel product type, exported volume in 2015 and assumed half-lives for China	112
Table A3.2.21	Harvested wood product type, waste and fuel product type, exported volume in 2015 and assumed half-lives for South Korea	112
Table A3.2.22	Harvested wood product type, waste and fuel product type, exported volume in 2015 and assumed half-lives for India	112
Table A3.2.23	Uncertainty in New Zealand’s 2021 carbon estimates from pre-1990 natural forest (including land in transition)	116
Table A3.2.24	Uncertainty in New Zealand’s 2021 carbon estimates from post-1989 natural forest (including land in transition)	116
Table A3.2.25	Uncertainty in New Zealand’s 2021 carbon estimates from pre-1990 planted forest (including land in transition)	117
Table A3.2.26	Uncertainty in New Zealand’s 2021 carbon estimates from post-1989 planted forest (including land in transition)	117
Table A3.2.27	Uncertainty in New Zealand’s 2020 Cropland carbon estimates (including land in transition)	118
Table A3.2.28	Uncertainty in New Zealand’s 2021 carbon estimates for the <i>Grassland</i> category (including land in transition)	119
Table A3.2.29	Uncertainty in New Zealand’s 2021 carbon estimates for the <i>Wetlands</i> category (including land in transition)	119
Table A3.2.30	Uncertainty in New Zealand’s 2021 carbon estimates for the <i>Settlements</i> category (including land in transition)	120
Table A3.2.31	Uncertainty in New Zealand’s 2021 carbon estimates for the land use category <i>Other land</i> (including land in transition)	120
Table A3.2.32	Uncertainty in <i>Harvested wood products</i> data and parameters	121

Table A3.2.33	Uncertainty in New Zealand’s 2021 carbon estimates from emissions associated with <i>Harvested wood products</i>	121
Table A3.2.34	Uncertainty in New Zealand’s 2020 estimates from nitrous oxide (N ₂ O) emissions associated with land-use change	122
Table A3.2.35	Uncertainty analysis for the LULUCF sector	123
Table A4.1	Gross carbon dioxide emission factors used for New Zealand’s energy sector in 2021	141
Table A4.2	Consumption-weighted average emission factors used for New Zealand’s sub-bituminous coal-fired electricity generation for 1990 to 2021	142
Table A4.3	Methane emission factors used for New Zealand’s energy sector for 1990 to 2021	142
Table A4.4	Nitrous oxide emission factors used for New Zealand’s energy sector for 1990 to 2021	143
Table A4.5	Carbon content (per cent mass) for liquid fuels for 1990 to 2021	144
Table A4.6	Emission factors for European gasoline and diesel vehicles – COPERT IV model (European Environment Agency, 2007)	145
Table A4.7	Split of energy use and non-energy use of natural gas in petajoules	148
Table A5.1.1	Pre-1990 planted forest data used to estimate emissions for the technically corrected FMRL	157
Table A5.1.2	Summary of the technical corrections to the FMRL	161
Table A5.3.1	Breakdown of carbon equivalent forests by domestic scheme application in 2021	165
Table A6.1.1	Quality assurance and quality control processes used in preparation of the inventory	169
Table A6.2.1	Summary of ‘NE’ (not estimated) entries in 2023 submission	172
Table A6.2.2	Emissions reported using the ‘IE’ (included elsewhere) notation key	175
Tokelau CRF Table 1.A.1.a: [1. Energy] [1.AA Fuel Combustion – Sectoral approach] [1.A.1 Energy Industries] [1.A.1.a Public Electricity and Heat Production] (Part 1 of 3)		181
Tokelau CRF Table 1.A.1.a: [1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.1 Energy Industries][1.A.1.a Public Electricity and Heat Production] (Part 2 of 3)		182
Tokelau CRF Table 1.A.1.a: [1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.1 Energy Industries][1.A.1.a Public Electricity and Heat Production] (Part 3 of 3)		183
Tokelau CRF Table 1.A.3.b.i: [1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Gasoline] (Part 1 of 3)		184
Tokelau CRF Table 1.A.3.b.i: [1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Gasoline] (Part 2 of 3)		185
Tokelau CRF Table 1.A.3.b.i: [1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Gasoline] (Part 3 of 3)		186
Tokelau CRF Table 1.A.3.b.i Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Diesel Oil] (Part 1 of 3)		187
Tokelau CRF Table 1.A.3.b.i Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Diesel Oil] (Part 2 of 3)		187
Tokelau CRF Table 1.A.3.b.i Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.3 Transport] [1.A.3.b Road Transportation][1.A.3.b.i Cars][Diesel Oil] (Part 3 of 3)		188
Tokelau CRF Table 1.A.3.d Gas/Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.3 Transport][1.A.3.d Domestic Navigation][Gas/Diesel Oil] (Part 1 of 3)		189

Tokelau CRF Table 1.A.3.d Gas/Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.3 Transport][1.A.3.d Domestic Navigation][Gas/Diesel Oil] (Part 2 of 3)	190
Tokelau CRF Table 1.A.3.d Gas/Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.3 Transport][1.A.3.d Domestic Navigation][Gas/Diesel Oil] (Part 3 of 3)	190
Tokelau CRF Table 1.A.4.b: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.4 Other Sectors][1.A.4.b Residential] (Part 1 of 3)	191
Tokelau CRF Table 1.A.4.b: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.4 Other Sectors][1.A.4.b Residential] (Part 2 of 3)	193
Tokelau CRF Table 1.A.4.b: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.4 Other Sectors][1.A.4.b Residential] (Part 3 of 3)	194
Tokelau CRF Table 1.A.4.c.iii Gas/Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.4 Other Sectors][1.A.4.c Agriculture/Forestry/Fishing][1.A.4.c.iii Fishing] [Gas/Diesel Oil] (Part 1 of 3)	196
Tokelau CRF Table 1.A.4.c.iii Gas/Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.4 Other Sectors][1.A.4.c Agriculture/Forestry/Fishing][1.A.4.c.iii Fishing] [Gas/Diesel Oil] (Part 2 of 3)	197
Tokelau CRF Table 1.A.4.c.iii Gas/Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.4 Other Sectors][1.A.4.c Agriculture/Forestry/Fishing][1.A.4.c.iii Fishing] [Gas/Diesel Oil] (Part 3 of 3)	198
Tokelau CRF Table 1.AB Gasoline: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Gasoline] (Part 1 of 3)	199
Tokelau CRF Table 1.AB Gasoline: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Gasoline] (Part 2 of 3)	199
Tokelau CRF Table 1.AB Gasoline: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Gasoline] (Part 3 of 3)	200
Tokelau CRF Table 1.AB Gas Diesel Oil: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Gas / Diesel Oil] (Part 1 of 3)	201
Tokelau CRF Table 1.AB Gas Diesel Oil: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Gas / Diesel Oil] (Part 2 of 3)	202
Tokelau CRF Table 1.AB Gas Diesel Oil: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Gas / Diesel Oil] (Part 3 of 3)	202
Tokelau CRF Table 1.AB Other Kerosene: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Other Kerosene] (Part 1 of 3)	203
Tokelau CRF Table 1.AB Other Kerosene: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Other Kerosene] (Part 2 of 3)	204
Tokelau CRF Table 1.AB Other Kerosene: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Other Kerosene] (Part 3 of 3)	205
Tokelau CRF Table 1.AB LPG: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels] [Liquefied Petroleum Gases (LPG)] (Part 1 of 3)	206
Tokelau CRF Table 1.AB LPG: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels] [Liquefied Petroleum Gases (LPG)] (Part 2 of 3)	206
Tokelau CRF Table 1.AB LPG: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels] [Liquefied Petroleum Gases (LPG)] (Part 3 of 3)	207
Tokelau CRF Table 1.AB Lubricants: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels] [Lubricants] (Part 1 of 3)	208
Tokelau CRF Table 1.AB Lubricants: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels] [Lubricants] (Part 2 of 3)	209
Tokelau CRF Table 1.AB Lubricants: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels] [Lubricants] (Part 3 of 3)	210

Tokelau CRF Table 2.F.1.b HFC-134a Product Uses as Substitutes for ODS: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.b Domestic Refrigeration] [HFC-134a] (Part 1 of 3)	211
Tokelau CRF Table 2.F.1.b HFC-134a Product Uses as Substitutes for ODS: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.b Domestic Refrigeration] [HFC-134a] (Part 2 of 3)	211
Tokelau CRF Table 2.F.1.b HFC-134a Product Uses as Substitutes for ODS: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.b Domestic Refrigeration] [HFC-134a] (Part 3 of 3)	212
Tokelau CRF Table 2.F.1.f: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air Conditioning] (Part 1 of 3)	213
Tokelau CRF Table 2.F.1.f: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air Conditioning] (Part 2 of 3)	213
Tokelau CRF Table 2.F.1.f: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air Conditioning] (Part 3 of 3)	214
Tokelau CRF Table 2.F.1.f HFC-32: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-32] (Part 1 of 3)	215
Tokelau CRF Table 2.F.1.f HFC-32: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-32] (Part 2 of 3)	216
Tokelau CRF Table 2.F.1.f HFC-32: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-32] (Part 3 of 3)	216
Tokelau CRF Table 2.F.1.f HFC-125: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-125] (Part 1 of 3)	217
Tokelau CRF Table 2.F.1.f HFC-125: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-125] (Part 2 of 3)	218
Tokelau CRF Table 2.F.1.f HFC-125: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-125] (Part 3 of 3)	218
Tokelau CRF Table 2.F.1.f HFC-134a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-134a] (Part 1 of 3)	219
Tokelau CRF Table 2.F.1.f HFC-134a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-134a] (Part 2 of 3)	220
Tokelau CRF Table 2.F.1.f HFC-134a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-134a] (Part 3 of 3)	220
Tokelau CRF Table 2.F.4.a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] (Part 1 of 3)	221
Tokelau CRF Table 2.F.4.a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] (Part 2 of 3)	222

Tokelau CRF Table 2.F.4.a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] (Part 3 of 3)	222
Tokelau CRF Table 2.F.4.a HFC-134a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] [HFC-134a] (Part 1 of 3)	223
Tokelau CRF Table 2.F.4.a HFC-134a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] [HFC-134a] (Part 2 of 3)	224
Tokelau CRF Table 2.F.4.a HFC-134a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] [HFC-134a] (Part 3 of 3)	224
Tokelau CRF Table 2.F.4.a HFC-227ea: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] [HFC-227ea] (Part 1 of 3)	225
Tokelau CRF Table 2.F.4.a HFC-227ea: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] [HFC-227ea] (Part 2 of 3)	225
Tokelau CRF Table 2.F.4.a HFC-227ea: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] [HFC-227ea] (Part 3 of 3)	226
Tokelau CRF Table 2.G.3.a: [2. Industrial Processes and Product Use] [2.G Other Product Manufacture and Use] [2.G.3 N2O from Product Uses] [2.G.3.a Medical Applications] (Part 1 of 3)	226
Tokelau CRF Table 2.G.3.a: [2. Industrial Processes and Product Use] [2.G Other Product Manufacture and Use] [2.G.3 N2O from Product Uses] [2.G.3.a Medical Applications] (Part 2 of 3)	227
Tokelau CRF Table 2.G.3.a: [2. Industrial Processes and Product Use] [2.G Other Product Manufacture and Use] [2.G.3 N2O from Product Uses] [2.G.3.a Medical Applications] (Part 3 of 3)	228
Tokelau CRF Table 3.A.3 Tokelau Swine: [3. Agriculture] [3.1 Livestock] [3.A Enteric Fermentation] [3.A.3 Swine] [Other (please specify)] [Tokelau_Swine] (Part 1 of 3)	228
Tokelau CRF Table 3.A.3 Tokelau Swine: [3. Agriculture] [3.1 Livestock] [3.A Enteric Fermentation] [3.A.3 Swine] [Other (please specify)] [Tokelau_Swine] (Part 2 of 3)	229
Tokelau CRF Table 3.A.3 Tokelau Swine: [3. Agriculture] [3.1 Livestock] [3.A Enteric Fermentation] [3.A.3 Swine] [Other (please specify)] [Tokelau_Swine] (Part 3 of 3)	230
Tokelau CRF Table 3.A.4 Tokelau Poultry: [3. Agriculture] [3.1 Livestock] [3.A Enteric Fermentation] [3.A.4 Other livestock] [Tokelau_Poultry] (Part 1 of 3)	230
Tokelau CRF Table 3.A.4 Tokelau Poultry: [3. Agriculture] [3.1 Livestock] [3.A Enteric Fermentation] [3.A.4 Other livestock] [Tokelau_Poultry] (Part 2 of 3)	231
Tokelau CRF Table 3.A.4 Tokelau Poultry: [3. Agriculture] [3.1 Livestock] [3.A Enteric Fermentation] [3.A.4 Other livestock] [Tokelau_Poultry] (Part 3 of 3)	232
Tokelau CRF Table 3.B.1.3 Tokelau Swine: [3. Agriculture] [3.1 Livestock] [3.B Manure Management] [3.B.1 CH4 Emissions] [3.B.1.3 Swine] [Other (please specify)] [Pigs] (Part 1 of 3)	233
Tokelau CRF Table 3.B.1.3 Tokelau Swine: [3. Agriculture] [3.1 Livestock] [3.B Manure Management] [3.B.1 CH4 Emissions] [3.B.1.3 Swine] [Other (please specify)] [Pigs] (Part 2 of 3)	233
Tokelau CRF Table 3.B.1.3 Tokelau Swine: [3. Agriculture] [3.1 Livestock] [3.B Manure Management] [3.B.1 CH4 Emissions] [3.B.1.3 Swine] [Other (please specify)] [Pigs] (Part 3 of 3)	234

Tokelau CRF Table 3.B.1.4 Tokelau Poultry: [3. Agriculture] [3.1 Livestock] [3.B Manure Management] [3.B.1 CH4 Emissions] [3.B.1.4 Other livestock] [Tokelau_Poultry] (Part 1 of 3)	235
Tokelau CRF Table 3.B.1.4 Tokelau Poultry: [3. Agriculture] [3.1 Livestock] [3.B Manure Management] [3.B.1 CH4 Emissions] [3.B.1.4 Other livestock] [Tokelau_Poultry] (Part 2 of 3)	235
Tokelau CRF Table 3.B.1.4 Tokelau Poultry: [3. Agriculture] [3.1 Livestock] [3.B Manure Management] [3.B.1 CH4 Emissions] [3.B.1.4 Other livestock] [Tokelau_Poultry] (Part 3 of 3)	236
Tokelau CRF Table 5.A: [5. Waste] [5.A Solid Waste Disposal] (Part 1 of 3)	237
Tokelau CRF Table 5.A: [5. Waste] [5.A Solid Waste Disposal] (Part 2 of 3)	238
Tokelau CRF Table 5.A: [5. Waste] [5.A Solid Waste Disposal] (Part 3 of 3)	238
Tokelau CRF Table 5.A.3: [5. Waste] [5.A Solid Waste Disposal] [5.A.3 Uncategorized Waste Disposal Sites] (Part 1 of 3)	239
Tokelau CRF Table 5.A.3: [5. Waste] [5.A Solid Waste Disposal] [5.A.3 Uncategorized Waste Disposal Sites] (Part 2 of 3)	240
Tokelau CRF Table 5.A.3: [5. Waste] [5.A Solid Waste Disposal] [5.A.3 Uncategorized Waste Disposal Sites] (Part 3 of 3)	241
Tokelau CRF Table 5.C.2.2.a: [5. Waste] [5.C Incineration and Open Burning of Waste] [5.C.2 Open Burning of Waste] [5.C.2.2 Non-biogenic] [5.C.2.2.a Municipal Solid Waste] (Part 1 of 3)	242
Tokelau CRF Table 5.C.2.2.a: [5. Waste] [5.C Incineration and Open Burning of Waste] [5.C.2 Open Burning of Waste] [5.C.2.2 Non-biogenic] [5.C.2.2.a Municipal Solid Waste] (Part 2 of 3)	242
Tokelau CRF Table 5.C.2.2.a: [5. Waste] [5.C Incineration and Open Burning of Waste] [5.C.2 Open Burning of Waste] [5.C.2.2 Non-biogenic] [5.C.2.2.a Municipal Solid Waste] (Part 3 of 3)	243
Tokelau CRF Table 5.D:[5. Waste] [5.D Wastewater Treatment and Discharge] (Part 1 of 3)	244
Tokelau CRF Table 5.D:[5. Waste] [5.D Wastewater Treatment and Discharge] (Part 2 of 3)	245
Tokelau CRF Table 5.D:[5. Waste] [5.D Wastewater Treatment and Discharge] (Part 3 of 3)	246
Tokelau CRF Table 5.D.1: [5. Waste] [5.D Wastewater Treatment and Discharge] [5.D.1 Domestic Wastewater] (Part 1 of 3)	247
Tokelau CRF Table 5.D.1: [5. Waste] [5.D Wastewater Treatment and Discharge] [5.D.1 Domestic Wastewater] (Part 2 of 3)	248
Tokelau CRF Table 5.D.1: [5. Waste] [5.D Wastewater Treatment and Discharge] [5.D.1 Domestic Wastewater] (Part 3 of 3)	249
Table A8.1.1 Direct and indirect emissions by fertiliser in 2021	250
Table A8.1.2 Use of fertilisers in New Zealand in 1990 and 2021	251
Table A8.1.3 Emissions from fertilisers in 1990 and 2021	252
Table A8.2.1 Total emissions by livestock type in 1990, 2020 and 2021	252

Figures


Figure A3.1.1	Proportion of urine nitrogen (N) applied to low (0–12 degree) slopes using a nutrient transfer model (equal proportion line shown in grey for comparison)	42
Figure A3.1.2	Proportion of urine nitrogen (N) applied to steep (more than 24 degree) slopes using a nutrient transfer model (equal proportion line shown in grey for comparison)	42
Figure A3.1.3	Proportion of land area, excretal nitrogen (N) and nitrous oxide (N ₂ O) emissions by hill slope category for sheep, beef cattle and deer farms in 2021	44
Figure A3.2.1	New Zealand’s land use mapping process for 1990 and 2008 land use maps	47
Figure A3.2.2	Identification of post-1989 forest in New Zealand	49
Figure A3.2.3	New Zealand’s land use mapping process for 2012 and 2016 land use maps	50
Figure A3.2.4	New Zealand’s identification of deforestation	53
Figure A3.2.5	Soil samples in the Soil CMS model calibration data set	75
Figure A3.2.6	Result of applying the Marcus multi-comparison test to the adopted model	78
Figure A3.2.7	Harvest area by age for pre-1990 planted forest in 2021	87
Figure A3.2.8	Harvest area by age for post-1989 planted forest in 2021	87
Figure A3.2.9	New Zealand’s planted forest inventory modelling process	94
Figure A3.2.10	Comparison of the LUCAS estimate for above-ground biomass removed on planted forest destocking and carbon stored in roundwood production from 1990 to 2021	98
Figure A3.2.11	Comparison of the planted forest age profile as estimated from LUCAS net stocked area for all planted forest (post-1989 and pre-1990) and the NEFD age profile as at 2021	99
Figure A3.2.12	Post-1989 planted forest carbon stock by age estimated from the yield table and plots measured in the 2016 to 2020 forest inventory	100
Figure A3.2.13	Pre-1990 planted forest – planted after 1990 – carbon stock by age estimated from the yield table and plots measured in the 2016 to 2020 forest inventory	101
Figure A3.2.14	Pre-1990 planted forest – planted before 1990 – carbon stock by age estimated from the yield table and plots measured in the 2016 to 2020 forest inventory	102
Figure A3.2.15	Post-1989 planted forest above-ground biomass carbon per hectare estimated from the forest inventory and LUCAS Calculation and Reporting Application model	103
Figure A3.2.16	Pre-1990 planted forest above-ground biomass carbon per hectare estimated from the forest inventory and LUCAS Calculation and Reporting Application model	104
Figure A3.2.17	New Zealand’s LUCAS data management system	126
Figure A3.2.18	New Zealand’s Geospatial System components	127
Figure A3.2.19	LUCAS Management Studio	129
Figure A3.2.20	LUCAS Gateway database	130
Figure A3.2.21	Improvements implementation quality control procedure	131
Figure A4.1	Reference and sectoral approaches to carbon dioxide by fuel type (kt CO ₂)	150
Figure A5.1.1	Pre-1990 planted forest age profile as at 2009, as reported for the original 2011 FMRL, the FMRLcorr and <i>Forest management</i>	156
Figure A5.1.2	Technical corrections to the FMRL by category	162

Figure A5.1.3	Comparison of the 2011 FMRL and total of technical corrections from 2013 to 2020	163
Figure A6.1.1	How the quality assurance and quality control processes and products align with the preparation of the inventory	170
Figure A6.1.2	Data processing, quality assurance and quality control processes applied to the inventory data from Tokelau and its integration into New Zealand's inventory	171

Annex 1: Key categories

A1.1 Methodology used for identifying key categories

The key categories in the inventory have been assessed using the Approach 1 level and trend methodologies from the *2006 IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC, 2006). The methodology applied was determined using the decision tree shown in figure A1.1.1. Because some categories in the inventory apply default uncertainty values for emission estimates, and developing country-specific uncertainty values is resource prohibitive, Approach 1 level and trend methodologies are used.



For this inventory submission, the Approach 1 level and trend assessments were applied, including and excluding the Land Use, Land-Use Change and Forestry (LULUCF) sector (IPCC, 2006). The level and trend assessments are calculated as per equations 4.1, 4.2 and 4.3 of the IPCC 2006 Guidelines (IPCC, 2006). Key categories are defined as those categories whose cumulative percentages, when summed in decreasing order of magnitude, contributed 95 per cent of the total level or trend.

A1.2 Disaggregation

The classification of categories follows the classification of the common reporting format (CRF) tables by:

- identifying categories using carbon dioxide equivalent emissions and considering each greenhouse gas from each category separately
- either including or excluding LULUCF categories at the level shown in the IPCC 2006 guidelines table 4.1 (IPCC, 2006).

The level of aggregation used for the key category analysis is similar to the default aggregation used for the key category analysis within the CRF tables, with adjustments to better reflect New Zealand's emissions profile. Specifically, a large proportion of emissions from the Energy and Agriculture sectors are disaggregated further than the key category analysis generated in the CRF tables, to allow for a more evenly proportioned analysis of categories.

A1.3 Tables 4.2 to 4.3 of the 2006 IPCC Guidelines (General Guidance and Reporting)

The following tables specify the level analyses for 2021 and 1990, and trend analyses, each including and excluding LULUCF. The tables show the categories that comprise 99 per cent of emissions for each analysis. Only the categories that comprise the top 95 per cent of emissions for the 2021 level analysis and the trend analysis are key categories. The 1990 level analysis tables are included for information only.

Table A1.3.1(a) Results of the key category level analysis for 99 per cent of the net emissions and removals for New Zealand in 2021

CRF category code	IPCC category	Gas	2021 estimate (kt CO ₂ -equivalent)	Level assessment (%)	Cumulative total (%)
3.A.1	Option A – Dairy Cattle	CH ₄	13,382.5	12.7	12.7
1.A.3.b	Transport – Road Transportation Liquid Fuels	CO ₂	12,555.0	11.9	24.7
4.A.1	Forest Land – Forest Land Remaining Forest Land	CO ₂	-12,163.5	11.6	36.3
4.G	Land Use, Land-Use Change and Forestry – Harvested Wood Products	CO ₂	-8,658.8	8.2	44.5
3.A.2	Other (please specify) – Sheep	CH ₄	8,011.9	7.6	52.1
3.A.1	Option A – Non-Dairy Cattle	CH ₄	5,944.0	5.7	57.8
4.A.2	Forest Land – Land Converted to Forest Land	CO ₂	-3,831.3	3.6	61.4
3.D.1.3	Direct N ₂ O Emissions From Managed Soils – Urine and Dung Deposited by Grazing Animals	N ₂ O	3,510.0	3.3	64.8
5.A	Waste – Solid Waste Disposal	CH ₄	2,578.2	2.5	67.2
1.A.1.a	Energy Industries – Public Electricity and Heat Production Solid Fuels	CO ₂	2,346.4	2.2	69.5

CRF category code	IPCC category	Gas	2021 estimate (kt CO ₂ -equivalent)	Level assessment (%)	Cumulative total (%)
1.A.1.a	Energy Industries – Public Electricity and Heat Production Gaseous Fuels	CO ₂	2,035.5	1.9	71.4
2.C.1	Metal Industry – Iron and Steel Production	CO ₂	1,718.1	1.6	73.0
4.C.2	Grassland – Land Converted to Grassland	CO ₂	1,485.2	1.4	74.4
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Solid Fuels	CO ₂	1,446.5	1.4	75.8
2.F.1	Product Uses as Substitutes for ODS – Refrigeration and Air conditioning	HFCs	1,401.8	1.3	77.2
3.D.1.1	Direct N ₂ O Emissions From Managed Soils – Inorganic N Fertilizers	N ₂ O	1,398.9	1.3	78.5
3.B.1.1	Option A – Dairy Cattle	CH ₄	1,341.1	1.3	79.8
1.A.4.b	Other Sectors – Residential Liquid Fuels	CO ₂	1,332.4	1.3	81.0
1.A.2.c	Manufacturing Industries and Construction – Chemicals Gaseous Fuels	CO ₂	1,267.1	1.2	82.2
1.A.4.c	Other Sectors – Agriculture/Forestry/Fishing Liquid Fuels	CO ₂	1,262.5	1.2	83.4
4.C.1	Grassland – Grassland Remaining Grassland	CO ₂	1,222.9	1.2	84.6
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Gaseous Fuels	CO ₂	1,053.3	1.0	85.6
3.D.2.1	Indirect N ₂ O Emissions From Managed Soils – Atmospheric Deposition	N ₂ O	854.7	0.8	86.4
1.A.3.a	Domestic Aviation – Jet Kerosene	CO ₂	797.2	0.8	87.2
1.A.4.a	Other Sectors – Commercial/Institutional Liquid Fuels	CO ₂	726.4	0.7	87.9
3.D.1.6	Direct N ₂ O Emissions From Managed Soils – Cultivation of Organic Soils	N ₂ O	667.4	0.6	88.5
1.A.1.b	Energy Industries – Petroleum Refining Liquid Fuels	CO ₂	617.3	0.6	89.1
3.D.2.2	Indirect N ₂ O Emissions From Managed Soils – Nitrogen Leaching and Run-off	N ₂ O	554.5	0.5	89.6
3.H	Agriculture – Urea Application	CO ₂	553.2	0.5	90.1
2.C.3	Metal Industry – Aluminium Production	CO ₂	541.8	0.5	90.7
1.A.2.g.v	Other (please specify) – Construction	CO ₂	522.6	0.5	91.2
3.A.4	Other livestock – Deer	CH ₄	468.3	0.4	91.6
1.B.2.d	Other (please specify) – Geothermal	CO ₂	431.6	0.4	92.0
1.A.4.a	Other Sectors – Commercial/Institutional Gaseous Fuels	CO ₂	412.5	0.4	92.4
1.A.4.b	Other Sectors – Residential Gaseous Fuels	CO ₂	385.5	0.4	92.8
2.A.1	Mineral Industry – Cement Production	CO ₂	369.6	0.4	93.1
1.A.2.g.iii	Other (please specify) – Mining (excluding fuels) and quarrying Liquid Fuels	CO ₂	367.6	0.3	93.5
3.G	Agriculture – Liming	CO ₂	355.6	0.3	93.8
4.B.1	Cropland – Cropland Remaining Cropland	CO ₂	320.7	0.3	94.1
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Liquid Fuels	CO ₂	287.1	0.3	94.4
5.D	Waste – Wastewater Treatment and Discharge	CH ₄	256.9	0.2	94.6
3.D.1.4	Direct N ₂ O Emissions From Managed Soils – Crop Residues	N ₂ O	255.7	0.2	94.9
1.A.1.c	Energy Industries – Manufacture of Solid Fuels and Other Energy Industries Gaseous Fuels	CO ₂	240.4	0.2	95.1
1.B.2.c.1.ii	Venting – Gas	CO ₂	234.2	0.2	95.3

CRF category code	IPCC category	Gas	2021 estimate (kt CO ₂ -equivalent)	Level assessment (%)	Cumulative total (%)
1.A.2.g.viii	Other (please specify) – Other (please specify) Liquid Fuels	CO ₂	221.0	0.2	95.5
1.A.3.d	Domestic Navigation – Residual Fuel Oil	CO ₂	201.3	0.2	95.7
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Gaseous Fuels	CO ₂	194.9	0.2	95.9
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Solid Fuels	CO ₂	187.7	0.2	96.1
1.A.4.c	Other Sectors – Agriculture/Forestry/Fishing Solid Fuels	CO ₂	139.3	0.1	96.2
1.B.2.b.2	Natural Gas – Production	CH ₄	126.8	0.1	96.4
4.A	Forest Land – Emissions and removals from drainage and rewetting and other management of organic and mineral soils	N ₂ O	124.6	0.1	96.5
5.D	Waste – Wastewater Treatment and Discharge	N ₂ O	120.7	0.1	96.6
1.A.3.c	Transport – Railways Liquid Fuels	CO ₂	117.6	0.1	96.7
1.B.2.d	Other (please specify) – Geothermal	CH ₄	114.6	0.1	96.8
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Gaseous Fuels	CO ₂	113.6	0.1	96.9
1.A.1.b	Energy Industries – Petroleum Refining Gaseous Fuels	CO ₂	111.7	0.1	97.0
1.A.2.g.viii	Other (please specify) – Other (please specify) Solid Fuels	CO ₂	104.4	0.1	97.1
2.A.2	Mineral Industry – Lime Production	CO ₂	103.7	0.1	97.2
1.A.2.a	Manufacturing Industries and Construction – Iron and Steel Gaseous Fuels	CO ₂	94.1	0.1	97.3
3.B.2.5	N ₂ O and NMVOC Emissions – Indirect N ₂ O Emissions	N ₂ O	93.6	0.1	97.4
5.C	Waste – Incineration and Open Burning of Waste	CO ₂	90.6	0.1	97.5
2.G.3	Other Product Manufacture and Use – N ₂ O from Product Uses	N ₂ O	89.5	0.1	97.6
3.B.1.2	CH ₄ Emissions – Sheep	CH ₄	87.6	0.1	97.7
1.A.3.b	Transport – Road Transportation Liquid Fuels	N ₂ O	84.3	0.1	97.7
3.B.1.1	Option A – Non-Dairy Cattle	CH ₄	82.6	0.1	97.8
4.F.2	Other Land – Land Converted to Other Land	CO ₂	81.6	0.1	97.9
5.C	Waste – Incineration and Open Burning of Waste	CH ₄	78.1	0.1	98.0
2.B.8	Chemical Industry – Petrochemical and Carbon Black Production	CH ₄	77.6	0.1	98.0
4.E.1	Settlements – Settlements Remaining Settlements	CO ₂	77.4	0.1	98.1
3.D.1.2	Direct N ₂ O Emissions From Managed Soils – Organic N Fertilizers	N ₂ O	73.8	0.1	98.2
2.F.4	Product Uses as Substitutes for ODS – Aerosols	HFCs	72.7	0.1	98.3
1.A.4.c	Other Sectors – Agriculture/Forestry/Fishing Gaseous Fuels	CO ₂	70.0	0.1	98.3
4.A.2	Forest Land – Land Converted to Forest Land	N ₂ O	65.1	0.1	98.4
1.B.1.a.2	Coal Mining and Handling – Surface Mines	CH ₄	62.4	0.1	98.4
2.A.4	Mineral Industry – Other Process Uses of Carbonates	CO ₂	55.9	0.1	98.5
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Liquid Fuels	CO ₂	53.1	0.1	98.5
4.B.2	Cropland – Land Converted to Cropland	CO ₂	51.4	0.0	98.6
2.C.3	Metal Industry – Aluminium Production	PFCs	50.7	0.0	98.6
1.A.4.b	Other Sectors – Residential Biomass	CH ₄	45.4	0.0	98.7

CRF category code	IPCC category	Gas	2021 estimate (kt CO ₂ -equivalent)	Level assessment (%)	Cumulative total (%)
1.A.4.a	Other Sectors – Commercial/Institutional Solid Fuels	CO ₂	44.8	0.0	98.7
4.E.2	Settlements – Land Converted to Settlements	CO ₂	44.8	0.0	98.8
2.B.10	Chemical Industry – Other (please specify)	CO ₂	43.4	0.0	98.8
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Liquid Fuels	CO ₂	43.2	0.0	98.9
2.D	Industrial Processes and Product Use – Non-energy Products from Fuels and Solvent Use	CO ₂	42.4	0.0	98.9
5.B	Waste – Biological Treatment of Solid Waste	CH ₄	42.0	0.0	98.9
1.A.3.e	Transport – Other Transportation (please specify) Gaseous Fuels	CO ₂	41.6	0.0	99.0
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Biomass	N ₂ O	39.4	0.0	99.0

Note: Key categories are those that comprise 95 per cent of the total. Removals from the LULUCF sector are shown as negatives in this table. The absolute values for those removals were used for the calculations.

Table A1.3.1(b) Results of the key category level analysis for 99 per cent of the gross emissions and removals for New Zealand in 2021

CRF category code	IPCC category	Gas	2021 estimate (kt CO ₂ -equivalent)	Level assessment (%)	Cumulative total (%)
3.A.1	Option A – Dairy Cattle	CH ₄	13,382.5	17.4	17.4
1.A.3.b	Transport – Road Transportation Liquid Fuels	CO ₂	12,555.0	16.3	33.8
3.A.2	Other (please specify) – Sheep	CH ₄	8,011.9	10.4	44.2
3.A.1	Option A – Non-Dairy Cattle	CH ₄	5,944.0	7.7	51.9
3.D.1.3	Direct N ₂ O Emissions From Managed Soils – Urine and Dung Deposited by Grazing Animals	N ₂ O	3,510.0	4.6	56.5
5.A	Waste – Solid Waste Disposal	CH ₄	2,578.2	3.4	59.9
1.A.1.a	Energy Industries – Public Electricity and Heat Production Solid Fuels	CO ₂	2,346.4	3.1	62.9
1.A.1.a	Energy Industries – Public Electricity and Heat Production Gaseous Fuels	CO ₂	2,035.5	2.6	65.6
2.C.1	Metal Industry – Iron and Steel Production	CO ₂	1,718.1	2.2	67.8
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Solid Fuels	CO ₂	1,446.5	1.9	69.7
2.F.1	Product Uses as Substitutes for ODS – Refrigeration and Air conditioning	HFCs	1,401.8	1.8	71.5
3.D.1.1	Direct N ₂ O Emissions From Managed Soils – Inorganic N Fertilizers	N ₂ O	1,398.9	1.8	73.3
3.B.1.1	Option A – Dairy Cattle	CH ₄	1,341.1	1.7	75.1
1.A.4.b	Other Sectors – Residential Liquid Fuels	CO ₂	1,332.4	1.7	76.8
1.A.2.c	Manufacturing Industries and Construction – Chemicals Gaseous Fuels	CO ₂	1,267.1	1.6	78.5
1.A.4.c	Other Sectors – Agriculture/Forestry/Fishing Liquid Fuels	CO ₂	1,262.5	1.6	80.1
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Gaseous Fuels	CO ₂	1,053.3	1.4	81.5
3.D.2.1	Indirect N ₂ O Emissions From Managed Soils – Atmospheric Deposition	N ₂ O	854.7	1.1	82.6
1.A.3.a	Domestic Aviation – Jet Kerosene	CO ₂	797.2	1.0	83.6
1.A.4.a	Other Sectors – Commercial/Institutional Liquid Fuels	CO ₂	726.4	0.9	84.6

CRF category code	IPCC category	Gas	2021 estimate (kt CO ₂ -equivalent)	Level assessment (%)	Cumulative total (%)
3.D.1.6	Direct N ₂ O Emissions From Managed Soils – Cultivation of Organic Soils	N ₂ O	667.4	0.9	85.4
1.A.1.b	Energy Industries – Petroleum Refining Liquid Fuels	CO ₂	617.3	0.8	86.2
3.D.2.2	Indirect N ₂ O Emissions From Managed Soils – Nitrogen Leaching and Run-off	N ₂ O	554.5	0.7	87.0
3.H	Agriculture – Urea Application	CO ₂	553.2	0.7	87.7
2.C.3	Metal Industry – Aluminium Production	CO ₂	541.8	0.7	88.4
1.A.2.g.v	Other (please specify) – Construction	CO ₂	522.6	0.7	89.1
3.A.4	Other livestock – Deer	CH ₄	468.3	0.6	89.7
1.B.2.d	Other (please specify) – Geothermal	CO ₂	431.6	0.6	90.2
1.A.4.a	Other Sectors – Commercial/Institutional Gaseous Fuels	CO ₂	412.5	0.5	90.8
1.A.4.b	Other Sectors – Residential Gaseous Fuels	CO ₂	385.5	0.5	91.3
2.A.1	Mineral Industry – Cement Production	CO ₂	369.6	0.5	91.8
1.A.2.g.iii	Other (please specify) – Mining (excluding fuels) and quarrying Liquid Fuels	CO ₂	367.6	0.5	92.2
3.G	Agriculture – Liming	CO ₂	355.6	0.5	92.7
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Liquid Fuels	CO ₂	287.1	0.4	93.1
5.D	Waste – Wastewater Treatment and Discharge	CH ₄	256.9	0.3	93.4
3.D.1.4	Direct N ₂ O Emissions From Managed Soils – Crop Residues	N ₂ O	255.7	0.3	93.7
1.A.1.c	Energy Industries – Manufacture of Solid Fuels and Other Energy Industries Gaseous Fuels	CO ₂	240.4	0.3	94.0
1.B.2.c.1.ii	Venting – Gas	CO ₂	234.2	0.3	94.4
1.A.2.g.viii	Other (please specify) – Other (please specify) Liquid Fuels	CO ₂	221.0	0.3	94.6
1.A.3.d	Domestic Navigation – Residual Fuel Oil	CO ₂	201.3	0.3	94.9
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Gaseous Fuels	CO ₂	194.9	0.3	95.2
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Solid Fuels	CO ₂	187.7	0.2	95.4
1.A.4.c	Other Sectors – Agriculture/Forestry/Fishing Solid Fuels	CO ₂	139.3	0.2	95.6
1.B.2.b.2	Natural Gas – Production	CH ₄	126.8	0.2	95.7
5.D	Waste – Wastewater Treatment and Discharge	N ₂ O	120.7	0.2	95.9
1.A.3.c	Transport – Railways Liquid Fuels	CO ₂	117.6	0.2	96.1
1.B.2.d	Other (please specify) – Geothermal	CH ₄	114.6	0.1	96.2
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Gaseous Fuels	CO ₂	113.6	0.1	96.4
1.A.1.b	Energy Industries – Petroleum Refining Gaseous Fuels	CO ₂	111.7	0.1	96.5
1.A.2.g.viii	Other (please specify) – Other (please specify) Solid Fuels	CO ₂	104.4	0.1	96.6
2.A.2	Mineral Industry – Lime Production	CO ₂	103.7	0.1	96.8
1.A.2.a	Manufacturing Industries and Construction – Iron and Steel Gaseous Fuels	CO ₂	94.1	0.1	96.9
3.B.2.5	N ₂ O and NMVOC Emissions – Indirect N ₂ O Emissions	N ₂ O	93.6	0.1	97.0
5.C	Waste – Incineration and Open Burning of Waste	CO ₂	90.6	0.1	97.1
2.G.3	Other Product Manufacture and Use – N ₂ O from Product Uses	N ₂ O	89.5	0.1	97.2
3.B.1.2	CH ₄ Emissions – Sheep	CH ₄	87.6	0.1	97.4

CRF category code	IPCC category	Gas	2021 estimate (kt CO ₂ -equivalent)	Level assessment (%)	Cumulative total (%)
1.A.3.b	Transport – Road Transportation Liquid Fuels	N ₂ O	84.3	0.1	97.5
3.B.1.1	Option A – Non-Dairy Cattle	CH ₄	82.6	0.1	97.6
5.C	Waste – Incineration and Open Burning of Waste	CH ₄	78.1	0.1	97.7
2.B.8	Chemical Industry – Petrochemical and Carbon Black Production	CH ₄	77.6	0.1	97.8
3.D.1.2	Direct N ₂ O Emissions From Managed Soils – Organic N Fertilizers	N ₂ O	73.8	0.1	97.9
2.F.4	Product Uses as Substitutes for ODS – Aerosols	HFCs	72.7	0.1	98.0
1.A.4.c	Other Sectors – Agriculture/Forestry/Fishing Gaseous Fuels	CO ₂	70.0	0.1	98.1
1.B.1.a.2	Coal Mining and Handling – Surface Mines	CH ₄	62.4	0.1	98.1
2.A.4	Mineral Industry – Other Process Uses of Carbonates	CO ₂	55.9	0.1	98.2
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Liquid Fuels	CO ₂	53.1	0.1	98.3
2.C.3	Metal Industry – Aluminium Production	PFCs	50.7	0.1	98.4
1.A.4.b	Other Sectors – Residential Biomass	CH ₄	45.4	0.1	98.4
1.A.4.a	Other Sectors – Commercial/Institutional Solid Fuels	CO ₂	44.8	0.1	98.5
2.B.10	Chemical Industry – Other (please specify)	CO ₂	43.4	0.1	98.5
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Liquid Fuels	CO ₂	43.2	0.1	98.6
2.D	Industrial Processes and Product Use – Non-energy Products from Fuels and Solvent Use	CO ₂	42.4	0.1	98.6
5.B	Waste – Biological Treatment of Solid Waste	CH ₄	42.0	0.1	98.7
1.A.3.e	Transport – Other Transportation (please specify) Gaseous Fuels	CO ₂	41.6	0.1	98.7
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Biomass	N ₂ O	39.4	0.1	98.8
1.B.2.b.5	Natural Gas – Distribution	CH ₄	39.1	0.1	98.8
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Other Fossil Fuels	CO ₂	37.6	0.0	98.9
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Other Fossil Fuels	CO ₂	37.0	0.0	98.9
3.B.1.3	CH ₄ Emissions – Swine	CH ₄	36.9	0.0	99.0
1.A.4.b	Other Sectors – Residential Liquid Fuels	N ₂ O	34.5	0.0	99.0

Note: Key categories are those that comprise 95 per cent of the total.

Table A1.3.2(a) Results of the level analysis for 99 per cent of the net emissions and removals for New Zealand in 1990 included for reference only

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	Level assessment (%)	Cumulative total (%)
4.A.2	Forest Land – Land Converted to Forest Land	CO ₂	-18,491.9	21.0	21.0
3.A.2	Other (please specify) – Sheep	CH ₄	14,407.4	16.4	37.3
1.A.3.b	Transport – Road Transportation Liquid Fuels	CO ₂	6,519.0	7.4	44.7
3.A.1	Option A – Dairy Cattle	CH ₄	6,012.9	6.8	51.6
3.A.1	Option A – Non-Dairy Cattle	CH ₄	5,861.6	6.7	58.2
5.A	Waste – Solid Waste Disposal	CH ₄	3,318.2	3.8	62.0
1.A.1.a	Energy Industries – Public Electricity and Heat Production Gaseous Fuels	CO ₂	2,999.6	3.4	65.4

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	Level assessment (%)	Cumulative total (%)
3.D.1.3	Direct N ₂ O Emissions From Managed Soils – Urine and Dung Deposited by Grazing Animals	N ₂ O	2,959.6	3.4	68.7
4.G	Land Use, Land-Use Change and Forestry – Harvested Wood Products	CO ₂	-2,481.2	2.8	71.6
1.A.1.c	Energy Industries – Manufacture of Solid Fuels and Other Energy Industries Gaseous Fuels	CO ₂	1,715.3	1.9	73.5
2.C.1	Metal Industry – Iron and Steel Production	CO ₂	1,306.7	1.5	75.0
1.A.4.c	Other Sectors – Agriculture/Forestry/Fishing Liquid Fuels	CO ₂	1,071.4	1.2	76.2
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Solid Fuels	CO ₂	938.6	1.1	77.3
2.C.3	Metal Industry – Aluminium Production	PFCs	909.9	1.0	78.3
1.A.3.a	Domestic Aviation – Jet Kerosene	CO ₂	892.6	1.0	79.3
1.A.4.b	Other Sectors – Residential Liquid Fuels	CO ₂	814.5	0.9	80.2
4.A.1	Forest Land – Forest Land Remaining Forest Land	CO ₂	-784.0	0.9	81.1
1.A.1.b	Energy Industries – Petroleum Refining Liquid Fuels	CO ₂	778.9	0.9	82.0
1.A.2.g.viii	Other (please specify) – Other (please specify) Solid Fuels	CO ₂	731.1	0.8	82.9
3.D.2.1	Indirect N ₂ O Emissions From Managed Soils – Atmospheric Deposition	N ₂ O	715.7	0.8	83.7
3.D.1.6	Direct N ₂ O Emissions From Managed Soils – Cultivation of Organic Soils	N ₂ O	658.7	0.7	84.4
1.A.2.c	Manufacturing Industries and Construction – Chemicals Gaseous Fuels	CO ₂	524.8	0.6	85.0
1.A.4.a	Other Sectors – Commercial/Institutional Liquid Fuels	CO ₂	500.7	0.6	85.6
1.A.1.a	Energy Industries – Public Electricity and Heat Production Solid Fuels	CO ₂	474.8	0.5	86.1
2.C.3	Metal Industry – Aluminium Production	CO ₂	449.0	0.5	86.6
2.A.1	Mineral Industry – Cement Production	CO ₂	448.7	0.5	87.1
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Gaseous Fuels	CO ₂	443.4	0.5	87.6
3.D.2.2	Indirect N ₂ O Emissions From Managed Soils – Nitrogen Leaching and Run-off	N ₂ O	441.2	0.5	88.1
3.A.4	Other Livestock – Deer	CH ₄	429.0	0.5	88.6
4.C.2	Grassland – Land Converted to Grassland	CO ₂	422.4	0.5	89.1
3.B.1.1	Option A – Dairy Cattle	CH ₄	413.8	0.5	89.6
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Solid Fuels	CO ₂	382.9	0.4	90.0
4.B.1	Cropland – Cropland Remaining Cropland	CO ₂	351.2	0.4	90.4
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Gaseous Fuels	CO ₂	347.6	0.4	90.8
1.A.4.b	Other Sectors – Residential Solid Fuels	CO ₂	344.9	0.4	91.2
3.G	Agriculture – Liming	CO ₂	296.5	0.3	91.5
1.B.1.a.1	Coal Mining and Handling – Underground Mines	CH ₄	289.6	0.3	91.9
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Liquid Fuels	CO ₂	281.1	0.3	92.2
1.B.2.b.5	Natural Gas – Distribution	CH ₄	277.5	0.3	92.5
1.A.2.g.v	Other (please specify) – Construction	CO ₂	245.0	0.3	92.8
1.A.4.a	Other Sectors – Commercial/Institutional Gaseous Fuels	CO ₂	235.2	0.3	93.0

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	Level assessment (%)	Cumulative total (%)
1.A.3.d	Domestic Navigation – Residual Fuel Oil	CO ₂	232.9	0.3	93.3
3.D.1.1	Direct N ₂ O Emissions From Managed Soils – Inorganic N Fertilizers	N ₂ O	230.3	0.3	93.6
1.B.2.d	Other (please specify) – Geothermal	CO ₂	228.6	0.3	93.8
5.D	Waste – Wastewater Treatment and Discharge	CH ₄	224.0	0.3	94.1
4.C.1	Grassland – Grassland Remaining Grassland	CO ₂	219.6	0.2	94.3
3.A.4	Other Livestock – Goats	CH ₄	196.6	0.2	94.5
1.A.4.b	Other Sectors – Residential Gaseous Fuels	CO ₂	184.9	0.2	94.8
3.D.1.4	Direct N ₂ O Emissions From Managed Soils – Crop Residues	N ₂ O	175.5	0.2	95.0
5.C	Waste – Incineration and Open Burning of Waste	CO ₂	158.9	0.2	95.1
2.B.10	Chemical Industry – Other (please specify)	CO ₂	152.3	0.2	95.3
3.B.1.2	CH ₄ Emissions – Sheep	CH ₄	146.8	0.2	95.5
1.B.2.b.2	Natural Gas – Production	CH ₄	143.5	0.2	95.6
1.A.4.a	Other Sectors – Commercial/Institutional Solid Fuels	CO ₂	142.2	0.2	95.8
1.A.3.b	Transport – Road Transportation Gaseous Fuels	CO ₂	140.3	0.2	96.0
5.C	Waste – Incineration and Open Burning of Waste	CH ₄	127.4	0.1	96.1
4.A.2	Forest Land – Land Converted to Forest Land	N ₂ O	124.2	0.1	96.2
4.B.2	Cropland – Land Converted to Cropland	CO ₂	117.9	0.1	96.4
1.A.2.a	Manufacturing Industries and Construction – Iron and Steel Gaseous Fuels	CO ₂	116.2	0.1	96.5
1.B.2.c.2.iii	Flaring – Combined	CO ₂	114.1	0.1	96.6
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Solid Fuels	CO ₂	109.5	0.1	96.8
1.B.2.c.1.ii	Venting – Gas	CO ₂	109.3	0.1	96.9
1.A.4.c	Other Sectors – Agriculture/Forestry/Fishing Gaseous Fuels	CO ₂	105.8	0.1	97.0
2.G.3	Other Product Manufacture and Use – N ₂ O from Product Uses	N ₂ O	102.4	0.1	97.1
1.B.2.c.1.iii	Venting – Combined	CH ₄	96.2	0.1	97.2
1.A.2.g.iii	Other (please specify) – Mining (excluding fuels) and Quarrying Liquid Fuels	CO ₂	94.1	0.1	97.3
1.A.3.b	Transport – Road Transportation Liquid Fuels	N ₂ O	89.3	0.1	97.4
2.A.2	Mineral Industry – Lime Production	CO ₂	82.6	0.1	97.5
5.D	Waste – Wastewater Treatment and Discharge	N ₂ O	82.0	0.1	97.6
3.B.1.1	Option A – Non-Dairy Cattle	CH ₄	81.1	0.1	97.7
1.A.3.c	Transport – Railways Liquid Fuels	CO ₂	78.4	0.1	97.8
4.A	Forest Land – Emissions and removals from drainage and rewetting and other management of organic and mineral soils	N ₂ O	77.7	0.1	97.9
1.A.3.b	Transport – Road Transportation Liquid Fuels	CH ₄	72.9	0.1	98.0
4.E.1	Settlements – Settlements Remaining Settlements	CO ₂	67.2	0.1	98.1
1.B.2.c.2.iii	Flaring – Combined	CH ₄	64.6	0.1	98.1
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Gaseous Fuels	CO ₂	64.1	0.1	98.2
1.A.2.g.vi	Other (please specify) – Textile and leather Gaseous Fuels	CO ₂	58.9	0.1	98.3
3.B.1.3	CH ₄ Emissions – Swine	CH ₄	58.6	0.1	98.3

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	Level assessment (%)	Cumulative total (%)
1.B.2.d	Other (please specify) – Geothermal	CH ₄	54.8	0.1	98.4
1.A.2.g.viii	Other (please specify) – Other (please specify) Liquid Fuels	CO ₂	52.3	0.1	98.5
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Liquid Fuels	CO ₂	50.1	0.1	98.5
1.A.4.b	Other Sectors – Residential Biomass	CH ₄	48.4	0.1	98.6
4.C.1	Grassland – Grassland Remaining Grassland	CH ₄	47.7	0.1	98.6
1.A.3.a	Domestic Aviation – Aviation Gasoline	CO ₂	47.7	0.1	98.7
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Liquid Fuels	CO ₂	46.0	0.1	98.7
3.A.4	Other livestock – Horses	CH ₄	42.3	0.0	98.8
1.A.2.g.i	Other (please specify) – Manufacturing of Machinery Gaseous Fuels	CO ₂	41.8	0.0	98.8
–	Land Use, Land-Use Change and Forestry – Indirect N ₂ O Emissions from Managed Soils	N ₂ O	40.8	0.0	98.9
3.H	Agriculture – Urea Application	CO ₂	39.2	0.0	98.9
1.B.1.a.2	Coal Mining and Handling – Surface Mines	CH ₄	38.5	0.0	99.0
3.D.1.2	Direct N ₂ O Emissions From Managed Soils – Organic N Fertilizers	N ₂ O	35.8	0.0	99.0

Note: Removals from the LULUCF sector are shown as negatives in this table. The absolute values for those removals were used for the calculations.

Table A1.3.2(b) Results of the level analysis for 99 per cent of the gross emissions and removals for New Zealand in 1990 included for reference only

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	Level assessment (%)	Cumulative total (%)
3.A.2	Other (please specify) – Sheep	CH ₄	14,407.4	22.3	22.3
1.A.3.b	Transport – Road Transportation Liquid Fuels	CO ₂	6,519.0	10.1	32.3
3.A.1	Option A – Dairy Cattle	CH ₄	6,012.9	9.3	41.6
3.A.1	Option A – Non-Dairy Cattle	CH ₄	5,861.6	9.1	50.7
5.A	Waste – Solid Waste Disposal	CH ₄	3,318.2	5.1	55.8
1.A.1.a	Energy Industries – Public Electricity and Heat Production Gaseous Fuels	CO ₂	2,999.6	4.6	60.4
3.D.1.3	Direct N ₂ O Emissions From Managed Soils – Urine and Dung Deposited by Grazing Animals	N ₂ O	2,959.6	4.6	65.0
1.A.1.c	Energy Industries – Manufacture of Solid Fuels and Other Energy Industries Gaseous Fuels	CO ₂	1,715.3	2.7	67.7
2.C.1	Metal Industry – Iron and Steel Production	CO ₂	1,306.7	2.0	69.7
1.A.4.c	Other Sectors – Agriculture/Forestry/Fishing Liquid Fuels	CO ₂	1,071.4	1.7	71.3
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Solid Fuels	CO ₂	938.6	1.5	72.8
2.C.3	Metal Industry – Aluminium Production	PFCs	909.9	1.4	74.2
1.A.3.a	Domestic Aviation – Jet Kerosene	CO ₂	892.6	1.4	75.6
1.A.4.b	Other Sectors – Residential Liquid Fuels	CO ₂	814.5	1.3	76.8
1.A.1.b	Energy Industries – Petroleum Refining Liquid Fuels	CO ₂	778.9	1.2	78.0
1.A.2.g.viii	Other (please specify) – Other (please specify) Solid Fuels	CO ₂	731.1	1.1	79.2
3.D.2.1	Indirect N ₂ O Emissions From Managed Soils – Atmospheric Deposition	N ₂ O	715.7	1.1	80.3

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	Level assessment (%)	Cumulative total (%)
3.D.1.6	Direct N ₂ O Emissions From Managed Soils – Cultivation of Organic Soils	N ₂ O	658.7	1.0	81.3
1.A.2.c	Manufacturing Industries and Construction – Chemicals Gaseous Fuels	CO ₂	524.8	0.8	82.1
1.A.4.a	Other Sectors – Commercial/Institutional Liquid Fuels	CO ₂	500.7	0.8	82.9
1.A.1.a	Energy Industries – Public Electricity and Heat Production Solid Fuels	CO ₂	474.8	0.7	83.6
2.C.3	Metal Industry – Aluminium Production	CO ₂	449.0	0.7	84.3
2.A.1	Mineral Industry – Cement Production	CO ₂	448.7	0.7	85.0
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Gaseous Fuels	CO ₂	443.4	0.7	85.7
3.D.2.2	Indirect N ₂ O Emissions From Managed Soils – Nitrogen Leaching and Run-off	N ₂ O	441.2	0.7	86.4
3.A.4	Other Livestock – Deer	CH ₄	429.0	0.7	87.0
3.B.1.1	Option A – Dairy Cattle	CH ₄	413.8	0.6	87.7
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Solid Fuels	CO ₂	382.9	0.6	88.3
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Gaseous Fuels	CO ₂	347.6	0.5	88.8
1.A.4.b	Other Sectors – Residential Solid Fuels	CO ₂	344.9	0.5	89.3
3.G	Agriculture – Liming	CO ₂	296.5	0.5	89.8
1.B.1.a.1	Coal Mining and Handling – Underground Mines	CH ₄	289.6	0.4	90.2
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Liquid Fuels	CO ₂	281.1	0.4	90.7
1.B.2.b.5	Natural Gas – Distribution	CH ₄	277.5	0.4	91.1
1.A.2.g.v	Other (please specify) – Construction	CO ₂	245.0	0.4	91.5
1.A.4.a	Other Sectors – Commercial/Institutional Gaseous Fuels	CO ₂	235.2	0.4	91.8
1.A.3.d	Domestic Navigation – Residual Fuel Oil	CO ₂	232.9	0.4	92.2
3.D.1.1	Direct N ₂ O Emissions From Managed Soils – Inorganic N Fertilizers	N ₂ O	230.3	0.4	92.6
1.B.2.d	Other (please specify) – Geothermal	CO ₂	228.6	0.4	92.9
5.D	Waste – Wastewater Treatment and Discharge	CH ₄	224.0	0.3	93.3
3.A.4	Other Livestock – Goats	CH ₄	196.6	0.3	93.6
1.A.4.b	Other Sectors – Residential Gaseous Fuels	CO ₂	184.9	0.3	93.8
3.D.1.4	Direct N ₂ O Emissions From Managed Soils – Crop Residues	N ₂ O	175.5	0.3	94.1
5.C	Waste – Incineration and Open Burning of Waste	CO ₂	158.9	0.2	94.4
2.B.10	Chemical Industry – Other (please specify)	CO ₂	152.3	0.2	94.6
3.B.1.2	CH ₄ Emissions – Sheep	CH ₄	146.8	0.2	94.8
1.B.2.b.2	Natural Gas – Production	CH ₄	143.5	0.2	95.0
1.A.4.a	Other Sectors – Commercial/Institutional Solid Fuels	CO ₂	142.2	0.2	95.3
1.A.3.b	Transport – Road Transportation Gaseous Fuels	CO ₂	140.3	0.2	95.5
5.C	Waste – Incineration and Open Burning of Waste	CH ₄	127.4	0.2	95.7
1.A.2.a	Manufacturing Industries and Construction – Iron and Steel Gaseous Fuels	CO ₂	116.2	0.2	95.9
1.B.2.c.2.iii	Flaring – Combined	CO ₂	114.1	0.2	96.0
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Solid Fuels	CO ₂	109.5	0.2	96.2
1.B.2.c.1.ii	Venting – Gas	CO ₂	109.3	0.2	96.4

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	Level assessment (%)	Cumulative total (%)
1.A.4.c	Other Sectors – Agriculture/Forestry/Fishing Gaseous Fuels	CO ₂	105.8	0.2	96.5
2.G.3	Other Product Manufacture and Use – N ₂ O from Product Uses	N ₂ O	102.4	0.2	96.7
1.B.2.c.1.iii	Venting – Combined	CH ₄	96.2	0.1	96.8
1.A.2.g.iii	Other (please specify) – Mining (excluding fuels) and Quarrying Liquid Fuels	CO ₂	94.1	0.1	97.0
1.A.3.b	Transport – Road Transportation Liquid Fuels	N ₂ O	89.3	0.1	97.1
2.A.2	Mineral Industry – Lime Production	CO ₂	82.6	0.1	97.3
5.D	Waste – Wastewater Treatment and Discharge	N ₂ O	82.0	0.1	97.4
3.B.1.1	Option A – Non-Dairy Cattle	CH ₄	81.1	0.1	97.5
1.A.3.c	Transport – Railways Liquid Fuels	CO ₂	78.4	0.1	97.6
1.A.3.b	Transport – Road Transportation Liquid Fuels	CH ₄	72.9	0.1	97.7
1.B.2.c.2.iii	Flaring – Combined	CH ₄	64.6	0.1	97.8
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Gaseous Fuels	CO ₂	64.1	0.1	97.9
1.A.2.g.vi	Other (please specify) – Textile and Leather Gaseous Fuels	CO ₂	58.9	0.1	98.0
3.B.1.3	CH ₄ Emissions – Swine	CH ₄	58.6	0.1	98.1
1.B.2.d	Other (please specify) – Geothermal	CH ₄	54.8	0.1	98.2
1.A.2.g.viii	Other (please specify) – Other (please specify) Liquid Fuels	CO ₂	52.3	0.1	98.3
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Liquid Fuels	CO ₂	50.1	0.1	98.4
1.A.4.b	Other Sectors – Residential Biomass	CH ₄	48.4	0.1	98.4
1.A.3.a	Domestic Aviation – Aviation Gasoline	CO ₂	47.7	0.1	98.5
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Liquid Fuels	CO ₂	46.0	0.1	98.6
3.A.4	Other Livestock – Horses	CH ₄	42.3	0.1	98.6
1.A.2.g.i	Other (please specify) – Manufacturing of Machinery Gaseous Fuels	CO ₂	41.8	0.1	98.7
3.H	Agriculture – Urea Application	CO ₂	39.2	0.1	98.8
1.B.1.a.2	Coal Mining and Handling – Surface Mines	CH ₄	38.5	0.1	98.8
3.D.1.2	Direct N ₂ O Emissions From Managed Soils – Organic N Fertilizers	N ₂ O	35.8	0.1	98.9
1.A.4.c	Other Sectors – Agriculture/Forestry/Fishing Solid Fuels	CO ₂	35.1	0.1	98.9
3.B.2.5	N ₂ O and NMVOC Emissions – Indirect N ₂ O Emissions	N ₂ O	33.7	0.1	99.0
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Biomass	N ₂ O	33.3	0.1	99.0

Table A1.3.3(a) Results of the key category trend analysis for 99 per cent of the net emissions and removals for New Zealand in 1990 to 2021

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	2021 estimate (kt CO ₂ -equivalent)	Trend assessment	Absolute contribution to trend (%)	Absolute cumulative total (%)
4.A.1	Forest Land – Forest Land Remaining Forest Land	CO ₂	-784.0	-12,163.5	0.131	15.5	15.5
3.A.2	Other (please specify) – Sheep	CH ₄	14,407.4	8,011.9	0.114	13.4	28.9
4.A.2	Forest Land – Land Converted to Forest Land	CO ₂	-18,491.9	-3,831.3	0.114	13.4	42.4

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	2021 estimate (kt CO ₂ -equivalent)	Trend assessment	Absolute contribution to trend (%)	Absolute cumulative total (%)
4.G	Land Use, Land-Use Change and Forestry – Harvested Wood Products	CO ₂	-2,481.2	-8,658.8	0.077	9.1	51.5
3.A.1	Option A – Dairy Cattle	CH ₄	6,012.9	13,382.5	0.066	7.9	59.3
1.A.3.b	Transport – Road Transportation Liquid Fuels	CO ₂	6,519.0	12,555.0	0.050	5.9	65.2
1.A.1.c	Energy Industries – Manufacture of Solid Fuels and Other Energy Industries Gaseous Fuels	CO ₂	1,715.3	240.4	0.022	2.6	67.8
1.A.1.a	Energy Industries – Public Electricity and Heat Production Solid Fuels	CO ₂	474.8	2,346.4	0.020	2.3	70.1
1.A.1.a	Energy Industries – Public Electricity and Heat Production Gaseous Fuels	CO ₂	2,999.6	2,035.5	0.020	2.3	72.4
5.A	Waste – Solid Waste Disposal	CH ₄	3,318.2	2,578.2	0.018	2.1	74.6
2.F.1	Product Uses as Substitutes for ODS – Refrigeration and Air conditioning	HFCs	0.0	1,401.8	0.016	1.9	76.4
3.A.1	Option A – Non-Dairy Cattle	CH ₄	5,861.6	5,944.0	0.016	1.9	78.3
3.D.1.1	Direct N ₂ O Emissions From Managed Soils – Inorganic N Fertilizers	N ₂ O	230.3	1,398.9	0.013	1.5	79.8
2.C.3	Metal Industry – Aluminium Production	PFCs	909.9	50.7	0.012	1.5	81.2
4.C.2	Grassland – Land Converted to Grassland	CO ₂	422.4	1,485.2	0.011	1.3	82.5
4.C.1	Grassland – Grassland Remaining Grassland	CO ₂	219.6	1,222.9	0.011	1.3	83.8
3.B.1.1	Option A – Dairy Cattle	CH ₄	413.8	1,341.1	0.009	1.1	84.9
1.A.2.g.viii	Other (please specify) – Other (please specify) Solid Fuels	CO ₂	731.1	104.4	0.009	1.1	86.0
1.A.2.c	Manufacturing Industries and Construction – Chemicals Gaseous Fuels	CO ₂	524.8	1,267.1	0.007	0.8	86.8
3.H	Agriculture – Urea Application	CO ₂	39.2	553.2	0.006	0.7	87.5
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Gaseous Fuels	CO ₂	443.4	1,053.3	0.006	0.7	88.1
1.A.4.b	Other Sectors – Residential Solid Fuels	CO ₂	344.9	21.8	0.005	0.5	88.7
1.B.1.a.1	Coal Mining and Handling – Underground Mines	CH ₄	289.6	0.0	0.004	0.5	89.2
1.A.1.b	Energy Industries – Petroleum Refining Liquid Fuels	CO ₂	778.9	617.3	0.004	0.5	89.7
1.A.3.a	Domestic Aviation – Jet Kerosene	CO ₂	892.6	797.2	0.004	0.4	90.1
1.A.4.b	Other Sectors – Residential Liquid Fuels	CO ₂	814.5	1,332.4	0.004	0.4	90.5
1.B.2.b.5	Natural Gas – Distribution	CH ₄	277.5	39.1	0.003	0.4	90.9

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	2021 estimate (kt CO ₂ -equivalent)	Trend assessment	Absolute contribution to trend (%)	Absolute cumulative total (%)
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Solid Fuels	CO ₂	382.9	187.7	0.003	0.4	91.3
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Solid Fuels	CO ₂	938.6	1,446.5	0.003	0.4	91.7
1.A.2.g.iii	Other (please specify) – Mining (excluding fuels) and Quarrying Liquid Fuels	CO ₂	94.1	367.6	0.003	0.3	92.0
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Gaseous Fuels	CO ₂	347.6	194.9	0.003	0.3	92.3
3.A.4	Other Livestock – Goats	CH ₄	196.6	26.1	0.002	0.3	92.6
1.A.2.g.v	Other (please specify) – Construction	CO ₂	245.0	522.6	0.002	0.3	92.9
3.D.1.3	Direct N ₂ O Emissions From Managed Soils – Urine and Dung Deposited by Grazing Animals	N ₂ O	2,959.6	3,510.0	0.002	0.3	93.2
2.A.1	Mineral Industry – Cement Production	CO ₂	448.7	369.6	0.002	0.3	93.4
1.A.3.b	Transport – Road Transportation Gaseous Fuels	CO ₂	140.3	0.0	0.002	0.2	93.7
3.D.1.6	Direct N ₂ O Emissions From Managed Soils – Cultivation of Organic Soils	N ₂ O	658.7	667.4	0.002	0.2	93.9
1.A.2.g.viii	Other (please specify) – Other (please specify) Liquid Fuels	CO ₂	52.3	221.0	0.002	0.2	94.1
1.A.4.b	Other Sectors – Residential Gaseous Fuels	CO ₂	184.9	385.5	0.002	0.2	94.3
2.B.10	Chemical Industry – Other (please specify)	CO ₂	152.3	43.4	0.002	0.2	94.5
1.B.2.d	Other (please specify) – Geothermal	CO ₂	228.6	431.6	0.002	0.2	94.7
1.A.4.a	Other Sectors – Commercial/Institutional Solid Fuels	CO ₂	142.2	44.8	0.002	0.2	94.9
4.B.1	Cropland – Cropland Remaining Cropland	CO ₂	351.2	320.7	0.001	0.2	95.0
1.A.4.a	Other Sectors – Commercial/Institutional Gaseous Fuels	CO ₂	235.2	412.5	0.001	0.2	95.2
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Solid Fuels	CO ₂	109.5	24.8	0.001	0.2	95.3
1.A.1.b	Energy Industries – Petroleum Refining Gaseous Fuels	CO ₂	0.0	111.7	0.001	0.1	95.5
1.B.2.c.2.iii	Flaring – Combined	CO ₂	114.1	34.3	0.001	0.1	95.6
5.C	Waste – Incineration and Open Burning of Waste	CO ₂	158.9	90.6	0.001	0.1	95.8
1.A.4.a	Other Sectors – Commercial/Institutional Liquid Fuels	CO ₂	500.7	726.4	0.001	0.1	95.9
1.B.2.c.1.ii	Venting – Gas	CO ₂	109.3	234.2	0.001	0.1	96.0
3.B.1.2	CH ₄ Emissions – Sheep	CH ₄	146.8	87.6	0.001	0.1	96.2

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	2021 estimate (kt CO ₂ -equivalent)	Trend assessment	Absolute contribution to trend (%)	Absolute cumulative total (%)
4.B.2	Cropland – Land Converted to Cropland	CO ₂	117.9	51.4	0.001	0.1	96.3
1.A.4.c	Other Sectors – Agriculture/ Forestry/Fishing Solid Fuels	CO ₂	35.1	139.3	0.001	0.1	96.4
4.A.2	Forest Land – Land Converted to Forest Land	N ₂ O	124.2	65.1	0.001	0.1	96.5
1.A.3.d	Domestic Navigation – Residual Fuel Oil	CO ₂	232.9	201.3	0.001	0.1	96.7
1.B.2.c.1.iii	Venting – Combined	CH ₄	96.2	31.4	0.001	0.1	96.8
2.C.1	Metal Industry – Iron and Steel Production	CO ₂	1,306.7	1,718.1	0.001	0.1	96.9
5.C	Waste – Incineration and Open Burning of Waste	CH ₄	127.4	78.1	0.001	0.1	97.0
1.A.4.c	Other Sectors – Agriculture/ Forestry/Fishing Liquid Fuels	CO ₂	1,071.4	1,262.5	0.001	0.1	97.1
1.A.3.b	Transport – Road Transportation Liquid Fuels	CH ₄	72.9	15.3	0.001	0.1	97.2
2.F.4	Product Uses as Substitutes for ODS – Aerosols	HFCs	0.0	72.7	0.001	0.1	97.3
1.B.2.c.2.iii	Flaring – Combined	CH ₄	64.6	11.2	0.001	0.1	97.4
3.A.4	Other Livestock – Deer	CH ₄	429.0	468.3	0.001	0.1	97.5
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Liquid Fuels	CO ₂	281.1	287.1	0.001	0.1	97.6
4.F.2	Other Land – Land Converted to Other Land	CO ₂	13.7	81.6	0.001	0.1	97.7
1.A.4.c	Other Sectors – Agriculture/ Forestry/Fishing Gaseous Fuels	CO ₂	105.8	70.0	0.001	0.1	97.8
1.A.2.g.vi	Other (please specify) – Textile and leather Gaseous Fuels	CO ₂	58.9	20.9	0.001	0.1	97.8
1.B.2.b.2	Natural Gas – Production	CH ₄	143.5	126.8	0.001	0.1	97.9
3.B.2.5	N ₂ O and NMVOC Emissions – Indirect N ₂ O Emissions	N ₂ O	33.7	93.6	0.001	0.1	98.0
1.A.2.a	Manufacturing Industries and Construction – Iron and Steel Gaseous Fuels	CO ₂	116.2	94.1	0.001	0.1	98.0
4.C.1	Grassland – Grassland Remaining Grassland	CH ₄	47.7	9.4	0.001	0.1	98.1
1.B.2.d	Other (please specify) – Geothermal	CH ₄	54.8	114.6	0.001	0.1	98.2
2.B.8	Chemical Industry – Petrochemical and Carbon Black Production	CH ₄	27.6	77.6	0.000	0.1	98.2
3.D.2.1	Indirect N ₂ O Emissions From Managed Soils – Atmospheric Deposition	N ₂ O	715.7	854.7	0.000	0.1	98.3
1.A.2.g.i	Other (please specify) – Manufacturing of machinery Gaseous Fuels	CO ₂	41.8	11.7	0.000	0.1	98.3
1.A.3.a	Domestic Aviation – Aviation Gasoline	CO ₂	47.7	20.4	0.000	0.1	98.4

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	2021 estimate (kt CO ₂ -equivalent)	Trend assessment	Absolute contribution to trend (%)	Absolute cumulative total (%)
2.G.3	Other Product Manufacture and Use – N ₂ O from Product Uses	N ₂ O	102.4	89.5	0.000	0.1	98.4
5.B	Waste – Biological Treatment of Solid Waste	CH ₄	2.7	42.0	0.000	0.1	98.5
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Other Fossil Fuels	CO ₂	0.0	37.6	0.000	0.1	98.5
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Other Fossil Fuels	CO ₂	0.0	37.0	0.000	0.0	98.6
3.B.1.3	CH ₄ Emissions – Swine	CH ₄	58.6	36.9	0.000	0.0	98.6
3.D.1.4	Direct N ₂ O Emissions From Managed Soils – Crop Residues	N ₂ O	175.5	255.7	0.000	0.0	98.7
3.A.4	Other Livestock – Horses	CH ₄	42.3	18.1	0.000	0.0	98.7
1.A.3.e	Transport – Other Transportation (please specify) Gaseous Fuels	CO ₂	5.5	41.6	0.000	0.0	98.8
4.E.2	Settlements – Land Converted to Settlements	CO ₂	8.5	44.8	0.000	0.0	98.8
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Gaseous Fuels	CO ₂	64.1	113.6	0.000	0.0	98.9
–	Land Use, Land-Use Change and Forestry – Indirect N ₂ O Emissions from Managed Soils	N ₂ O	40.8	18.5	0.000	0.0	98.9
1.A.4.b	Other Sectors – Residential Solid Fuels	CH ₄	27.3	1.7	0.000	0.0	99.0
4.C.2	Grassland – Land Converted to Grassland	N ₂ O	28.6	5.5	0.000	0.0	99.0
4.C.1	Grassland – Grassland Remaining Grassland	N ₂ O	35.4	15.1	0.000	0.0	99.0

Note: Key categories are those that comprise 95 per cent of the total. Removals from the LULUCF sector are shown as negatives in this table. The absolute values for those removals were used for the calculations.

Table A1.3.3(b) Results of the key category trend analysis for 99 per cent of the gross emissions and removals for New Zealand in 1990–2021

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	2021 estimate (kt CO ₂ -equivalent)	Trend assessment	Absolute contribution to trend (%)	Absolute cumulative total (%)
3.A.2	Other (please specify) – Sheep	CH ₄	14,407.4	8,011.9	0.140	21.5	21.5
3.A.1	Option A – Dairy Cattle	CH ₄	6,012.9	13,382.5	0.096	14.8	36.3
1.A.3.b	Transport – Road Transportation Liquid Fuels	CO ₂	6,519.0	12,555.0	0.074	11.4	47.7
1.A.1.c	Energy Industries – Manufacture of Solid Fuels and Other Energy Industries Gaseous Fuels	CO ₂	1,715.3	240.4	0.028	4.3	52.0
1.A.1.a	Energy Industries – Public Electricity and Heat Production Solid Fuels	CO ₂	474.8	2,346.4	0.028	4.2	56.2

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	2021 estimate (kt CO ₂ -equivalent)	Trend assessment	Absolute contribution to trend (%)	Absolute cumulative total (%)
1.A.1.a	Energy Industries – Public Electricity and Heat Production Gaseous Fuels	CO ₂	2,999.6	2,035.5	0.024	3.6	59.8
2.F.1	Product Uses as Substitutes for ODS – Refrigeration and Air Conditioning	HFCs	0.0	1,401.8	0.022	3.3	63.1
5.A	Waste – Solid Waste Disposal	CH ₄	3,318.2	2,578.2	0.021	3.2	66.4
3.D.1.1	Direct N ₂ O Emissions From Managed Soils – Inorganic N Fertilizers	N ₂ O	230.3	1,398.9	0.017	2.7	69.0
2.C.3	Metal Industry – Aluminium Production	PFCs	909.9	50.7	0.016	2.4	71.5
3.A.1	Option A – Non-Dairy Cattle	CH ₄	5,861.6	5,944.0	0.016	2.4	73.9
3.B.1.1	Option A – Dairy Cattle	CH ₄	413.8	1,341.1	0.013	2.0	75.9
1.A.2.g.viii	Other (please specify) – Other (please specify) Solid Fuels	CO ₂	731.1	104.4	0.012	1.8	77.7
1.A.2.c	Manufacturing Industries and Construction – Chemicals Gaseous Fuels	CO ₂	524.8	1,267.1	0.010	1.5	79.2
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Gaseous Fuels	CO ₂	443.4	1,053.3	0.008	1.2	80.5
3.H	Agriculture – Urea Application	CO ₂	39.2	553.2	0.008	1.2	81.7
1.A.4.b	Other Sectors – Residential Solid Fuels	CO ₂	344.9	21.8	0.006	0.9	82.6
1.A.4.b	Other Sectors – Residential Liquid Fuels	CO ₂	814.5	1,332.4	0.006	0.9	83.5
1.B.1.a.1	Coal Mining and Handling – Underground Mines	CH ₄	289.6	0.0	0.005	0.8	84.3
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Solid Fuels	CO ₂	938.6	1,446.5	0.005	0.8	85.1
1.A.1.b	Energy Industries – Petroleum Refining Liquid Fuels	CO ₂	778.9	617.3	0.005	0.7	85.8
1.B.2.b.5	Natural Gas – Distribution	CH ₄	277.5	39.1	0.004	0.7	86.5
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Solid Fuels	CO ₂	382.9	187.7	0.004	0.6	87.1
1.A.3.a	Domestic Aviation – Jet Kerosene	CO ₂	892.6	797.2	0.004	0.6	87.7
1.A.2.g.iii	Other (please specify) – Mining (excluding fuels) and Quarrying Liquid Fuels	CO ₂	94.1	367.6	0.004	0.6	88.3
1.A.2.g.v	Other (please specify) – Construction	CO ₂	245.0	522.6	0.004	0.5	88.9
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Gaseous Fuels	CO ₂	347.6	194.9	0.003	0.5	89.4
3.A.4	Other Livestock – Goats	CH ₄	196.6	26.1	0.003	0.5	89.9

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	2021 estimate (kt CO ₂ -equivalent)	Trend assessment	Absolute contribution to trend (%)	Absolute cumulative total (%)
2.C.1	Metal Industry – Iron and Steel Production	CO ₂	1,306.7	1,718.1	0.003	0.4	90.3
1.A.3.b	Transport – Road Transportation Gaseous Fuels	CO ₂	140.3	0.0	0.003	0.4	90.7
1.A.4.b	Other Sectors – Residential Gaseous Fuels	CO ₂	184.9	385.5	0.003	0.4	91.1
2.A.1	Mineral Industry – Cement Production	CO ₂	448.7	369.6	0.003	0.4	91.5
1.B.2.d	Other (please specify) – Geothermal	CO ₂	228.6	431.6	0.002	0.4	91.8
1.A.2.g.viii	Other (please specify) – Other (please specify) Liquid Fuels	CO ₂	52.3	221.0	0.002	0.4	92.2
2.B.10	Chemical Industry – Other (please specify)	CO ₂	152.3	43.4	0.002	0.3	92.5
1.A.4.a	Other Sectors – Commercial/Institutional Gaseous Fuels	CO ₂	235.2	412.5	0.002	0.3	92.8
1.A.4.a	Other Sectors – Commercial/Institutional Liquid Fuels	CO ₂	500.7	726.4	0.002	0.3	93.2
1.A.4.a	Other Sectors – Commercial/Institutional Solid Fuels	CO ₂	142.2	44.8	0.002	0.3	93.5
3.D.1.6	Direct N ₂ O Emissions From Managed Soils – Cultivation of Organic Soils	N ₂ O	658.7	667.4	0.002	0.3	93.7
1.A.1.b	Energy Industries – Petroleum Refining Gaseous Fuels	CO ₂	0.0	111.7	0.002	0.3	94.0
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Solid Fuels	CO ₂	109.5	24.8	0.002	0.2	94.2
1.B.2.c.1.ii	Venting – Gas	CO ₂	109.3	234.2	0.002	0.2	94.5
1.B.2.c.2.iii	Flaring – Combined	CO ₂	114.1	34.3	0.002	0.2	94.7
5.C	Waste – Incineration and Open Burning of Waste	CO ₂	158.9	90.6	0.002	0.2	95.0
1.A.4.c	Other Sectors – Agriculture/Forestry/Fishing Solid Fuels	CO ₂	35.1	139.3	0.002	0.2	95.2
3.B.1.2	CH ₄ Emissions – Sheep	CH ₄	146.8	87.6	0.001	0.2	95.4
1.B.2.c.1.iii	Venting – Combined	CH ₄	96.2	31.4	0.001	0.2	95.6
1.A.3.d	Domestic Navigation – Residual Fuel Oil	CO ₂	232.9	201.3	0.001	0.2	95.8
5.C	Waste – Incineration and Open Burning of Waste	CH ₄	127.4	78.1	0.001	0.2	95.9
2.F.4	Product Uses as Substitutes for ODS – Aerosols	HFCs	0.0	72.7	0.001	0.2	96.1
1.A.3.b	Transport – Road Transportation Liquid Fuels	CH ₄	72.9	15.3	0.001	0.2	96.3
1.B.2.c.2.iii	Flaring – Combined	CH ₄	64.6	11.2	0.001	0.2	96.4
1.A.4.c	Other Sectors – Agriculture/Forestry/Fishing Gaseous Fuels	CO ₂	105.8	70.0	0.001	0.1	96.6
3.B.2.5	N ₂ O and NMVOC Emissions – Indirect N ₂ O Emissions	N ₂ O	33.7	93.6	0.001	0.1	96.7
1.B.2.d	Other (please specify) – Geothermal	CH ₄	54.8	114.6	0.001	0.1	96.8

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	2021 estimate (kt CO ₂ -equivalent)	Trend assessment	Absolute contribution to trend (%)	Absolute cumulative total (%)
1.A.2.g.vi	Other (please specify) – Textile and Leather Gaseous Fuels	CO ₂	58.9	20.9	0.001	0.1	96.9
3.D.1.4	Direct N ₂ O Emissions From Managed Soils – Crop Residues	N ₂ O	175.5	255.7	0.001	0.1	97.0
1.A.2.e	Manufacturing Industries and Construction – Food Processing, Beverages and Tobacco Liquid Fuels	CO ₂	281.1	287.1	0.001	0.1	97.2
2.B.8	Chemical Industry – Petrochemical and Carbon Black Production	CH ₄	27.6	77.6	0.001	0.1	97.3
1.A.2.a	Manufacturing Industries and Construction – Iron and Steel Gaseous Fuels	CO ₂	116.2	94.1	0.001	0.1	97.4
1.B.2.b.2	Natural Gas – Production	CH ₄	143.5	126.8	0.001	0.1	97.5
3.A.4	Other Livestock – Deer	CH ₄	429.0	468.3	0.001	0.1	97.6
5.B	Waste – Biological Treatment of Solid Waste	CH ₄	2.7	42.0	0.001	0.1	97.7
1.A.2.g.i	Other (please specify) – Manufacturing of Machinery Gaseous Fuels	CO ₂	41.8	11.7	0.001	0.1	97.7
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Other Fossil Fuels	CO ₂	0.0	37.6	0.001	0.1	97.8
1.A.2.f	Manufacturing Industries and Construction – Non-metallic Minerals Gaseous Fuels	CO ₂	64.1	113.6	0.001	0.1	97.9
1.A.2.d	Manufacturing Industries and Construction – Pulp, Paper and Print Other Fossil Fuels	CO ₂	0.0	37.0	0.001	0.1	98.0
1.A.3.a	Domestic Aviation – Aviation Gasoline	CO ₂	47.7	20.4	0.001	0.1	98.1
1.A.3.e	Transport – Other Transportation (please specify) Gaseous Fuels	CO ₂	5.5	41.6	0.001	0.1	98.2
3.B.1.3	CH ₄ Emissions – Swine	CH ₄	58.6	36.9	0.001	0.1	98.3
3.A.4	Other Livestock – Horses	CH ₄	42.3	18.1	0.000	0.1	98.3
2.G.3	Other Product Manufacture and Use – N ₂ O from Product Uses	N ₂ O	102.4	89.5	0.000	0.1	98.4
3.D.1.2	Direct N ₂ O Emissions From Managed Soils – Organic N Fertilizers	N ₂ O	35.8	73.8	0.000	0.1	98.5
1.A.4.b	Other Sectors – Residential Solid Fuels	CH ₄	27.3	1.7	0.000	0.1	98.6
3.D.2.2	Indirect N ₂ O Emissions From Managed Soils – Nitrogen Leaching and Run-off	N ₂ O	441.2	554.5	0.000	0.1	98.6
5.B	Waste – Biological Treatment of Solid Waste	N ₂ O	2.0	30.1	0.000	0.1	98.7
1.A.3.c	Transport – Railways Liquid Fuels	CO ₂	78.4	117.6	0.000	0.1	98.8

CRF category code	IPCC category	Gas	1990 estimate (kt CO ₂ -equivalent)	2021 estimate (kt CO ₂ -equivalent)	Trend assessment	Absolute contribution to trend (%)	Absolute cumulative total (%)
5.D	Waste – Wastewater Treatment and Discharge	N ₂ O	82.0	120.7	0.000	0.1	98.8
1.A.2.g.vi	Other (please specify) – Textile and Leather Liquid Fuels	CO ₂	19.3	1.1	0.000	0.1	98.9
1.A.3.b	Transport – Road Transportation Liquid Fuels	N ₂ O	89.3	84.3	0.000	0.1	98.9
2.A.4	Mineral Industry – Other Process Uses of Carbonates	CO ₂	30.5	55.9	0.000	0.0	99.0
1.B.1.a.2	Coal Mining and Handling – Surface Mines	CH ₄	38.5	62.4	0.000	0.0	99.0
5.C	Waste – Incineration and Open Burning of Waste	N ₂ O	29.5	18.3	0.000	0.0	99.0

Note: Key categories are those that comprise 95 per cent of the total.

Annex 1: References

IPCC. 2006. Eggleston HS, Buendia L, Miwa K, Ngara T, Tanabe K (eds). *2006 IPCC Guidelines for National Greenhouse Gas Inventories. Volume 1. General Guidance and Reporting*. IPCC National Greenhouse Gas Inventories Programme. Japan: Published for the IPCC by the Institute for Global Environmental Strategies.

Annex 2: Uncertainty analysis

Uncertainty estimates are an essential element of a complete greenhouse gas inventory. Uncertainty information helps prioritise efforts to improve the accuracy of inventories in the future and guides decisions on methodological choice.

New Zealand has followed Approach 1 for uncertainty analysis, as required by the inventory reporting guidelines under United Nations Framework Convention on Climate Change (UNFCCC, 2013) and Intergovernmental Panel on Climate Change (IPCC) methodological guidelines (IPCC, 2006). Uncertainties in the categories are combined in the uncertainty analysis to provide uncertainty estimates for the entire inventory in any year and the uncertainty in the overall inventory trend over time. Uncertainties for the categories themselves are described in the sector chapters 3 to 8 and chapter 11, as well as chapter 1 section 1.6.

A2.1 Approach 1 uncertainty calculation

The uncertainty in activity data and emission and/or removal factors presented in tables A2.1.1 and A2.1.2 are equal to half the 95 per cent confidence interval divided by the mean and expressed as a percentage. The reason for halving the 95 per cent confidence interval is that the value corresponds to the familiar plus or minus value when uncertainties are loosely quoted as 'plus or minus x per cent'.

Where uncertainty is highly asymmetrical, the larger percentage difference between the mean and the confidence limit is entered. Where only the total uncertainty is known for a category, then:

- if uncertainty is correlated across years, the uncertainty is entered as the emission or the removal factor uncertainty and as zero in the activity data uncertainty
- if uncertainty is not correlated across years, the uncertainty is entered as the uncertainty in the activity data and as zero in the emission or the removal factor uncertainty.

In Approach 1, uncertainties in the trend are estimated using two sensitivities.

- Type A sensitivity is the change in the difference of total emissions between the base year and the current year, expressed as a percentage. Further, this change results from a 1 per cent increase in emissions of a given source category and a greenhouse gas in both the base year and the current year.
- Type B sensitivity is the change in the difference of total emissions between the base year and the current year, expressed as a percentage. Further, this change results from a 1 per cent increase in emissions of a given source category and gas in the current year only.

Uncertainties that are fully correlated between years are associated with Type A sensitivities, and uncertainties that are not correlated between years are associated with Type B sensitivities. Once the uncertainties introduced into the national inventory by Type A and Type B sensitivities have been calculated, they are summed using equation 3.1 (IPCC, 2006) to give the overall uncertainty in the trend.

In tables A2.1.1 and A2.1.2, the columns presenting trend uncertainties provide an estimate of the total uncertainty in the trend in emissions since the base year. This is expressed as the number of percentage points in the 95 per cent confidence interval in the per cent change in emissions since the base year. The values for individual categories are an estimate of the uncertainty introduced into the trend by the category in question.

In 2021 an internal review of the methods used to calculate the uncertainties was undertaken. The review identified an anomaly in the application of the methodology applied to the Land Use, Land-Use Change and Forestry (LULUCF) categories that comprised net removals, where they were converted to absolute values. This method resulted in reduced percentage uncertainty estimates for the base year, final year and trend, when compared with using unmodified values. The calculation method has been revised for this submission. The revision affects table A2.1.1 only.

Table A2.1.1 and table A2.1.2 present uncertainties for net emissions and gross emissions respectively.

Table A2.1.1 Uncertainty calculation (including LULUCF) for New Zealand's Greenhouse Gas Inventory 1990–2021 (IPCC, 2006, Approach 1)

IPCC source category	Gas	1990 emissions or absolute value of removals (kt CO ₂ -e)	2021 emissions or absolute value of removals (kt CO ₂ -e)	Activity data uncertainty (%)	Emission or removal factor uncertainty (%)	Combined uncertainty (%)	Combined uncertainty as a per cent of the national total in 1990 (%)	Combined uncertainty as a per cent of the national total in 2021 (%)	Type A sensitivity (%)	Type B sensitivity (%)	Uncertainty in the trend in national total introduced by emission or removal factor uncertainty (%)	Uncertainty in trend in national total introduced by activity data uncertainty (%)	Uncertainty introduced into the trend in the national total (%)	Combined uncertainty of the national total in 1990	Combined uncertainty of the national total in 2021
Energy – Gaseous fuels	CO ₂	7,027.14	6,111.00	4.4	2.4	5.0	0.7917	0.5502	0.0601	0.1372	0.1449	0.8541	0.8663	0.6268	0.3027
Energy – Liquid fuels	CO ₂	11,788.74	19,199.14	0.6	0.5	0.8	0.2042	0.2657	0.0996	0.4310	0.0498	0.3581	0.3616	0.0417	0.0706
Energy – Other fossil fuels	CO ₂	7.31	74.84	5.0	5.0	7.1	0.0012	0.0095	0.0015	0.0017	0.0074	0.0119	0.0140	0.0000	0.0001
Energy – Solid fuels	CO ₂	3,203.74	4,333.31	3.0	2.2	3.7	0.2684	0.2901	0.0073	0.0973	0.0158	0.4172	0.4175	0.0720	0.0841
Energy – Fugitive – oil exploration	CO ₂	0.00	0.00	0.6	100.0	100.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Energy – Fugitive – oil production	CO ₂	0.00	0.00	0.6	100.0	100.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Energy – Fugitive – oil transport	CO ₂	0.01	0.00	0.6	100.0	100.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Energy – Fugitive – gas production	CO ₂	0.21	0.18	4.4	100.0	100.1	0.0005	0.0003	0.0000	0.0000	0.0002	0.0000	0.0002	0.0000	0.0000
Energy – Fugitive – gas transmission and storage	CO ₂	0.01	0.16	4.4	100.0	100.1	0.0000	0.0003	0.0000	0.0000	0.0003	0.0000	0.0003	0.0000	0.0000
Energy – Fugitive – gas distribution	CO ₂	1.45	0.37	4.4	100.0	100.1	0.0033	0.0007	0.0000	0.0000	0.0032	0.0001	0.0032	0.0000	0.0000
Energy – Fugitive – venting and flaring	CO ₂	229.48	272.90	4.4	2.4	5.0	0.0259	0.0246	0.0003	0.0061	0.0008	0.0381	0.0381	0.0007	0.0006
Energy – Fugitive – other forms of energy production	CO ₂	228.58	431.59	5.0	5.0	7.1	0.0363	0.0547	0.0033	0.0097	0.0163	0.0685	0.0704	0.0013	0.0030
2.A.1 Cement production	CO ₂	448.75	369.63	1.0	1.0	1.4	0.0142	0.0094	0.0043	0.0083	0.0043	0.0117	0.0125	0.0002	0.0001
2.A.2 Lime production	CO ₂	82.60	103.68	2.0	2.0	2.8	0.0052	0.0053	0.0000	0.0023	0.0000	0.0066	0.0066	0.0000	0.0000
2.A.4.a Ceramics	CO ₂	0.01	0.01	50.0	20.0	53.9	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
2.A.4.b Other uses of soda ash	CO ₂	5.87	6.36	3.0	2.0	3.6	0.0005	0.0004	0.0000	0.0001	0.0000	0.0006	0.0006	0.0000	0.0000
2.A.4.d Other – Other uses of limestone	CO ₂	24.63	49.52	3.0	2.0	3.6	0.0020	0.0032	0.0004	0.0011	0.0008	0.0047	0.0048	0.0000	0.0000
2.B.1 Ammonia production	CO ₂	21.68	16.68	2.0	6.0	6.3	0.0031	0.0019	0.0002	0.0004	0.0014	0.0011	0.0018	0.0000	0.0000
2.B.5.b Calcium carbide	CO ₂	1.43	1.43	50.0	50.0	70.7	0.0023	0.0018	0.0000	0.0000	0.0004	0.0023	0.0023	0.0000	0.0000
2.B.10 Hydrogen production	CO ₂	152.29	43.37	2.0	6.0	6.3	0.0216	0.0049	0.0033	0.0010	0.0198	0.0028	0.0200	0.0005	0.0000
2.C.1 Iron and steel	CO ₂	1,306.73	1,718.13	5.0	7.0	8.6	0.2523	0.2651	0.0019	0.0386	0.0130	0.2727	0.2730	0.0637	0.0703
2.C.3.a Aluminium	CO ₂	448.98	541.80	5.0	2.0	5.4	0.0543	0.0523	0.0004	0.0122	0.0009	0.0860	0.0860	0.0029	0.0027
2.C.5 Secondary lead production	CO ₂	1.80	0.00	50.0	50.0	70.7	0.0029	0.0000	0.0001	0.0000	0.0025	0.0000	0.0025	0.0000	0.0000
2.D.1 Lubricant use	CO ₂	22.83	36.25	20.0	50.0	53.9	0.0276	0.0350	0.0002	0.0008	0.0086	0.0230	0.0246	0.0008	0.0012
2.D.2 Paraffin wax	CO ₂	2.35	2.35	20.0	100.0	102.0	0.0054	0.0043	0.0000	0.0001	0.0013	0.0015	0.0020	0.0000	0.0000

IPCC source category	Gas	1990 emissions or absolute value of removals (kt CO ₂ -e)	2021 emissions or absolute value of removals (kt CO ₂ -e)	Activity data uncertainty (%)	Emission or removal factor uncertainty (%)	Combined uncertainty (%)	Combined uncertainty as a per cent of the national total in 1990 (%)	Combined uncertainty as a per cent of the national total in 2021 (%)	Type A sensitivity (%)	Type B sensitivity (%)	Uncertainty in the trend in national total introduced by emission or removal factor uncertainty (%)	Uncertainty in trend in national total introduced by activity data uncertainty (%)	Uncertainty introduced into the trend in the national total (%)	Combined uncertainty of the national total in 1990	Combined uncertainty of the national total in 2021
2.D.3 Other – Urea catalyst in road transport	CO ₂	0.00	3.82	50.0	10.0	51.0	0.0000	0.0035	0.0001	0.0001	0.0009	0.0061	0.0061	0.0000	0.0000
Agriculture – Liming	CO ₂	296.48	355.64	3.4	50.0	50.1	0.3335	0.3197	0.0003	0.0080	0.0173	0.0384	0.0421	0.1112	0.1022
Agriculture – Urea application	CO ₂	39.19	553.19	10.0	50.0	51.0	0.0449	0.5060	0.0113	0.0124	0.5658	0.1756	0.5925	0.0020	0.2560
LULUCF – Forest land	CO ₂	-19,275.86	-15,994.80	0.0	80.8	80.8	-34.9703	-23.1891	0.1832	0.3590	14.8065	0.0000	14.8065	1222.9193	537.7331
LULUCF – Cropland	CO ₂	469.02	372.19	0.0	71.3	71.3	0.7507	0.4760	0.0048	0.0084	0.3436	0.0000	0.3436	0.5635	0.2266
LULUCF – Grassland	CO ₂	641.98	2,708.07	0.0	42.5	42.5	0.6122	2.0637	0.0427	0.0608	1.8161	0.0000	1.8161	0.3748	4.2590
LULUCF – Wetlands	CO ₂	-8.44	11.38	0.0	77.7	77.7	-0.0147	0.0159	0.0005	0.0003	0.0383	0.0000	0.0383	0.0002	0.0003
LULUCF – Settlements	CO ₂	75.68	122.16	0.0	63.4	63.4	0.1077	0.1389	0.0006	0.0027	0.0391	0.0000	0.0391	0.0116	0.0193
LULUCF – Other Land	CO ₂	13.66	81.63	0.0	86.9	86.9	0.0267	0.1273	0.0014	0.0018	0.1259	0.0000	0.1259	0.0007	0.0162
LULUCF – Harvested wood products	CO ₂	-2,481.21	-8,658.83	0.0	68.2	68.2	-3.8006	-10.5992	0.1247	0.1944	8.5121	0.0000	8.5121	14.4449	112.3425
Waste – Incineration and open burning of waste	CO ₂	158.91	90.62	50.0	40.0	64.0	0.2284	0.1041	0.0024	0.0020	0.0972	0.1438	0.1736	0.0522	0.0108
Tokelau Energy Industries – Sectoral approach – liquid	CO ₂	0.23	0.52	10.0	7.0	12.2	0.0001	0.0001	0.0000	0.0000	0.0000	0.0002	0.0002	0.0000	0.0000
Tokelau Gas Diesel Oil – Sectoral approach – liquid	CO ₂	0.90	1.37	50.0	1.5	50.0	0.0010	0.0012	0.0000	0.0000	0.0000	0.0022	0.0022	0.0000	0.0000
Tokelau Other/Residential – Sectoral approach – liquid	CO ₂	0.12	0.09	20.0	7.0	21.2	0.0001	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.0000	0.0000
Tokelau Waste – Incineration and open burning of waste	CO ₂	0.05	0.04	50.0	40.0	64.0	0.0001	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.0000	0.0000
Energy – Gaseous fuels	CH ₄	9.05	3.98	4.4	50.0	50.2	0.0102	0.0036	0.0002	0.0001	0.0082	0.0006	0.0083	0.0001	0.0000
Energy – Liquid fuels	CH ₄	87.51	35.44	0.6	50.0	50.0	0.0982	0.0318	0.0017	0.0008	0.0831	0.0007	0.0831	0.0096	0.0010
Energy – Other fossil fuels	CH ₄	0.03	0.46	5.0	50.0	50.2	0.0000	0.0004	0.0000	0.0000	0.0005	0.0001	0.0005	0.0000	0.0000
Energy – Solid fuels	CH ₄	36.35	17.81	3.0	50.0	50.1	0.0409	0.0160	0.0006	0.0004	0.0311	0.0017	0.0311	0.0017	0.0003
Energy – Biomass	CH ₄	69.44	69.60	50.0	50.0	70.7	0.1102	0.0883	0.0004	0.0016	0.0194	0.1105	0.1122	0.0121	0.0078
Energy – Fugitive – coal handling	CH ₄	328.03	62.44	3.0	50.0	50.1	0.3688	0.0561	0.0078	0.0014	0.3906	0.0060	0.3906	0.1360	0.0031
Energy – Fugitive – oil exploration	CH ₄	0.00	0.00	0.6	50.0	50.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Energy – Fugitive – oil production	CH ₄	0.06	0.02	0.6	50.0	50.0	0.0001	0.0000	0.0000	0.0000	0.0001	0.0000	0.0001	0.0000	0.0000
Energy – Fugitive – oil transport	CH ₄	1.68	0.82	0.6	50.0	50.0	0.0019	0.0007	0.0000	0.0000	0.0014	0.0000	0.0014	0.0000	0.0000
Energy – Fugitive – oil refining	CH ₄	2.73	2.49	0.6	50.0	50.0	0.0031	0.0022	0.0000	0.0001	0.0010	0.0000	0.0010	0.0000	0.0000

IPCC source category	Gas	1990 emissions or absolute value of removals (kt CO ₂ -e)	2021 emissions or absolute value of removals (kt CO ₂ -e)	Activity data uncertainty (%)	Emission or removal factor uncertainty (%)	Combined uncertainty (%)	Combined uncertainty as a per cent of the national total in 1990 (%)	Combined uncertainty as a per cent of the national total in 2021 (%)	Type A sensitivity (%)	Type B sensitivity (%)	Uncertainty in the trend in national total introduced by emission or removal factor uncertainty (%)	Uncertainty in trend in national total introduced by activity data uncertainty (%)	Uncertainty introduced into the trend in the national total (%)	Combined uncertainty of the national total in 1990	Combined uncertainty of the national total in 2021
Energy – Fugitive – gas production	CH ₄	143.45	126.83	4.4	50.0	50.2	0.1616	0.1142	0.0012	0.0028	0.0591	0.0177	0.0617	0.0261	0.0130
Energy – Fugitive – gas transmission and storage	CH ₄	2.47	19.23	4.4	100.0	100.1	0.0056	0.0345	0.0004	0.0004	0.0362	0.0027	0.0363	0.0000	0.0012
Energy – Fugitive – gas distribution	CH ₄	277.49	39.07	4.4	100.0	100.1	0.6235	0.0702	0.0069	0.0009	0.6917	0.0055	0.6917	0.3887	0.0049
Energy – Fugitive – venting and flaring	CH ₄	160.83	42.56	4.4	50.0	50.2	0.1812	0.0383	0.0036	0.0010	0.1781	0.0059	0.1782	0.0328	0.0015
Energy – Fugitive – other forms of energy production	CH ₄	54.79	114.56	5.0	50.0	50.2	0.0618	0.1033	0.0010	0.0026	0.0516	0.0182	0.0547	0.0038	0.0107
2.B.8 Methanol	CH ₄	27.60	77.62	2.0	80.0	80.0	0.0496	0.1114	0.0010	0.0017	0.0774	0.0049	0.0775	0.0025	0.0124
Agriculture – Enteric fermentation	CH ₄	26,960.43	27,859.05	3.9	15.5	16.0	9.6830	7.9959	0.1312	0.6254	2.0351	3.4491	4.0048	93.7607	63.9350
Agriculture – Manure management	CH ₄	722.02	1,571.92	5.0	20.0	20.6	0.3341	0.5813	0.0150	0.0353	0.3000	0.2495	0.3902	0.1116	0.3379
Agriculture – Burning of residues	CH ₄	22.62	18.71	6.0	20.0	20.9	0.0106	0.0070	0.0002	0.0004	0.0043	0.0036	0.0056	0.0001	0.0000
CH ₄ emissions associated with biomass burning (CO ₂ -e)	CH ₄	68.36	32.64	0.3	41.6	41.6	0.0638	0.0243	0.0012	0.0007	0.0494	0.0003	0.0494	0.0041	0.0006
Waste – Solid waste disposal	CH ₄	3,318.20	2,578.15	87.0	40.0	95.8	7.1348	4.4300	0.0353	0.0579	1.4123	7.1235	7.2621	50.9051	19.6252
Waste – Wastewater treatment and discharge	CH ₄	224.04	256.95	10.0	40.0	41.2	0.2074	0.1900	0.0005	0.0058	0.0210	0.0816	0.0842	0.0430	0.0361
Waste – Biological treatment of solid waste	CH ₄	2.74	42.02	100.0	100.0	141.4	0.0087	0.1066	0.0009	0.0009	0.0866	0.1334	0.1591	0.0001	0.0114
Waste – Incineration and open burning of waste	CH ₄	127.35	78.13	50.0	100.0	111.8	0.3196	0.1567	0.0018	0.0018	0.1823	0.1240	0.2205	0.1022	0.0246
Tokelau Energy Industries – Sectoral approach – liquid	CH ₄	0.00	0.00	10.0	50.0	51.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau Gas Diesel Oil – Sectoral approach – liquid	CH ₄	0.00	0.00	50.0	50.0	70.7	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau Other/Residential – Sectoral approach – liquid	CH ₄	0.00	0.00	20.0	50.0	53.9	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau Agriculture – Enteric fermentation	CH ₄	0.09	0.06	20.0	50.0	53.9	0.0001	0.0001	0.0000	0.0000	0.0001	0.0000	0.0001	0.0000	0.0000
Tokelau Agriculture – Manure management	CH ₄	1.06	0.76	20.0	30.0	36.1	0.0009	0.0005	0.0000	0.0000	0.0004	0.0005	0.0006	0.0000	0.0000
Tokelau Waste – Solid waste disposal	CH ₄	0.39	0.31	140.0	40.0	145.6	0.0013	0.0008	0.0000	0.0000	0.0002	0.0014	0.0014	0.0000	0.0000
Tokelau Waste – Wastewater treatment and discharge	CH ₄	0.15	0.27	10.0	40.0	41.2	0.0001	0.0002	0.0000	0.0000	0.0001	0.0001	0.0001	0.0000	0.0000

IPCC source category	Gas	1990 emissions or absolute value of removals (kt CO ₂ -e)	2021 emissions or absolute value of removals (kt CO ₂ -e)	Activity data uncertainty (%)	Emission or removal factor uncertainty (%)	Combined uncertainty (%)	Combined uncertainty as a per cent of the national total in 1990 (%)	Combined uncertainty as a per cent of the national total in 2021 (%)	Type A sensitivity (%)	Type B sensitivity (%)	Uncertainty in the trend in national total introduced by emission or removal factor uncertainty (%)	Uncertainty in trend in national total introduced by activity data uncertainty (%)	Uncertainty introduced into the trend in the national total (%)	Combined uncertainty of the national total in 1990	Combined uncertainty of the national total in 2021
Tokelau Waste – Incineration and open burning of waste	CH ₄	0.09	0.07	50.0	100.0	111.8	0.0002	0.0001	0.0000	0.0000	0.0001	0.0001	0.0001	0.0000	0.0000
Energy – Gaseous fuels	N ₂ O	5.53	3.15	4.4	50.0	50.2	0.0062	0.0028	0.0001	0.0001	0.0042	0.0004	0.0043	0.0000	0.0000
Energy – Liquid fuels	N ₂ O	157.91	180.67	0.6	50.0	50.0	0.1772	0.1621	0.0004	0.0041	0.0190	0.0034	0.0193	0.0314	0.0263
Energy – Other fossil fuels	N ₂ O	0.24	0.76	5.0	50.0	50.2	0.0003	0.0007	0.0000	0.0000	0.0005	0.0001	0.0005	0.0000	0.0000
Energy – Solid fuels	N ₂ O	14.90	19.91	3.0	50.0	50.1	0.0168	0.0179	0.0000	0.0004	0.0014	0.0019	0.0024	0.0003	0.0003
Energy – Biomass	N ₂ O	41.09	46.81	50.0	50.0	70.7	0.0652	0.0594	0.0001	0.0011	0.0052	0.0743	0.0745	0.0043	0.0035
Energy – Fugitive – venting and flaring	N ₂ O	0.06	0.03	4.4	100.0	100.1	0.0001	0.0001	0.0000	0.0000	0.0001	0.0000	0.0001	0.0000	0.0000
2.G.3 N ₂ O from product uses	N ₂ O	102.45	89.55	15.0	0.0	15.0	0.0345	0.0241	0.0009	0.0020	0.0000	0.0426	0.0426	0.0012	0.0006
Agriculture – Agricultural soils	N ₂ O	5,216.82	7,315.16	11.4	54.1	55.3	6.4759	7.2567	0.0176	0.1642	0.9549	2.6494	2.8162	41.9370	52.6590
Agriculture – Manure management	N ₂ O	49.72	108.60	5.0	100.0	100.1	0.1117	0.1951	0.0010	0.0024	0.1041	0.0172	0.1055	0.0125	0.0380
Agriculture – Burning of residues	N ₂ O	4.77	3.87	6.0	20.0	20.9	0.0022	0.0014	0.0000	0.0001	0.0009	0.0007	0.0012	0.0000	0.0000
Direct and indirect N ₂ O emissions (CO ₂ -e)	N ₂ O	299.96	225.19	0.0	58.3	58.3	0.3929	0.2357	0.0034	0.0051	0.1967	0.0000	0.1967	0.1544	0.0556
N ₂ O emissions associated with biomass burning (CO ₂ -e)	N ₂ O	25.62	22.22	0.3	41.6	41.6	0.0239	0.0166	0.0002	0.0005	0.0092	0.0002	0.0092	0.0006	0.0003
Waste – Wastewater treatment and discharge	N ₂ O	81.97	120.65	10.0	90.0	90.6	0.1666	0.1960	0.0004	0.0027	0.0365	0.0383	0.0529	0.0278	0.0384
Waste – Incineration and open burning of waste	N ₂ O	29.46	18.34	50.0	100.0	111.8	0.0739	0.0368	0.0004	0.0004	0.0416	0.0291	0.0508	0.0055	0.0014
Waste – Biological treatment of solid waste	N ₂ O	1.96	30.06	100.0	150.0	180.3	0.0079	0.0972	0.0006	0.0007	0.0930	0.0954	0.1332	0.0001	0.0094
Tokelau Energy Industries – Sectoral approach – liquid	N ₂ O	0.00	0.00	10.0	50.0	51.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau Gas Diesel Oil – Sectoral approach – liquid	N ₂ O	0.01	0.01	50.0	50.0	70.7	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau Other/Residential – Sectoral approach – liquid	N ₂ O	0.00	0.00	20.0	50.0	53.9	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau IPPU – Other product manufacture and use	N ₂ O	0.05	0.02	15.0	0.0	15.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau Waste – Wastewater treatment and discharge	N ₂ O	0.02	0.00	10.0	90.0	90.6	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000

IPCC source category	Gas	1990 emissions or absolute value of removals (kt CO ₂ -e)	2021 emissions or absolute value of removals (kt CO ₂ -e)	Activity data uncertainty (%)	Emission or removal factor uncertainty (%)	Combined uncertainty (%)	Combined uncertainty as a per cent of the national total in 1990 (%)	Combined uncertainty as a per cent of the national total in 2021 (%)	Type A sensitivity (%)	Type B sensitivity (%)	Uncertainty in the trend in national total introduced by emission or removal factor uncertainty (%)	Uncertainty in trend in national total introduced by activity data uncertainty (%)	Uncertainty introduced into the trend in the national total (%)	Combined uncertainty of the national total in 1990	Combined uncertainty of the national total in 2021
Tokelau Waste – Incineration and open burning of waste	N ₂ O	0.01	0.01	50.0	100.0	111.8	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
2.F.1 Refrigeration and air conditioning	HFCs	0.00	1401.83	24.0	0.0	24.0	0.0000	0.6035	0.0315	0.0315	0.0000	1.0680	1.0680	0.0000	0.3642
2.F.2 Foam blowing agents	HFCs	0.00	6.31	12.0	50.0	51.4	0.0000	0.0058	0.0001	0.0001	0.0071	0.0024	0.0075	0.0000	0.0000
2.F.3 Fire protection	HFCs	0.00	2.18	30.0	30.0	42.4	0.0000	0.0017	0.0000	0.0000	0.0015	0.0021	0.0025	0.0000	0.0000
2.F.4 Aerosols	HFCs	0.00	72.67	25.0	10.0	26.9	0.0000	0.0351	0.0016	0.0016	0.0163	0.0577	0.0599	0.0000	0.0012
2.C.3.a Aluminium	PFCs	909.95	50.68	5.0	30.0	30.4	0.6212	0.0277	0.0244	0.0011	0.7325	0.0080	0.7326	0.3859	0.0008
2.F.1 Refrigeration and air conditioning	PFCs	0.00	0.00	25.0	0.0	25.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
2.G.2 Other product use	PFCs	0.00	0.01	80.0	0.0	80.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
2.C.4 Magnesium production	SF ₆	2.74	0.00	100.0	30.0	104.4	0.0064	0.0000	0.0001	0.0000	0.0023	0.0000	0.0023	0.0000	0.0000
2.G.1 Electrical equipment	SF ₆	14.50	12.99	20.0	30.0	36.1	0.0117	0.0084	0.0001	0.0003	0.0035	0.0083	0.0089	0.0001	0.0001
2.G.2 Other product use	SF ₆	2.74	2.74	80.0	0.0	80.0	0.0049	0.0039	0.0000	0.0001	0.0000	0.0069	0.0069	0.0000	0.0000
Tokelau IPPU – Product uses as substitutes for ODS	HFCs	0.00	0.23	32.0	0.0	32.0	0.0000	0.0001	0.0000	0.0000	0.0000	0.0002	0.0002	0.0000	0.0000
Total emissions/removals		44,548.8	55,746.4				37.8%		Uncertainty in the final year		28.2%	Uncertainty in the trend		19.4%	

Table A2.1.2 Uncertainty calculation (excluding LULUCF) for New Zealand's Greenhouse Gas Inventory 1990–2021 (IPCC, 2006, Approach 1)

IPCC source category	Gas	1990 emissions or absolute value of removals (kt CO ₂ -e)	2021 emissions or absolute value of removals (kt CO ₂ -e)	Activity data uncertainty (%)	Emission or removal factor uncertainty (%)	Combined uncertainty (%)	Combined uncertainty as a per cent of the national total in 2021 (%)	Combined uncertainty as a per cent of the national total in the last year (%)	Type A sensitivity (%)	Type B sensitivity (%)	Uncertainty in the trend in national total introduced by emission or removal factor uncertainty (%)	Uncertainty in trend in national total introduced by activity data uncertainty (%)	Uncertainty introduced into the trend in the national total (%)	Combined uncertainty of the national total in 1990	Combined uncertainty of the national total in 2021
Energy – Gaseous fuels	CO ₂	7,027.14	6,111.00	4.4	2.4	5.0	0.5450	0.3992	0.0344	0.0944	0.0830	0.5879	0.5937	0.2970	0.1594
Energy – Liquid fuels	CO ₂	11,788.74	19,199.14	0.6	0.5	0.8	0.1405	0.1928	0.0803	0.2966	0.0401	0.2465	0.2498	0.0198	0.0372
Energy – Other fossil fuels	CO ₂	7.31	74.84	5.0	5.0	7.1	0.0008	0.0069	0.0010	0.0012	0.0051	0.0082	0.0096	0.0000	0.0000
Energy – Solid fuels	CO ₂	3,203.74	4,333.31	3.0	2.2	3.7	0.1847	0.2105	0.0082	0.0670	0.0178	0.2872	0.2877	0.0341	0.0443
Energy – Fugitive – oil exploration	CO ₂	0.00	0.00	0.6	100.0	100.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Energy – Fugitive – oil production	CO ₂	0.00	0.00	0.6	100.0	100.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Energy – Fugitive – oil transport	CO ₂	0.01	0.00	0.6	100.0	100.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Energy – Fugitive – gas production	CO ₂	0.21	0.18	4.4	100.0	100.1	0.0003	0.0002	0.0000	0.0000	0.0001	0.0000	0.0001	0.0000	0.0000
Energy – Fugitive – gas transmission and storage	CO ₂	0.01	0.16	4.4	100.0	100.1	0.0000	0.0002	0.0000	0.0000	0.0002	0.0000	0.0002	0.0000	0.0000
Energy – Fugitive – gas distribution	CO ₂	1.45	0.37	4.4	100.0	100.1	0.0022	0.0005	0.0000	0.0000	0.0021	0.0000	0.0021	0.0000	0.0000
Energy – Fugitive – venting and flaring	CO ₂	229.48	272.90	4.4	2.4	5.0	0.0178	0.0178	0.0000	0.0042	0.0000	0.0263	0.0263	0.0003	0.0003
Energy – Fugitive – other forms of energy production	CO ₂	228.58	431.59	5.0	5.0	7.1	0.0250	0.0397	0.0025	0.0067	0.0124	0.0472	0.0488	0.0006	0.0016
2.A.1 Cement production	CO ₂	448.75	369.63	1.0	1.0	1.4	0.0	0.0	0.0025	0.0057	0.0025	0.0081	0.0085	0.0001	0.0000
2.A.2 Lime production	CO ₂	82.60	103.68	2.0	2.0	2.8	0.0	0.0	0.0001	0.0016	0.0002	0.0045	0.0045	0.0000	0.0000
2.A.4.a Ceramics	CO ₂	0.01	0.01	50.0	20.0	53.9	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
2.A.4.b Other uses of soda ash	CO ₂	5.87	6.36	3.0	2.0	3.6	0.0	0.0	0.0000	0.0001	0.0000	0.0004	0.0004	0.0000	0.0000
2.A.4.d Other – Other uses of limestone	CO ₂	24.63	49.52	3.0	2.0	3.6	0.0	0.0	0.0003	0.0008	0.0006	0.0032	0.0033	0.0000	0.0000
2.B.1 Ammonia production	CO ₂	21.68	16.68	2.0	6.0	6.3	0.0	0.0	0.0001	0.0003	0.0008	0.0007	0.0011	0.0000	0.0000
2.B.5.b Calcium carbide	CO ₂	1.43	1.43	50.0	50.0	70.7	0.0	0.0	0.0000	0.0000	0.0002	0.0016	0.0016	0.0000	0.0000
2.B.10 Hydrogen production	CO ₂	152.29	43.37	2.0	6.0	6.3	0.0	0.0	0.0021	0.0007	0.0127	0.0019	0.0129	0.0002	0.0000
2.C.1 Iron and steel	CO ₂	1,306.73	1,718.13	5.0	7.0	8.6	0.2	0.2	0.0026	0.0265	0.0181	0.1877	0.1886	0.0302	0.0370
2.C.3.a Aluminium	CO ₂	448.98	541.80	5.0	2.0	5.4	0.0	0.0	0.0001	0.0084	0.0003	0.0592	0.0592	0.0014	0.0014

IPCC source category	Gas	1990 emissions or absolute value of removals (kt CO ₂ -e)	2021 emissions or absolute value of removals (kt CO ₂ -e)	Activity data uncertainty (%)	Emission or removal factor uncertainty (%)	Combined uncertainty (%)	Combined uncertainty as a per cent of the national total in 2021 (%)	Combined uncertainty as a per cent of the national total in the last year (%)	Type A sensitivity (%)	Type B sensitivity (%)	Uncertainty in the trend in national total introduced by emission or removal factor uncertainty (%)	Uncertainty in trend in national total introduced by activity data uncertainty (%)	Uncertainty introduced into the trend in the national total (%)	Combined uncertainty of the national total in 1990	Combined uncertainty of the national total in 2021
2.C.5 Secondary lead production	CO ₂	1.80	0.00	50.0	50.0	70.7	0.0	0.0	0.0000	0.0000	0.0017	0.0000	0.0017	0.0000	0.0000
2.D.1 Lubricant use	CO ₂	22.83	36.25	20.0	50.0	53.9	0.0	0.0	0.0001	0.0006	0.0071	0.0158	0.0173	0.0004	0.0006
2.D.2 Paraffin wax	CO ₂	2.35	2.35	20.0	100.0	102.0	0.0	0.0	0.0000	0.0000	0.0007	0.0010	0.0012	0.0000	0.0000
2.D.3 Other: Urea catalyst in road transport	CO ₂	0.00	3.82	50.0	10.0	51.0	0.0	0.0	0.0001	0.0001	0.0006	0.0042	0.0042	0.0000	0.0000
Agriculture – Liming	CO ₂	296.48	355.64	3.4	50.0	50.1	0.2	0.2	0.0001	0.0055	0.0029	0.0264	0.0266	0.0527	0.0538
Agriculture – Urea application	CO ₂	39.19	553.19	10.0	50.0	51.0	0.0	0.4	0.0078	0.0085	0.3914	0.1209	0.4097	0.0010	0.1348
Waste – Incineration and open burning of waste	CO ₂	158.91	90.62	50.0	40.0	64.0	0.2	0.1	0.0015	0.0014	0.0606	0.0990	0.1161	0.0247	0.0057
Tokelau Energy Industries – Sectoral approach – liquid	CO ₂	0.23	0.52	10.0	7.0	12.2	0.0	0.0	0.0000	0.0000	0.0000	0.0001	0.0001	0.0000	0.0000
Tokelau Gas Diesel Oil – Sectoral approach – liquid	CO ₂	0.90	1.37	50.0	1.5	50.0	0.0	0.0	0.0000	0.0000	0.0000	0.0015	0.0015	0.0000	0.0000
Tokelau Other/Residential – Sectoral approach – liquid	CO ₂	0.12	0.09	20.0	7.0	21.2	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau Waste – Incineration and open burning of waste	CO ₂	0.05	0.04	50.0	40.0	64.0	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Energy – Gaseous fuels	CH ₄	9.05	3.98	4.4	50.0	50.2	0.0	0.0	0.0001	0.0001	0.0052	0.0004	0.0052	0.0000	0.0000
Energy – Liquid fuels	CH ₄	87.51	35.44	0.6	50.0	50.0	0.1	0.0	0.0011	0.0005	0.0529	0.0005	0.0529	0.0046	0.0005
Energy – Other fossil fuels	CH ₄	0.03	0.46	5.0	50.0	50.2	0.0	0.0	0.0000	0.0000	0.0003	0.0001	0.0003	0.0000	0.0000
Energy – Solid fuels	CH ₄	36.35	17.81	3.0	50.0	50.1	0.0	0.0	0.0004	0.0003	0.0196	0.0012	0.0196	0.0008	0.0001
Energy – Biomass	CH ₄	69.44	69.60	50.0	50.0	70.7	0.1	0.1	0.0002	0.0011	0.0099	0.0760	0.0767	0.0058	0.0041
Energy – Fugitive – coal handling	CH ₄	328.03	62.44	3.0	50.0	50.1	0.3	0.0	0.0051	0.0010	0.2526	0.0041	0.2526	0.0645	0.0017
Energy – Fugitive – oil exploration	CH ₄	0.00	0.00	0.6	50.0	50.0	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Energy – Fugitive – oil production	CH ₄	0.06	0.02	0.6	50.0	50.0	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Energy – Fugitive – oil transport	CH ₄	1.68	0.82	0.6	50.0	50.0	0.0	0.0	0.0000	0.0000	0.0009	0.0000	0.0009	0.0000	0.0000
Energy – Fugitive – oil refining	CH ₄	2.73	2.49	0.6	50.0	50.0	0.0	0.0	0.0000	0.0000	0.0006	0.0000	0.0006	0.0000	0.0000
Energy – Fugitive – gas production	CH ₄	143.45	126.83	4.4	50.0	50.2	0.1	0.1	0.0007	0.0020	0.0336	0.0122	0.0357	0.0124	0.0069

IPCC source category	Gas	1990 emissions or absolute value of removals (kt CO ₂ -e)	2021 emissions or absolute value of removals (kt CO ₂ -e)	Activity data uncertainty (%)	Emission or removal factor uncertainty (%)	Combined uncertainty (%)	Combined uncertainty as a per cent of the national total in 2021 (%)	Combined uncertainty as a per cent of the national total in the last year (%)	Type A sensitivity (%)	Type B sensitivity (%)	Uncertainty in the trend in national total introduced by emission or removal factor uncertainty (%)	Uncertainty in trend in national total introduced by activity data uncertainty (%)	Uncertainty introduced into the trend in the national total (%)	Combined uncertainty of the national total in 1990	Combined uncertainty of the national total in 2021
Energy – Fugitive – gas transmission and storage	CH ₄	2.47	19.23	4.4	100.0	100.1	0.0	0.0	0.0003	0.0003	0.0252	0.0018	0.0252	0.0000	0.0006
Energy – Fugitive – gas distribution	CH ₄	277.49	39.07	4.4	100.0	100.1	0.4	0.1	0.0045	0.0006	0.4485	0.0038	0.4486	0.1842	0.0026
Energy – Fugitive – venting and flaring	CH ₄	160.83	42.56	4.4	50.0	50.2	0.1	0.0	0.0023	0.0007	0.1146	0.0041	0.1147	0.0156	0.0008
Energy – Fugitive – other forms of energy production	CH ₄	54.79	114.56	5.0	50.0	50.2	0.0	0.1	0.0008	0.0018	0.0383	0.0125	0.0403	0.0018	0.0056
2.B.8 Methanol	CH ₄	27.60	77.62	2.0	80.0	80.0	0.0	0.1	0.0007	0.0012	0.0555	0.0034	0.0556	0.0012	0.0065
Agriculture – Enteric fermentation	CH ₄	26,960.43	27,859.05	3.9	15.5	16.0	6.7	5.8	0.0638	0.4305	0.9894	2.3741	2.5721	44.4238	33.6645
Agriculture – Manure management	CH ₄	722.02	1,571.92	5.0	20.0	20.6	0.2	0.4	0.0110	0.0243	0.2209	0.1717	0.2798	0.0529	0.1779
Agriculture – Burning of residues	CH ₄	22.62	18.71	6.0	20.0	20.9	0.0	0.0	0.0001	0.0003	0.0025	0.0025	0.0035	0.0001	0.0000
Waste – Solid waste disposal	CH ₄	3,318.20	2,578.15	87.0	40.0	95.8	4.9	3.2	0.0210	0.0398	0.8405	4.9033	4.9748	24.1188	10.3335
Waste – Wastewater treatment and discharge	CH ₄	224.04	256.95	10.0	40.0	41.2	0.1	0.1	0.0001	0.0040	0.0056	0.0561	0.0564	0.0204	0.0190
Waste – Biological treatment of solid waste	CH ₄	2.74	42.02	100.0	100.0	141.4	0.0	0.1	0.0006	0.0006	0.0599	0.0918	0.1096	0.0000	0.0060
Waste – Incineration and open burning of waste	CH ₄	127.35	78.13	50.0	100.0	111.8	0.2	0.1	0.0011	0.0012	0.1129	0.0854	0.1415	0.0484	0.0129
Tokelau Energy industries – Sectoral approach – liquid	CH ₄	0.00	0.00	10.0	50.0	51.0	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau Gas Diesel Oil – Sectoral approach – liquid	CH ₄	0.00	0.00	50.0	50.0	70.7	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau Other/Residential – Sectoral approach – liquid	CH ₄	0.00	0.00	20.0	50.0	53.9	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau Agriculture – Enteric fermentation	CH ₄	0.09	0.06	20.0	50.0	53.9	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau Agriculture – Manure management	CH ₄	1.06	0.76	20.0	30.0	36.1	0.0	0.0	0.0000	0.0000	0.0002	0.0003	0.0004	0.0000	0.0000
Tokelau Waste – Solid waste disposal	CH ₄	0.39	0.31	140.0	40.0	145.6	0.0	0.0	0.0000	0.0000	0.0001	0.0009	0.0009	0.0000	0.0000

IPCC source category	Gas	1990 emissions or absolute value of removals (kt CO ₂ -e)	2021 emissions or absolute value of removals (kt CO ₂ -e)	Activity data uncertainty (%)	Emission or removal factor uncertainty (%)	Combined uncertainty (%)	Combined uncertainty as a per cent of the national total in 2021 (%)	Combined uncertainty as a per cent of the national total in the last year (%)	Type A sensitivity (%)	Type B sensitivity (%)	Uncertainty in the trend in national total introduced by emission or removal factor uncertainty (%)	Uncertainty in trend in national total introduced by activity data uncertainty (%)	Uncertainty introduced into the trend in the national total (%)	Combined uncertainty of the national total in 1990	Combined uncertainty of the national total in 2021
Tokelau Waste – Wastewater treatment and discharge	CH ₄	0.15	0.27	10.0	40.0	41.2	0.0	0.0	0.0000	0.0000	0.0001	0.0001	0.0001	0.0000	0.0000
Tokelau Waste – Incineration and open burning of waste	CH ₄	0.09	0.07	50.0	100.0	111.8	0.0	0.0	0.0000	0.0000	0.0000	0.0001	0.0001	0.0000	0.0000
Energy – Gaseous fuels	N ₂ O	5.53	3.15	4.4	50.0	50.2	0.0	0.0	0.0001	0.0000	0.0026	0.0003	0.0027	0.0000	0.0000
Energy – Liquid fuels	N ₂ O	157.91	180.67	0.6	50.0	50.0	0.1	0.1	0.0001	0.0028	0.0052	0.0023	0.0057	0.0149	0.0138
Energy – Other fossil fuels	N ₂ O	0.24	0.76	5.0	50.0	50.2	0.0	0.0	0.0000	0.0000	0.0004	0.0001	0.0004	0.0000	0.0000
Energy – Solid fuels	N ₂ O	14.90	19.91	3.0	50.0	50.1	0.0	0.0	0.0000	0.0003	0.0017	0.0013	0.0022	0.0001	0.0002
Energy – Biomass	N ₂ O	41.09	46.81	50.0	50.0	70.7	0.0	0.0	0.0000	0.0007	0.0015	0.0511	0.0512	0.0020	0.0019
Energy – Fugitive – venting and flaring	N ₂ O	0.06	0.03	4.4	100.0	100.1	0.0	0.0	0.0000	0.0000	0.0001	0.0000	0.0001	0.0000	0.0000
2.G.3 N ₂ O from product uses	N ₂ O	102.45	89.55	15.0	0.0	15.0	0.0	0.0	0.0005	0.0014	0.0000	0.0294	0.0294	0.0006	0.0003
Agriculture – Agricultural soils	N ₂ O	5,216.82	7,315.16	11.4	54.1	55.3	4.5	5.3	0.0173	0.1130	0.9379	1.8237	2.0507	19.8697	27.7272
Agriculture – Manure management	N ₂ O	49.72	108.60	5.0	100.0	100.1	0.1	0.1	0.0008	0.0017	0.0766	0.0119	0.0775	0.0059	0.0200
Agriculture – Burning of residues	N ₂ O	4.77	3.87	6.0	20.0	20.9	0.0	0.0	0.0000	0.0001	0.0006	0.0005	0.0007	0.0000	0.0000
Waste – Wastewater treatment and discharge	N ₂ O	81.97	120.65	10.0	90.0	90.6	0.1	0.1	0.0004	0.0019	0.0325	0.0264	0.0418	0.0132	0.0202
Waste – Incineration and open burning of waste	N ₂ O	29.46	18.34	50.0	100.0	111.8	0.1	0.0	0.0003	0.0003	0.0257	0.0200	0.0326	0.0026	0.0007
Waste – Biological treatment of solid waste	N ₂ O	1.96	30.06	100.0	150.0	180.3	0.0	0.1	0.0004	0.0005	0.0643	0.0657	0.0919	0.0000	0.0050
Tokelau Energy Industries – Sectoral approach – liquid	N ₂ O	0.00	0.00	10.0	50.0	51.0	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau Gas Diesel Oil – Sectoral approach – liquid	N ₂ O	0.01	0.01	50.0	50.0	70.7	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau Other/Residential – Sectoral approach – liquid	N ₂ O	0.00	0.00	20.0	50.0	53.9	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau IPPU – Other product manufacture and use	N ₂ O	0.05	0.02	15.0	0.0	15.0	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
Tokelau Waste – wastewater treatment and discharge	N ₂ O	0.02	0.00	10.0	90.0	90.6	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000

IPCC source category	Gas	1990 emissions or absolute value of removals (kt CO ₂ -e)	2021 emissions or absolute value of removals (kt CO ₂ -e)	Activity data uncertainty (%)	Emission or removal factor uncertainty (%)	Combined uncertainty (%)	Combined uncertainty as a per cent of the national total in 2021 (%)	Combined uncertainty as a per cent of the national total in the last year (%)	Type A sensitivity (%)	Type B sensitivity (%)	Uncertainty in the trend in national total introduced by emission or removal factor uncertainty (%)	Uncertainty in trend in national total introduced by activity data uncertainty (%)	Uncertainty introduced into the trend in the national total (%)	Combined uncertainty of the national total in 1990	Combined uncertainty of the national total in 2021
Tokelau Waste – Incineration and open burning of waste	N ₂ O	0.01	0.01	50.0	100.0	111.8	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
2.F.1 Refrigeration and air conditioning	HFCs	0.00	1401.83	24.0	0.0	24.0	0.0	0.4	0.0217	0.0217	0.0000	0.7352	0.7352	0.0000	0.1918
2.F.2 Foam blowing agents	HFCs	0.00	6.31	12.0	50.0	51.4	0.0	0.0	0.0001	0.0001	0.0049	0.0017	0.0051	0.0000	0.0000
2.F.3 Fire protection	HFCs	0.00	2.18	30.0	30.0	42.4	0.0	0.0	0.0000	0.0000	0.0010	0.0014	0.0017	0.0000	0.0000
2.F.4 Aerosols	HFCs	0.00	72.67	25.0	10.0	26.9	0.0	0.0	0.0011	0.0011	0.0112	0.0397	0.0413	0.0000	0.0006
2.C.3.a Aluminium	PFCs	909.95	50.68	5.0	30.0	30.4	0.4	0.0	0.0159	0.0008	0.4771	0.0055	0.4772	0.1829	0.0004
2.F.1 Refrigeration and air conditioning	PFCs	0.00	0.00	25.0	0.0	25.0	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
2.G.2 Other product use	PFCs	0.00	0.01	80.0	0.0	80.0	0.0	0.0	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
2.C.4 Magnesium production	SF ₆	2.74	0.00	100.0	30.0	104.4	0.0	0.0	0.0001	0.0000	0.0015	0.0000	0.0015	0.0000	0.0000
2.G.1 Electrical equipment	SF ₆	14.50	12.99	20.0	30.0	36.1	0.0	0.0	0.0001	0.0002	0.0020	0.0057	0.0060	0.0001	0.0000
2.G.2 Other product use	SF ₆	2.74	2.74	80.0	0.0	80.0	0.0	0.0	0.0000	0.0000	0.0000	0.0048	0.0048	0.0000	0.0000
Tokelau IPPU – Product uses as substitutes for ODS	HFCs	0.00	0.23	32.0	0.0	32.0	0.0	0.0	0.0000	0.0000	0.0000	0.0002	0.0002	0.0000	0.0000
Total emissions/removals		64,720.1	76,824.6				9.5%		Uncertainty in the base year		8.5%		Uncertainty in the trend	6.1%	

Annex 2: References

IPCC. 2006. Eggleston HS, Buendia L, Miwa K, Ngara T, Tanabe K (eds). *2006 IPCC Guidelines for National Greenhouse Gas Inventories. Volume 1. General Guidance and Reporting*. IPCC National Greenhouse Gas Inventories Programme. Japan: Published for the IPCC by the Institute for Global Environmental Strategies.

UNFCCC. 2013. FCCC/CP/2013/Add.3. *Guidelines for the preparation of national communications by Parties included in Annex I to the Convention, Part I: UNFCCC reporting guidelines on annual greenhouse gas inventories (addendum to Decision 24/CP.19)*.

Annex 3: Detailed methodological information for other sectors

A3.1 Supplementary information for the Agriculture sector

A3.1.1 Livestock population data

Agricultural Production Census 2020 and Agricultural Production Survey 2021

Details of the Agricultural Production Census (APC) and Agricultural Production Survey (APS) are included to provide an understanding of the livestock statistics process and uncertainty values. The information here is provided by Stats NZ, with full details available from the Stats NZ website (www.stats.govt.nz/information-releases/agricultural-production-statistics-year-to-june-2021-final).

Stats NZ conducts the APC every five years, with the most recent census held in 2017. The final results from the 2022 APC are due to be released in mid-2023. In all other years, Stats NZ carries out the APS, which applies a similar method to the APC but targets about half of the businesses involved in agriculture or forestry production. The National Inventory Report is compiled with data from the APC and APS.

The 2021 APS used a stratified sample design to select a sample from the target population (all registered businesses that were engaged in agricultural production activity, including livestock, cropping, horticulture and forestry) or that owned land intended for agricultural activity during the year ended 30 June 2021). The response rate, or the estimated proportion of eligible businesses that responded to the 2021 APS, was 80 per cent.

The imputation levels of the 2020 APS and 2021 APS are provided in table A3.1.1. Full details on APC and APS data collection methodology can be found on the Stats NZ website (datainfolplus.stats.govt.nz).

Sampling error arises in the APS from selecting a sample of businesses and weighting the results rather than taking a complete enumeration (i.e. census). Non-sampling error arises from biases in the patterns of response and non-response, inaccuracies in reporting by respondents and errors in the recording and classification of data. Stats NZ adopts procedures to detect and minimise these types of errors, but they may still occur and are not easy to quantify.

Table A3.1.1 Imputation levels and sampling errors for recent Agricultural Production Surveys

Statistic	Proportion of total estimate imputed (%)		Relative sampling errors at 95% confidence interval (%)	
	2020	2021	2020	2021
Survey year				
Ewe hoggets put to ram	16	19	6	6
Breeding ewes, two tooth and over	17	22	3	3
Total number of sheep	17	21	3	3
Lambs born to ewe hoggets	16	19	5	6
Lambs born to ewes	17	22	3	3

Statistic Survey year	Proportion of total estimate imputed (%)		Relative sampling errors at 95% confidence interval (%)	
	2020	2021	2020	2021
Total number of lambs	17	21	3	3
Calves born alive to dairy heifers and/or cows	26	30	4	4
Dairy cows and heifers, in milk or calf	25	29	4	4
Total number of dairy cattle	24	28	3	4
Calves born alive to beef heifers and/or cows	18	23	4	3
Beef cows and heifers (in calf) one to two years	20	22	8	9
Beef cows and heifers (in calf) two years and over	18	23	4	3
Total number of beef cattle	19	23	3	3
Female deer mated	14	15	5	6
Total number of deer	16	16	4	6
Fawns born on farm and alive at four months	14	15	5	5
Total pigs	5	1	1	0
Area of wheat harvested	15	15	5	5
Area of barley harvested	14	20	8	8
Area of oat grain harvested	20	20	32	18
Area of maize grain harvested	18	19	11	13

Livestock characterisation in New Zealand's Tier 2 modelling

The delineation of the major livestock categories in New Zealand's Tier 2 livestock nutritional and energy requirements modelling (see table A3.1.2) are taken from population data collected by the APC and APS and Ministry for Primary Industries slaughter statistics.

Table A3.1.2 Characterisation of major livestock subcategories (dairy cattle, beef cattle, sheep and deer) in New Zealand's Tier 2 livestock modelling

Livestock category	Subcategory
Dairy cattle	Milking cows and heifers
	Growing females less than one year
	Growing females one to two years
	Breeding bulls
	Northland
	Auckland
	Waikato
	Bay of Plenty
	Gisborne
	Hawke's Bay
	Taranaki
	Manawatu–Whanganui
	Wellington
	Tasman
	Nelson
	Marlborough
	West Coast
	Canterbury
	Otago
Southland	

Livestock category	Subcategory
Beef cattle categories	Breeding growing cows less than one year
	Breeding growing cows one to two years
	Breeding growing cows two to three years
	Breeding mature cows
	Breeding bulls – mixed age
	Slaughter heifers less than one year
	Slaughter heifers one to two years
	Slaughter steers less than one year
	Slaughter steers one to two years
	Slaughter bulls less than one year
Slaughter bulls one to two years	
Sheep categories	Dry ewes
	Mature breeding ewes
	Growing breeding sheep
	Growing non-breeding sheep
	Wethers
	Lambs
Deer categories	Rams
	Breeding hinds
	Hinds less than one year
	Hinds one to two years
	Stags less than one year
	Stags one to two years
	Stags two to three years
Mixed age and breeding stags	

A3.1.2 Key parameters and emission factors used in the Agriculture sector

For the major livestock categories, milk yield varies over the course of a year, which affects energy requirements, feed intake and greenhouse gas emissions. Table A3.1.3 shows the proportions that are used to calculate milk yield for different months over the course of a year. Table A3.1.4 shows the emission factors used to calculate methane emissions from minor livestock species, while tables A3.1.5 and A3.1.6 show the emission factors used to calculate nitrous oxide emissions from agriculture. Table A3.1.7 shows some of the parameter values used to calculate nitrous oxide emissions.

Table A3.1.3 Proportion of annual milk yield each month for major livestock categories

Month	Dairy cattle	Beef cattle	Sheep	Deer
July	0.0088	0.0000	0.0000	0.0000
August	0.0578	0.0000	0.0000	0.0000
September	0.1213	0.1670	0.1639	0.0000
October	0.1503	0.1670	0.2541	0.0000
November	0.1425	0.1670	0.2459	0.1000
December	0.1282	0.1670	0.2541	0.2583
January	0.1109	0.1670	0.0820	0.2583
February	0.0900	0.1670	0.0000	0.2333

Month	Dairy cattle	Beef cattle	Sheep	Deer
March	0.0851	0.0000	0.0000	0.1500
April	0.0654	0.0000	0.0000	0.0000
May	0.0335	0.0000	0.0000	0.0000
June	0.0061	0.0000	0.0000	0.0000

Source: Suttie (2012) and Pickering and Fick (2015)

Note: All values presented in the table are rounded to four decimal places for presentation purposes and more precise values are available upon request.

Table A3.1.4 Methane emission factors for Tier 1 enteric fermentation livestock and manure management

Emission factor	Emission type	Source	Parameter value (kg CH ₄ /head/yr)
EF _{GOATS}	Enteric fermentation – goats	Lassey (2011)	9.0 ¹
EF _{HORSES}	Enteric fermentation – horses	IPCC (2006a), table 10.10	18.0
EF _{MULES}	Enteric fermentation – mules and asses	IPCC (2006a), table 10.10	10.0
EF _{SWINE}	Enteric fermentation – swine	Hill (2012)	1.06
EF _{ALPACA}	Enteric fermentation – alpaca	IPCC (2006a), table 10.10	8.0
MM _{GOATS}	Manure management – goats	IPCC (2006a), table 10.15	0.20
MM _{HORSES}	Manure management – horses	IPCC (2006a), table 10.15	2.34
MM _{MULES}	Manure management – mules and asses	IPCC (2006a), table 10.15	1.1
MM _{SWINE}	Manure management – swine	Hill (2012); IPCC (2000)	5.94
MM _{BROILERS}	Manure management – broilers	Fick et al. (2011)	0.022
MM _{LAYERS}	Manure management – layer hens	Fick et al. (2011)	0.016
MM _{OTHER POULTRY}	Manure management – other poultry	IPCC (1996), table 4.5	0.117
MM _{ALPACA}	Manure management – alpaca	New Zealand 1990 sheep value	0.103

Table A3.1.5 Emission factors for New Zealand's agricultural nitrous oxide emissions

Emission factor	Emissions	Source	Parameter value
EF ₁ (kg N ₂ O-N/kg N)	Direct emissions from nitrogen input to soil	Kelliher and de Klein (unpublished)	0.01
EF _{1-UREA} (kg N ₂ O-N/kg N)	Direct emissions from nitrogen input to soil from urea fertiliser	van der Weerden et al. (2016)	0.0059
EF _{1-DAIRY} (kg N ₂ O-N/kg N)	Direct emissions from nitrogen input to soil from dairy cattle manure	van der Weerden et al. (2016)	0.0025
EF ₂ (kg N ₂ O-N/ha-yr)	Direct emissions from organic soil mineralisation due to cultivation	IPCC (2006a), table 11.1	8.00
EF _{3SSD} (kg N ₂ O-N/kg N excreted)	Direct emissions from waste in solid waste and dry lot animal waste management systems	IPCC (2000), table 4.12	0.02
EF _{3(PRP-MINOR)} (kg N ₂ O-N/kg N excreted)	Direct emissions from manure (dung and urine) from minor grazing animals (i.e., <i>excluding</i> cattle, sheep and deer) in pasture, range and paddock systems	Carran et al. (1995); Muller et al. (1995); de Klein et al. (2003)	0.01

¹ Value is for 2020. In 1990, the value was EF 7.4 kg CH₄/head/year. Values for the intermediate years between 1990 and 2018 are calculated based on the estimated proportion of dairy goats in the overall goat population.

Emission factor	Emissions	Source	Parameter value
EF _{3(PRP DUNG)} (kg N ₂ O-N/kg N excreted)	Direct emissions from dung in pasture, range and paddock systems for cattle, sheep and deer (direct emission factors for urine are reported in table A3.1.6)	van der Weerden et al. (2019)	0.0012
EF _{3(OTHER)} (kg N ₂ O-N/kg N excreted)	Direct emissions from waste in other animal waste management systems	IPCC (2000), table 4.13	0.005
EF _{3(POULTRY)} (kg N ₂ O-N/kg N excreted)	Direct emissions from waste in other animal waste management systems – poultry specific	Fick et al. (2011)	0.001
EF ₄ (kg N ₂ O-N/kg NH _x -N)	Indirect emissions from volatilising nitrogen	IPCC (2006a), table 11.3	0.01
EF ₅ (kg N ₂ O-N/kg N leached and runoff)	Indirect emissions from leaching nitrogen	IPCC (2006a), table 11.3	0.0075

Table A3.1.6 Direct nitrous oxide emission factors for urine deposited by cattle, sheep and deer, by livestock type and slope

Livestock type	Emission factor by topography (kg N ₂ O–N/kg N excreted)	
	Flat and low sloped land (less than 12° gradient)	Medium and steep sloped land (greater than 12° gradient)
	EF _{3(PRP-FLAT)}	EF _{3(PRP-STEEP)}
All cattle (includes dairy and non-dairy)	0.0098	0.0033
Deer	0.0074	0.0020
Sheep	0.0050	0.0008

Source: Values used as calculated by van der Weerden et al. (2019)

Table A3.1.7 Parameter values for New Zealand’s agriculture nitrous oxide emissions

Parameter (fraction)	Fraction of the parameter	Source	Parameter value
Fra _{CGASF} (kg NH ₃ -N + NO _x -N/kg of synthetic fertiliser N applied)	Total of synthetic fertiliser emitted as NO _x or NH ₃	IPCC (2006a) verified by Sherlock et al. (2008)	0.1
Fra _{CGASM} (kg NH ₃ -N + NO _x -N/kg of N excreted by livestock)	Total of nitrogen emitted as NO _x or NH ₃	Sherlock et al. (2008)	0.1
Fra _{CLEACH(-H)} (kg N/kg fertiliser or manure N)	Nitrogen input to soils that is lost through leaching and run-off	Welten et al. (2021)	0.10 (Cropland) 0.08 (Grassland) 0.082 (Synthetic N fertiliser)
Fra _{CBURN} (kg N/kg crop-N)	Crop residue burned in fields	Thomas et al. (2008), table 14	Crop specific survey data
Fra _{CBURNL} (kg N/kg legume-N)	Legume crop residue burned in fields	Thomas et al. (2008) Practice does not occur in New Zealand	0
Fra _{CRENEW}	Fraction of land undergoing pasture renewal	Thomas et al. (2014)	Year specific
Fra _{REMOVE}	Fraction of nitrogen in above-ground residues removed for bedding, feed or construction	Thomas et al. (2014) Practice does not occur in New Zealand	0
Fra _{CFUEL} (N/kg N excreted)	Livestock nitrogen excretion in excrements burned for fuel	Practice does not occur in New Zealand	0

Some of the parameters used to calculate *Nitrous oxide emissions from crop residue returned to soil* and emissions from *Field burning of agricultural residues* are summarised in table A3.1.8. These values are taken from research conducted by Thomas et al. (2008, 2011).

Table A3.1.8 Parameter values for New Zealand's cropping emissions

Crop	HI	dmf	AG _N	Root Shoot ratio R _{BG}	BG _N
Wheat	0.41	0.86	0.005	0.1	0.009
Barley	0.46	0.86	0.005	0.1	0.009
Oats	0.30	0.86	0.005	0.1	0.009
Maize grain	0.50	0.86	0.007	0.1	0.007
Field seed peas	0.50	0.21	0.02	0.1	0.015
Lentils	0.50	0.86	0.02	0.1	0.015
Peas fresh and processed	0.45	0.86	0.03	0.1	0.015
Potatoes	0.90	0.22	0.02	0.1	0.01
Onions	0.80	0.11	0.02	0.1	0.01
Sweet corn	0.55	0.24	0.009	0.1	0.007
Squash	0.80	0.20	0.02	0.1	0.01
Herbage seeds	0.11	0.85	0.015	0.1	0.01
Legume seeds	0.09	0.85	0.04	0.1	0.01
Brassica seeds	0.20	0.85	0.01	0.1	0.008

Source: Thomas et al. (2008, 2011)

Note: AG_N = above-ground nitrogen residue; BG_N = below-ground nitrogen residue; dmf = dry-matter conversion factor; HI = harvest index; R_{BG} = ratio of below-ground residues to the harvest yield.

A3.1.3 Methodology and data used to allocate livestock excreta to different hill slopes, for cattle, sheep and deer

The emission factors used to calculate direct nitrous oxide (N₂O) emissions from all cattle, sheep and deer were described in detail in chapter 5, section 5.5.2. These pages explained the research behind the revised emission factors and how they were applied to estimate emissions from cattle, sheep and deer on different hill slopes.

These revised emission factors are disaggregated by slope (as well as by livestock type), and a methodology is used to calculate the amount of nitrogen (in the form of urine or dung) deposited on these different slopes. The steps described below are used to do this.

The nutrient transfer model outlined by Saggart et al. (2015) is used to allocate total dung and urine (calculated elsewhere in the inventory model) between low, medium and steep slopes. The nutrient transfer model was discussed by the Agriculture Inventory Advisory Panel in 2015, which agreed that the methodology used in the nutrient transfer model was appropriate. Beef + Lamb New Zealand Ltd provides data (on the topography and number of animals on different farm types) used in the nutrient transfer model.

Dairy excreta are not allocated to different slope types because the Agriculture Inventory assumes that all dairy cattle graze on flatland. The flatland/low slope emission factor for cattle urine (EF_{3(PRP FLAT)} = 0.0098) is applied to all dairy cattle urine.

Step 1: Calculations of total nitrogen excretion rates for each animal category

Total nitrogen excretion rates (N_{ex}) for each animal category are calculated using the methods described in chapter 5, section 5.3.2 of the National Inventory Report (*Nitrogen excretion rates for the major livestock categories*), and in chapter 5 of Pickering et al. (2020).

Step 2: Split of nitrogen between urine and dung

The total N_{ex} calculated in step 1 is split into urine and dung using the method described by Pacheco et al. (unpublished), and section 5.2.4 (beef cattle), section 5.3.5 (sheep) and section 5.4.5 (deer) of Pickering and Gibbs (2019).

Step 3: Allocating urine excreta to different hill slopes

The nutrient transfer model (described by Saggari et al. (2015)) uses Beef + Lamb New Zealand Ltd data on the proportion of sheep and beef farmland on different hill slopes to allocate urine excreta to different hill slopes. The nutrient transfer model accounts for the preference of animals to spend more time on flatter slopes. Using this model, the proportion of excreta deposited on low slopes is greater than the proportion of low slope land area, because animals spend more time on flatter land.

The equations and variables needed to allocate excreta to different slopes are outlined in table A3.1.9 and figures A3.1.1 and A3.1.2. For example, an area with 60 per cent low slopes and 25 per cent steep slopes will have 72 per cent of livestock urine deposited on low slope land ($0.45 \times 60 \text{ per cent} + 0.45 = 72 \text{ per cent}$) and 14 per cent of livestock urine deposited on steep slope land. After the allocation of excreta to low and high slope areas, the remainder (14 per cent) is assumed to be deposited onto medium sloped land.

A single dung emission factor ($EF_{3(PRP-DUNG)} = 0.0012$) is used across all slope categories for cattle, sheep and deer. Therefore, dung excreta do not need to be allocated to different slopes.

Table A3.1.9 Allocation of urine deposition to low slope (0–12 degrees) and steep slope (more than 24 degrees), split by the percentage of low slope and steep slope land available

Allocation to flat land	
Percentage of low land area	Fraction urine deposition
Less than 1%	27x
1–5%	0.27
5–9%	0.405
9–35%	0.55
35–85%	$(0.45x + 0.45)$
Greater than 85%	$(0.5x + 0.5)$
Allocation to steep land	
Percentage of steep land area	Fraction urine deposition
Less than 1%	10x
1–20%	0.10
20–40%	0.14
40–60%	0.21
60–85%	0.28
Greater than 85%	$4.8x - 3.8$

Figure A3.1.1 Proportion of urine nitrogen (N) applied to low (0–12 degree) slopes using a nutrient transfer model (equal proportion line shown in grey for comparison)





Figure A3.1.2 Proportion of urine nitrogen (N) applied to steep (more than 24 degree) slopes using a nutrient transfer model (equal proportion line shown in grey for comparison)



Tables A3.1.10, A3.1.11, A3.1.12 and figure A3.1.3 provide examples of how this nutrient allocation methodology uses Beef + Lamb New Zealand Ltd data to allocate urine nitrogen (N) to different hill slopes. First, data on the number of sheep, beef cattle and deer in each farm class are used to allocate total urine N (calculated using the methods described in chapter 5, section 5.3.2 of the National Inventory Report) to these different farm classes (tables A3.1.11 and A3.1.12).

Table A3.1.10 Share of livestock population, and amount of urine nitrogen (N) deposition in 2021, by Beef + Lamb New Zealand Ltd farm class

Farm class	Percentage of sheep population on farm class (%)	Amount of sheep urine N on farm class (kg N)	Percentage of beef cattle population on farm class (%)	Amount of beef cattle urine N on farm class (kg N)	Percentage of deer population on farm class (%)	Amount of deer urine N on farm class (kg N)
1. South Island High Country	7.6	25,203,753	4.5	10,264,321	9.8	1,839,837
2. South Island Hill Country	11.8	38,867,028	6.8	15,559,265	7.9	1,492,620
3. North Island Hard Hill Country	16.6	54,606,833	14.6	33,102,990	6.5	1,216,719
4. North Island Hill Country	25.7	84,551,401	39.9	90,612,020	35.2	6,611,052
5. North Island Intensive Finishing	5.2	17,176,630	11.1	25,337,902	0.4	82,628
6. South Island Finishing Breeding	20.0	65,998,155	15.9	36,082,063	30.1	5,661,843
7. South Island Intensive Finishing	10.5	34,705,890	3.2	7,302,269	10.1	1,902,388
8. South Island Mixed Finishing	2.5	8,370,924	4.0	9,119,056	0.0	0
Total		329,480,614		227,379,885		18,807,088

Each farm class has a different proportion of land in low, medium and steep slopes, as shown in table A3.1.11. These data are combined with the nutrient transfer methodology to calculate total urine N that is estimated to be deposited on different hill slopes for different animal categories. From this point, direct N₂O emissions can be calculated using the emission factors in chapter 5, table 5.5.3.

Table A3.1.11 Proportion of total sheep, beef and deer land on different hill slopes, by Beef + Lamb New Zealand Ltd farm class, for 2020/21

Farm class	Land type by slope		
	Flat/low (0–12° slope) (%)	Rolling/medium (12–24° slope) (%)	Steep (>24° slope) (%)
1. South Island High Country	7.9	28.5	63.6
2. South Island Hill Country	16.7	25.1	58.2
3. North Island Hard Hill Country	8.5	33.8	57.6
4. North Island Hill Country	16.4	53.4	30.2
5. North Island Intensive Finishing	44.4	50.8	4.8
6. South Island Finishing Breeding	33.8	46.0	20.2
7. South Island Intensive Finishing	62.1	37.9	0.0
8. South Island Mixed Finishing	93.5	6.5	0.0
Total sheep, beef and deer land	21.4	37.9	40.8


Note: The percentages may not add up to 100 per cent due to rounding.

Table A3.1.12 Proportion of total sheep, beef and deer urine nitrogen deposited on different hill slopes, by Beef + Lamb New Zealand Ltd farm class, for 2021

Farm class	Flat/low	Rolling/medium	Steep
1. South Island High Country	0.41	0.32	0.28
2. South Island Hill Country	0.55	0.24	0.21
3. North Island Hard Hill Country	0.41	0.39	0.21
4. North Island Hill Country	0.55	0.31	0.14
5. North Island Intensive Finishing	0.65	0.25	0.10
6. South Island Finishing Breeding	0.55	0.31	0.14
7. South Island Intensive Finishing	0.73	0.27	0.00
8. South Island Mixed Finishing	0.97	0.03	0.00
Total sheep urine	0.55	0.30	0.15
Total beef urine	0.56	0.30	0.15
Total deer urine	0.55	0.31	0.15
Total sheep, beef and deer urine	0.55	0.30	0.15

Note: The proportions may not add up to 1 due to rounding.

Figure A3.1.3 Proportion of land area, excretal nitrogen (N) and nitrous oxide (N₂O) emissions by hill slope category for sheep, beef cattle and deer farms in 2021



A3.2 Supplementary information for the LULUCF sector

A3.2.1 Land use mapping methodology

Areas of land use and land-use change between 1990 and 2021 are based on four wall-to-wall land use maps derived from satellite imagery at nominal mapping dates of 31 December 1989, 31 December 2007, 31 December 2012 and 31 December 2016. Area information from these maps is interpolated and extrapolated to obtain a complete time series of land-use change occurring between 1990 and 2021.

Satellite image acquisition and pre-processing

Each of the national land use maps is based on a collection of either Landsat, SPOT² or Sentinel-2 satellite imagery acquired over the summer periods (October to March) as described in table A3.2.1. Acquisition is limited to the summer months, because a high sun angle is required to reduce shadowing and increase the dynamic range of the signal received from the ground.

Table A3.2.1 Satellite imagery used for land use mapping in 1990, 2008, 2012 and 2016

Land use map	Satellite imagery	Resolution (metres)	Acquisition period
1990	Landsat 4 and Landsat 5	30	November 1988 – February 1993
2008	SPOT 5	10	November 2006 – April 2008
2012	SPOT 5	10	October 2011 – March 2013
2016	Sentinel-2	10	October 2016 – March 2017

All the imagery was orthorectified and atmospherically corrected and then standardised for spectral reflectance using the Ecosat algorithms documented in Dymond et al. (2001), Shepherd and Dymond (2003) and Dymond and Shepherd (2004). This standardisation process removes the effect of terrain slope from the imagery, effectively ‘flattening’ it, so that individual land cover types are a more consistent colour across the whole image. By minimising the effects of terrain, a more accurate and consistent classification of land use is possible. This is particularly important in New Zealand, due to the extensive areas of steep terrain.

The final step in image preparation was the mosaicking of the satellite image scenes into a seamless national image. To minimise the effect of cloud and cloud shadows in the mosaic, cloud masks were generated for each scene. These masks were then used to prioritise the order of inclusion of each scene in the mosaic to obtain a near cloud-free image of New Zealand at each mapping date.

Creating the first two land use maps: 1990 and 2008

Mapping approach

The 1990 and 2008 land use maps were created using a common mapping approach based on difference detection from an intermediate reference land-cover layer that was derived from Landsat 7 ETM+ imagery acquired in 2000 and 2001. A semi-automated approach was used to

² From French ‘Satellite pour l’Observation de la Terre’.

classify woody land cover³ in the 1990 and 2008 image mosaics. These layers were then differenced from the 2001 reference layer to create a 1990 to 2001 potential woody change layer and a 2001 to 2008 potential woody change layer.

The potential woody change layers were visually checked to confirm change and then the changes were combined with the 2001 reference layer to create the 1990 and 2008 woody land cover layers. By using this approach, it was possible to obtain a consistent resolution of change detection even though there was a significant difference between the resolutions of the source imagery at the two mapping dates: 30 metres at 1990 versus 10 metres at 2008.


Area and proximity rules were used to convert these layers from woody land cover to woody land use, making allowances for unstocked areas within forest extents and areas of regenerating vegetation in a forest context. This process is described in Shepherd and Newsome (unpublished(b)).

To determine the spatial location of the other land uses as at 1990 and 2008, information from two Land Cover Databases, LCDB1 (1996) and LCDB2 (2001) (Thompson et al., 2004), hydrological data from Land Information New Zealand (a government agency) and the New Zealand Land Resource Inventory (NZLRI) (Eyles, 1977) were used (Shepherd and Newsome, unpublished(a)).

The NZLRI database defined the area of high and low producing grassland. Areas tagged as 'improved pasture' in the NZLRI vegetation records were classified as high producing grassland in the land use maps. All other areas were classified as low producing grassland. Figure A3.2.1 illustrates this mapping process.

³ Land cover consistent with pre-1990 natural forest, pre-1990 planted forest, post-1989 forest and grassland with woody biomass land uses.

Figure A3.2.1 New Zealand's land use mapping process for 1990 and 2008 land use maps



Note: LINZ = Toitū Te Whenua Land Information New Zealand.

An interpretation guide for automated and visual interpretation of satellite imagery was prepared and used to ensure a consistent basis for all mapping processes (Ministry for the Environment, 2012). During the mapping process, independent quality control checks were performed to ensure consistent image interpretation. This involved an independent agency looking at randomly selected points across New Zealand and using the same data as the original operator to decide within what land use the point fell. The two operators were in agreement at least 95 per cent of the time. This is described in more detail in Joyce (unpublished).

Decision process for mapping post-1989 forests

The use of remotely sensed imagery has some limitations, in particular, a limited ability to map planted forest of less than three years of age. Where trees are planted within three years of the image acquisition date, they (and their surrounding vegetation) are unlikely to show a distinguishable spectral signature in satellite imagery. This occurs particularly with coarse-resolution (30 metres) Landsat 4 and 5 imagery captured around 1990. This situation is compounded by the lack of ancillary data at 1990 to support land use classification decisions. However, since 2009, the New Zealand Emissions Trading Scheme (NZ ETS) has provided valuable spatial information that has been used to confirm 1990 forest land use classifications.


Owners of post-1989 forest may apply to lodge their forests within the NZ ETS to obtain credit for increases in carbon stock since 1 January 2008. Mapping received by Te Uru Rākau – Ministry for Primary Industries for these applications is used to improve the Land Use and Carbon Analysis System (LUCAS) land use maps.

Mapping from the NZ ETS (and other post-1989 forestry schemes) has also provided a significant source of planting date information, which helps determine the correct classification of planted forest. The Forestry Allocation Plan, which forms part of the NZ ETS, compensates private owners of pre-1990 planted forest for the loss in land value arising from the introduction of penalties for deforesting pre-1990 forest land. Forest owners must apply for this compensation, providing detailed mapping and evidence of their forest planting date. These mapping data are used regularly to improve the classification accuracy of the LUCAS land use maps.

To help the decision-making process, nationwide cloud-free 1996 SPOT and 2001 Landsat 7 satellite image mosaics are also used to determine the age of forests that have been planted within two to three years of 1990. Figure A3.2.2 shows how mapping operators use the spectral signature in later imagery and ancillary information to determine the status of an area of planted forest established around 1990.

Where possible, information obtained directly from forest owners and the national planted forest plot network is also used to improve the accuracy of the pre-1990 and post-1989 forest classification.

Figure A3.2.2 Identification of post-1989 forest in New Zealand




Images:	1990 Landsat 4 (top left) 1996 SPOT 2 (top right) 2000 Landsat 7 ETM+ (bottom left) 2008 SPOT 5 (bottom right)
Location:	2,017,800, 5,730,677 (NZTM)
1990 land use:	Low producing grassland
2008 land use:	Post-1989 forest
Explanation:	In the Landsat 1990 imagery acquired on 2 December 1990, there is little evidence of the forest within the blue box that is clearly apparent in later imagery. The strength of the spectral response in the SPOT 1996 imagery suggests that the forest must have been planted near to 1990. Final confirmation of the planting date is provided via the NZ ETS application (delineated in green in the 2008 imagery), which states that the forest was planted in 1990 and, therefore, is classed as a post-1989 forest.

Adding land use maps to the time series: 2012 and 2016 land use maps

The 2012 and 2016 land use maps were created by detecting change between satellite imagery acquired for each mapping year (2008, 2012 and 2016) (Newsome et al., 2013; 2018). The 2012 map was created by using the 2008 map as a starting point and mapping in all the change detected between 2008 and 2012. Similarly, the 2012 map was used as a starting point for the 2016 map, with all areas of change detected between 2012 and 2016 mapped in to the 2012 map to create a snapshot of land use as at 2016. Figure A3.2.3 illustrates this mapping process.

Figure A3.2.3 New Zealand’s land use mapping process for 2012 and 2016 land use maps



A multi-date image segmentation process was used to identify areas of potential change. This process is described in Shepherd et al. (2019). These areas of potential change were confirmed using two separate approaches: one for areas mapped as non-forest at the start of the period and one for areas mapped as forest at the start of the period. These approaches are discussed further in the subsequent sections.

Mapping approach: non-forest areas

Potential changes in areas mapped as non-forest were manually checked in the satellite imagery to determine whether a land-use change had occurred since the previous land use map. Operators used the 2008 and 2012 SPOT imagery and 2016 Sentinel-2 imagery, along with other imagery data sets as listed in table A3.2.2, to establish whether land-use change had occurred. Once change was confirmed, the area of change was delineated in the land use map.

Table A3.2.2 Ancillary mosaicked imagery data sets used in land use mapping

Satellite imagery	Resolution (m)	Coverage	Acquisition period
Landsat 7	30	North Island, South Island and Stewart Island	September 1999 – February 2003 October 2011 – February 2012 October 2012 – March 2013
SPOT maps products	2.5 and 1.5	North Island, South Island and Stewart Island	January 2008 – June 2009 October 2012 – April 2014
Disaster Monitoring Constellation	22	North Island, South Island and Stewart Island	November 2009 – March 2010
SPOT 5	10	Four priority areas: Northland, Waikato, Marlborough and Southland	October 2010 – March 2011
Landsat 8	30	North Island, South Island and Stewart Island	November 2013 – February 2014 October 2014 – March 2015 October 2015 – March 2016
Sentinel-2	10	North Island, South Island and Stewart Island	November 2015 – March 2016
Aerial photography	Variable	All of North Island and Stewart Island and most of South Island	Various

As part of the 2016 mapping process, high and low producing grassland classes were mapped at 2008, 2012 and 2016 using a data fusion technique described in Manderson et al. (2018). This technique brought together a range of biophysical and land use data sets to create a probability map for high producing grassland at each mapping date. This map was used to classify grassland into high and low producing areas in the 2016 land use map and back-correct the 2012 and 2008 maps to maintain time-series consistency. Before the 2016 map, grassland areas had been split into high-producing and low-producing based on the mapping of high-producing grassland in the NZLRI, which was completed in the mid-1980s. No changes between grassland subcategories had been mapped throughout the time series, and any change to grassland from other land uses were classified into low or high producing grassland based on imagery and context. The 1990 land use map was assumed to contain a fair representation of the split between high and low producing grassland, based on the original mapping of this data set using the NZLRI as described above.

Mapping approach: forest areas

Areas of potential change within the forest extent were considered to be potential destocking.⁴ These areas were first screened, to ensure they represented actual change, as opposed to false change related to cloud contamination or image misregistration.

The next step was to determine which areas of destocking represented land-use change (deforestation) as opposed to temporary forest loss (e.g., harvesting activity occurring as part of ongoing forestry land use).

Where possible, areas of destocking were first checked in pre-existing aerial orthophotography to determine whether replanting may have occurred. Cases of replanting were then classified as 'harvested' and excluded from further consideration.

Because it is rarely possible to determine whether deforestation has occurred using currently available satellite imagery alone, high-resolution vertical or oblique aerial photography is necessary to provide a detailed view of land use activity occurring subsequently on the ground.

All remaining unclassified areas of destocking were field checked by obtaining aerial photography over each site.

Based on the aerial photographic evidence and supplemental deforestation data from the NZ ETS, each area was given one of the following destock classifications.

- Harvested: the area shows evidence of ongoing forestry land use such as replanting, preparation for planting or a context consistent with replanting, such as being surrounded by plantation forestry.
- Harvested and converted: the forest stand is registered in the NZ ETS using the Carbon Equivalent Forest option to harvest but replanted in a different location.
- Deforested: the area shows evidence of land-use change, such as the removal of stumps, pasture establishment, fencing and stock, or earthworks.

⁴ 'Destocking' is defined here as forest loss for any reason including harvesting, deforestation or some type of non-anthropogenic change, such as wind damage or erosion.

- Awaiting: the area has been destocked for less than four years⁵ and/or there is no clear evidence of land-use change or replanting. That is, the area is lying fallow or, in the case of natural forest areas, the vegetation has been sprayed but not cleared.⁶
- No change: the area has not been sufficiently destocked and was incorrectly identified as meaningful change (may include thinning activity).
- Never forest: the area in fact did not meet the forest definition at the beginning of the change period. These areas required correction to a non-forest land use in the land use map from the beginning of the change period.
- Non-anthropogenic change: destocking was not directly human induced – for example, erosion – and there has been no land-use change.


For each deforested area, further information was then recorded, such as the year in which the deforestation occurred. This was determined by examining the ancillary imagery data sets listed in table A3.2.2. Figure A3.2.4 shows the process of confirming deforestation and establishing the year in which it occurred. Further information on the mapping of forest change can be found in Indufor Asia Pacific (2018).

The final step in the 2012 and 2016 land use mapping process was to add the confirmed areas of deforestation into the land use map.

⁵ To distinguish between deforestation and temporary tree crown cover removal in forest land, New Zealand has defined the expected period between the removal of tree cover and successful natural regeneration or planting as four years.

⁶ Often regenerating shrubland areas are sprayed but land use conversion is not completed by clearing the area. In these instances, the vegetation regenerates and recovers, therefore, land-use change has not occurred.

Figure A3.2.4 New Zealand's identification of deforestation



Note: DMC = Disaster Monitoring Constellation.

Quality assurance/quality control (QA/QC) and verification

During the mapping process, the 1990, 2008, 2012 and 2016 land use maps were checked to determine that the mapping was consistent with the satellite image classification specification set out in *Land Use and Carbon Analysis System: Satellite imagery interpretation guide for land use classes* (Ministry for the Environment, 2012).

The quality-control checks performed on the 1990 and 2008 land use maps included checking around 28,000 randomly selected points in areas mapped as forest and grassland with woody biomass. These were evaluated by independent assessors. In this exercise, independent assessors agreed with the original classification 91 per cent of the time. Where there was disagreement, the points were recorded in a register and this was used to plan improvements to the 1990 and 2008 land use maps. These improvements were subsequently completed.

Two distinct quality-control checks were performed on the 2012 land use map. The first of these checked every polygon where land-use change had occurred from a non-forest land use between 2008 and 2012. The acceptance criterion for this check was that the land use classification had to be correct at both mapping dates at least 90 per cent of the time. This means that the land use, both at the start of the land-use change event and at the end of the land-use change event, had to be correct. The second quality-control measure was to check the accuracy of destock detection in areas that were in a forest land use at 2008. Sampling for this check was designed to test that at least 90 per cent of the destocking had been detected at the 95 per cent confidence level. Checks were completed on each of the 16 regions of New Zealand individually and all regions passed. During this process, 14,443 points were checked.

Quality-control checking for the 2016 land use map was carried out region-by-region looking at all areas of expected change (based on mapping targets sent to the mapping supplier) and actual change supplied in the map. Checks were also made for invalid change, for example, a pre-1990 planted forest cannot change to a post-1989 forest. Spatial checks were performed to ensure that the integrity of the map had been maintained. These included checking for gaps and overlaps as well as that the total area of the map had not changed.

Each mapping improvement activity carried out on the 1990, 2008, 2012 and 2016 maps has been subjected to quality-assurance checks, to ensure accuracy and consistency. Quality assurance strategies have been tailored to each improvement activity, usually including a combination of random sampling of updated areas and analysis of the changes in land use areas.

The approach used to implement quality-assurance processes is documented in the LUCAS Data Quality Framework (PricewaterhouseCoopers, unpublished).

Uncertainties and time-series consistency

In 2014, an accuracy assessment was completed for the 2012 land use map. A stratified random sample of 2,000 points was made, and the land use classification was independently assessed at each point location. SPOT-6 natural colour 1.5-metre resolution imagery was used as the reference data source. This imagery met the criteria for a reference data source, having better resolution than the SPOT-5 10-metre resolution imagery used to create the 2012 land use map, and being acquired over a similar period.⁷

The overall map accuracy was found to be 95.2 per cent (Poyry Management Consulting (NZ) Ltd, unpublished). The user and producer accuracies for the three forest classes were all over 94 per cent. For all forest classes, the total mapped area fell within the 95 per cent confidence interval of the total class area as determined by the accuracy assessment.

Non-forest land uses generally had user and producer accuracies of over 90 per cent. Exceptions were the *Wetlands* and *Grassland with woody biomass* categories, for which producer accuracies were 85 per cent and 60 per cent respectively (Poyry Management Consulting (NZ) Ltd, unpublished). The *Wetlands* category was slightly under-mapped. This is because vegetated wetland and *Grassland with woody biomass* are sometimes difficult to distinguish in imagery where the extent of flooding varies seasonally. *Grassland with woody biomass* appears to be more substantially under-mapped, with accuracy assessment operators identifying areas of high and low producing grassland that should have been mapped as

⁷ The SPOT-6 natural colour 1.5-metre resolution imagery was acquired in the summers of 2012/13 and 2013/14 making it generally one year later than the SPOT-5 multi-spectral 10-metre resolution imagery used to create the 2012 land use map.

Grassland with woody biomass. This is also a difficult judgement call, because the boundary between areas of low producing and high producing grassland and *Grassland with woody biomass* can be hard to define and can shift with grazing.

A3.2.2 Annual land-use change

Annual land-use change areas are interpolated and extrapolated from the four national land use maps using a number of supporting data sets to inform the trends occurring between the wall-to-wall mapping dates of 1990, 2008, 2012 and 2016.

Land-use change before 1990

Data from a variety of sources were used to determine land areas before 1990. Data sources suitable for determining land use at a national level typically comprise one of the following:

- maps or scaled images depicting land use or proxies for land use (e.g., a ‘map of forest areas’)
- tabulated land use area data collected for an administrative area (e.g., county, district or region)
- production sector (e.g., the area of orchard crops).

This methodology was peer reviewed by Hunter and McNeill (unpublished), who provided independent subject-matter expertise. They noted that the methodology was sound, and the choice of historical data sets was reasonable. They judged that the method reasonably met the standards of the IPCC Guidelines for National Greenhouse Gas Inventories (IPCC, 2006a).

Annual land-use changes from 1990 to 2007

Annual land-use changes from 1990 to 2007 are interpolated between the 1990 and 2008 land use maps, which provide the total area of change over that period. Most of the land-use changes are interpolated linearly between mapping dates; however, some of the land-use changes make use of surrogate data sets to better reflect land-use change trends within this period. This approach follows methodology outlined in section 3.3.1 of the 2006 IPCC Guidelines (IPCC, 2006a).

The surrogate data sets used between 1 January 1990 and 31 December 2007 are as follows.

- Deforestation trends between 1990 and 1 January 2008 for pre-1990 planted forest and post-1989 forest are based on the 2008 Deforestation Intentions Survey (Manley, 2009) and unpublished work by Scion (the New Zealand Forest Research Institute). The work by Scion is referred to in Wakelin (unpublished(c))
- Afforestation trends for post-1989 planted forest are based on estimates from the National Exotic Forest Description (NEFD) (Ministry for Primary Industries, 2021)
- Afforestation trends for post-1989 natural forest are based on the plot analysis in Paul et al. (unpublished(b)). The age of vegetation on plots was used to estimate the year afforestation occurred. Afforestation area was then assigned annually by taking the number of new post-1989 natural forest plots per year (estimated using a five-year rolling average) as a proportion of the total number of post-1989 natural forest plots in 2007 and multiplying by the mapped area of post-1989 natural forest in 2007.

Annual land-use changes from 2008 to 2016

Annual land-use changes from 2008 to 2016 are generally linearly interpolated between the 2008, 2012 and 2016 land use maps. The only exceptions to this are:

- deforestation occurring between 2008 and 2016, which is mapped
- afforestation, which uses a mixture of mapped and surveyed data as detailed in table A3.2.3. This is because not all new planting will have been detected in satellite imagery and mapped into the 2016 map yet. New planting can take up to four years to be visible in satellite imagery, therefore, afforestation mapping up to 2016 will not be finalised until 2020.

Table A3.2.3 Methods used to estimate Afforestation total area and trends between 2008 and 2016

Afforestation type	Reporting years: 2008 to 2012		Reporting years: 2013 to 2016	
	Estimate of total afforestation for the period	Trend in afforestation within the period	Estimate of total afforestation for the period	Trend in afforestation within the period
Post-1989 planted forest	Based on afforestation mapped between 2008 and 2012	Based on new planting data from national survey (National Exotic Forest Description)	Based on afforestation mapped between 2013 and 2016	Based on new planting data from national survey (National Exotic Forest Description)
Post-1989 natural forest	Based on afforestation mapped between 2008 and 2012	Linear interpolation	Based on afforestation mapped between 2013 and 2016	Linear interpolation

Estimating land-use change for 2017 to 2021

Activity data for the five most recent years of this inventory from 2017 to 2021, have been estimated from a survey for afforestation (Ministry for Primary Industries, 2021) and extrapolated from the most recent mapping for all other land-use changes.

Deforestation

Annual deforestation has now been mapped for the years from 2017 to 2019 inclusive. The approach used for mapping deforestation in these years is similar to the approach used in the period of 2008 to 2016. Forest clearing is first detected through automated change detection in annual national Sentinel-2 satellite imagery mosaics. These areas are then overflowed to capture 0.3-metre resolution aerial imagery over each area of interest. Machine learning techniques are used on this aerial imagery to identify the land cover present at each site (Lynker Analytics Consortium, 2020). The combination of land covers is used to infer the possible land use of the site. For example, the presence of newly planted seedlings implies ongoing forestry land use and hence confirms harvesting rather than deforestation activity, whereas the presence of grassland implies deforestation to a grazing land use. These inferred land uses are visually checked to confirm where deforestation has occurred.

The estimated area of deforestation of pre-1990 planted forest and post-1989 planted forest occurring during 2020 and 2021 has been extrapolated from the annual mapped deforestation over the period 2013 to 2019. Over this period there has been a downward trend in both the total annual planted forest deforestation and the annual pre-1990 planted forest deforestation. This downward trend in deforestation is in response to the increasing domestic price of carbon and therefore increased cost associated with deforestation.

The estimated area of deforestation of pre-1990 natural forest and post-1989 natural forest occurring in 2020 and 2021 has been calculated as the average of the three previous mapped

years. Natural forest deforestation has occurred at relatively low levels in recent years but there is not sufficient evidence of a downward trend in the data to justify using the extrapolation approach employed for the planted forest classes. Natural forest deforestation is not subject to the same carbon price liabilities as planted forest deforestation although there are other environmental constraints to clearing this type of forest.

The destination land use for areas of estimated deforestation has been pro-rated based on the mapped destination land uses of deforestation occurring in the period 2017 to 2019. Only the major destination land uses were included in this pro-rating process. Major destination land uses were considered to be those that were consistently reported for each year across the whole mapped period for all types of forest loss. This reduced the destination land uses from 12 to five classes: *High producing grassland*, *Low producing grassland*, *Grassland with woody biomass*, *Settlements* and *Other land*.

Afforestation

The annual area of afforestation of post-1989 planted forest for 2017 to 2021 is based on estimates from the NEFD (Ministry for Primary Industries, 2021). The annual area of afforestation of post-1989 natural forest for 2017 to 2019 is estimated from the Ministry for Primary Industries afforestation scheme data. The area of post-1989 natural afforestation for 2020 and 2021 is estimated from the Afforestation and Deforestation Intentions Survey for 2020 by taking the total area of 'natural reversion' and 'indigenous tall planted' (Manley, 2022). For post-1989 natural forest dominated by wilding exotic conifers, a linear extrapolation of the mapped area of land use change between 2012 and 2016 (for this forest type) was used to estimate afforestation for 2017 to 2021.

The land use before afforestation has been pro-rated across all non-forest land uses in the same proportions as for post-1989 afforestation that has been mapped between 2012 and 2016.

Other land-use changes

All other land-use changes for 2017 to 2021 have been linearly extrapolated from the changes mapped between 2012 and 2016.

Uncertainties and time-series consistency

Time-series consistency is maintained by using a combination of linear interpolation and extrapolation between mapping dates, and from the last mapping date, as described in section 5.3 of volume 1 of the 2006 IPCC General Guidance and Reporting (IPCC, 2006b).

It is difficult to quantify the uncertainty introduced by the interpolation and extrapolation process. The error introduced by extrapolation from the last mapping date depends on how consistent the rate of change in land use is between the mapped period, which is used to establish the trend, and the extrapolated period.

When New Zealand introduced the 2016 land use map into the reporting cycle for the Inventory submitted in 2019, replacing 2013 to 2016 extrapolated activity data with interpolated data with a mapped end point at 2016, an emission reduction of 9 per cent was reported as the recalculation for 2016. This recalculation also included other updates, however, it is not substantially different to the recalculations reported in other years, indicating that the error introduced by extrapolation is unlikely to be large.

A3.2.3 Annual land-use change summary

This section contains a summary of the annual land-use change from 1990 to 2021 (table A3.2.4).

Table A3.2.4 Annual land-use changes (units in 000s hectares)

From Pre-1990 natural forest, to...	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Natural forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pre-1990 planted forest	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4
Post-1989 forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Post-1989 natural regenerating forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Cropland – annual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cropland – perennial	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – high producing	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
Grassland – low producing	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9
Grassland – with woody biomass	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Wetland – open water	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – vegetative non-forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Settlements	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other land	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
From Pre-1990 planted forest, to...																
Natural forest	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Pre-1990 planted forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Post-1989 forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Post-1989 natural regenerating forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Cropland – annual	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.0	0.0	0.0	0.1
Cropland – perennial	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.0	0.0	0.0	0.1
Grassland – high producing	-	-	-	-	-	-	-	-	-	-	1.8	1.7	1.3	2.4	5.1	9.9
Grassland – low producing	-	-	-	-	-	-	-	-	-	-	0.6	0.5	0.5	0.7	1.1	1.7
Grassland – with woody biomass	-	-	-	-	-	-	-	-	-	-	0.3	0.3	0.3	0.4	0.7	0.9
Wetland – open water	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.0	0.0	0.0	0.0

From Pre-1990 natural forest, to...	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Wetland – vegetative non-forest	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.0	0.0	0.0	0.0
Settlements	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.0	0.0	0.1	0.1
Other land	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.0	0.0	0.0	0.1
From Post-1989 planted forest, to...																
Natural forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pre-1990 planted forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Post-1989 forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Post-1989 natural regenerating forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Cropland – annual	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.0	0.0
Cropland – perennial	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.0	0.0
Grassland – high producing	-	-	-	-	-	-	-	-	-	-	-	-	0.5	1.5	1.3	1.5
Grassland – low producing	-	-	-	-	-	-	-	-	-	-	-	-	0.1	0.3	0.2	0.3
Grassland – with woody biomass	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.1	0.1	0.1
Wetland – open water	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.0	0.0
Wetland – vegetative non-forest	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.0	0.0
Settlements	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.0	0.0
Other land	-	-	-	-	-	-	-	-	-	-	-	-	0.0	0.0	0.0	0.0
From Post-1989 natural forest, to...																
Natural forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pre-1990 planted forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Post-1989 forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Post-1989 natural regenerating forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Cropland – annual	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Cropland – perennial	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Grassland – high producing	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Grassland – low producing	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Grassland – with woody biomass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Wetland – open water	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

From Pre-1990 natural forest, to...	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Wetland – vegetative non-forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Settlements	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Other land	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
From Cropland – annual, to...																
Natural forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pre-1990 planted forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Post-1989 forest	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Post-1989 natural regenerating forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cropland – annual	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Cropland – perennial	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Grassland – high producing	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Grassland – low producing	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – with woody biomass	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – open water	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – vegetative non-forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Settlements	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other land	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
From Cropland – perennial, to...																
Natural forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pre-1990 planted forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Post-1989 forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Post-1989 natural regenerating forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cropland – annual	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Cropland – perennial	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Grassland – high producing	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Grassland – low producing	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – with woody biomass	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – open water	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

From Pre-1990 natural forest, to...	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Wetland – vegetative non-forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Settlements	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other land	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
From Grassland – high producing, to...																
Natural forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Pre-1990 planted forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 forest	2.7	2.6	8.5	10.4	16.6	12.5	14.1	10.8	8.7	6.8	5.7	5.1	3.7	3.4	1.8	1.0
Post-1989 natural regenerating forest	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.3
Cropland – annual	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.3	1.3	1.3	1.3	1.3	1.2	1.2	1.2	1.2
Cropland – perennial	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.7	1.7	1.7	1.7	1.7	1.6	1.6	1.6	1.6
Grassland – high producing	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Grassland – low producing	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – with woody biomass	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.9	0.9	0.9	0.9	0.9	0.7	0.7	0.7	0.7
Wetland – open water	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – vegetative non-forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Settlements	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Other land	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
From Grassland – low producing, to...																
Natural forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Pre-1990 planted forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 forest	8.0	7.8	25.4	31.2	49.7	37.4	42.4	32.3	25.9	20.3	17.0	15.2	11.2	10.1	5.4	3.0
Post-1989 natural regenerating forest	0.4	0.5	0.6	0.5	0.4	0.4	0.4	1.2	1.3	1.5	1.8	2.2	1.8	2.0	2.6	3.1
Cropland – annual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cropland – perennial	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Grassland – high producing	54.7	54.7	54.7	54.7	54.7	54.7	54.7	56.1	56.1	56.1	56.1	56.1	54.7	54.7	54.7	54.7
Grassland – low producing	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Grassland – with woody biomass	3.3	3.3	3.3	3.3	3.3	3.3	3.3	4.9	4.9	4.9	4.9	4.9	3.3	3.3	3.3	3.3
Wetland – open water	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1

From Pre-1990 natural forest, to...	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Wetland – vegetative non-forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Settlements	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Other land	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
From Grassland – with woody biomass, to...																
Natural forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Pre-1990 planted forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 forest	2.8	2.8	9.0	11.1	17.7	13.3	15.0	11.5	9.2	7.2	6.0	5.4	4.0	3.6	1.9	1.1
Post-1989 natural regenerating forest	0.5	0.6	0.7	0.6	0.5	0.4	0.4	1.3	1.5	1.6	2.0	2.5	2.0	2.2	2.9	3.4
Cropland – annual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cropland – perennial	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – high producing	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.3	1.3	1.3	1.3	1.3	1.1	1.1	1.1	1.1
Grassland – low producing	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.7	1.7	1.7	1.7	1.7	1.2	1.2	1.2	1.2
Grassland – with woody biomass	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Wetland – open water	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – vegetative non-forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Settlements	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other land	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
From Wetland – open water, to...																
Natural forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Pre-1990 planted forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Post-1989 natural regenerating forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – annual	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – perennial	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – high producing	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – low producing	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – with woody biomass	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – open water	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–

From Pre-1990 natural forest, to...	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Wetland – vegetative non-forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Settlements	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other land	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
From Wetland – vegetative non-forest, to...																
Natural forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Pre-1990 planted forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 forest	0.0	0.0	0.1	0.1	0.2	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0
Post-1989 natural regenerating forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cropland – annual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cropland – perennial	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – high producing	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Grassland – low producing	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Grassland – with woody biomass	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – open water	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – vegetative non-forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Settlements	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other land	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
From Settlements, to...																
Natural forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Pre-1990 planted forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Post-1989 natural regenerating forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cropland – annual	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – perennial	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – high producing	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – low producing	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Grassland – with woody biomass	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – open water	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

From Pre-1990 natural forest, to...	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Wetland – vegetative non-forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Settlements	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Other land	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
From Other land, to...																
Natural forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Pre-1990 planted forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 forest	0.1	0.1	0.2	0.3	0.4	0.3	0.3	0.3	0.2	0.2	0.1	0.1	0.1	0.1	0.0	0.0
Post-1989 natural regenerating forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.1
Cropland – annual	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – perennial	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Grassland – high producing	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – low producing	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – with woody biomass	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Wetland – open water	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – vegetative non-forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Settlements	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other land	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–

From Pre-1990 natural forest, to...	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Natural forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Pre-1990 planted forest	1.4	1.4	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Post-1989 forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 natural regenerating forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – annual	0.0	0.0	–	–	0.0	–	–	0.0	0.0	0.0	–	0.0	–	–	–	–
Cropland – perennial	0.0	0.0	–	–	–	–	–	–	–	–	–	–	–	0.0	–	–
Grassland – high producing	0.6	0.6	0.3	0.6	0.5	0.2	0.1	0.2	0.1	0.2	0.1	0.1	0.2	0.1	0.2	0.2
Grassland – low producing	0.9	0.9	0.3	0.8	0.5	0.3	0.3	0.5	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.2
Grassland – with woody biomass	0.3	0.3	0.2	1.0	0.9	0.5	0.6	0.4	0.2	0.5	0.3	0.2	0.1	0.3	0.2	0.2

From Pre-1990 natural forest, to...	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Wetland – open water	0.0	0.0	–	0.0	–	0.0	–	–	–	0.0	–	–	–	–	–	–
Wetland – vegetative non-forest	0.0	0.0	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Settlements	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other land	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0
From Pre-1990 planted forest, to...																
Natural forest	0.1	0.1	0.4	0.4	0.4	0.4	0.4	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Pre-1990 planted forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 natural regenerating forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – annual	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	–	–
Cropland – perennial	0.1	0.1	–	0.0	–	–	–	–	0.0	–	0.0	–	0.0	0.0	–	–
Grassland – high producing	12.4	16.5	2.4	3.2	3.5	2.6	4.6	5.8	4.0	2.6	2.4	2.0	1.3	0.6	0.5	0.3
Grassland – low producing	2.1	2.8	1.0	1.3	2.2	1.9	2.4	3.1	2.5	1.7	1.9	0.9	0.4	0.4	0.3	0.1
Grassland – with woody biomass	1.1	1.5	0.3	1.0	0.7	0.6	0.5	0.7	0.4	0.7	0.5	0.5	0.2	0.2	0.1	0.1
Wetland – open water	0.0	0.0	0.0	0.0	0.0	0.0	–	0.0	0.0	0.0	0.0	–	–	–	–	–
Wetland – vegetative non-forest	0.0	0.0	–	0.0	0.0	0.0	–	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–
Settlements	0.2	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other land	0.1	0.1	0.1	0.0	0.0	0.1	0.1	0.1	0.1	0.2	0.2	0.0	0.0	0.0	0.0	0.0
From Post-1989 planted forest, to...																
Natural forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Pre-1990 planted forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 natural regenerating forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – annual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	–	–
Cropland – perennial	0.0	0.0	–	0.0	–	–	–	–	–	–	–	–	0.0	0.0	–	–
Grassland – high producing	1.3	3.1	0.9	0.9	1.0	1.2	0.8	1.6	1.3	1.6	1.8	1.7	2.2	1.1	1.0	0.7
Grassland – low producing	0.2	0.5	0.2	0.9	0.5	0.8	0.4	0.8	0.7	0.7	1.2	1.8	1.3	0.6	0.7	0.5
Grassland – with woody biomass	0.1	0.3	0.1	0.2	0.2	0.1	0.4	0.3	1.1	0.4	0.5	0.7	0.5	0.7	0.4	0.3

From Pre-1990 natural forest, to...	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Wetland – open water	0.0	0.0	0.0	–	–	0.0	0.0	0.0	–	0.0	–	–	–	–	–	–
Wetland – vegetative non-forest	0.0	0.0	–	–	0.0	–	–	–	0.0	–	0.0	0.0	0.0	–	–	–
Settlements	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	0.0	0.0	0.0	0.0
Other land	0.0	0.0	–	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0
From Post-1989 natural forest, to...																
Natural forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Pre-1990 planted forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 natural regenerating forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – annual	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – perennial	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Grassland – high producing	–	–	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0
Grassland – low producing	–	–	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – with woody biomass	–	–	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0
Wetland – open water	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Wetland – vegetative non-forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Settlements	–	–	–	–	–	–	–	–	–	0.0	0.0	0.0	–	0.0	0.0	0.0
Other land	–	–	–	–	0.0	–	–	–	–	–	–	0.0	–	0.0	0.0	0.0
From Cropland – annual, to...																
Natural forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Pre-1990 planted forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–	–	–	–	–
Post-1989 natural regenerating forest	0.0	0.0	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – annual	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – perennial	0.3	0.3	0.1	0.1	0.1	0.1	0.1	–	–	–	–	–	–	–	–	–
Grassland – high producing	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–
Grassland – low producing	0.0	0.0	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Grassland – with woody biomass	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–	–	–	–	–

From Pre-1990 natural forest, to...	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Wetland – open water	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – vegetative non-forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Settlements	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0
Other land	0.0	0.0	–	–	–	–	–	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
From Cropland – perennial, to...																
Natural forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Pre-1990 planted forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 forest	0.0	0.0	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 natural regenerating forest	0.0	0.0	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – annual	0.1	0.1	0.2	0.2	0.2	0.2	0.2	–	–	–	–	–	–	–	–	–
Cropland – perennial	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Grassland – high producing	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – low producing	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–	–	–	–	–
Grassland – with woody biomass	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–	–	–	–	–
Wetland – open water	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – vegetative non-forest	0.0	0.0	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Settlements	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other land	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
From Grassland – high producing, to...																
Natural forest	–	–	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–
Pre-1990 planted forest	–	–	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–
Post-1989 forest	0.4	0.4	1.1	1.6	2.0	3.2	3.1	1.0	0.8	0.8	0.6	0.6	0.7	1.8	3.4	4.5
Post-1989 natural regenerating forest	0.3	0.3	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.2	0.2	0.2	0.2	0.2
Cropland – annual	1.2	1.2	0.0	0.0	0.0	0.0	0.0	–	–	–	0.0	–	–	–	–	–
Cropland – perennial	1.6	1.6	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Grassland – high producing	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Grassland – low producing	0.0	0.0	0.6	0.6	0.6	0.6	0.6	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2
Grassland – with woody biomass	0.7	0.7	0.6	0.6	0.6	0.6	0.6	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0

From Pre-1990 natural forest, to...	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Wetland – open water	0.0	0.0	0.3	0.3	0.3	0.3	0.3	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Wetland – vegetative non-forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Settlements	0.8	0.8	0.5	0.5	0.5	0.5	0.5	0.8	0.8	0.8	0.8	0.7	0.7	0.7	0.7	0.7
Other land	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
From Grassland – low producing, to...																
Natural forest	–	–	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–
Pre-1990 planted forest	–	–	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.0	–	–	–	–
Post-1989 forest	1.3	1.2	1.8	3.9	5.4	10.7	10.3	4.4	3.1	3.1	1.6	4.5	5.9	16.1	28.8	38.1
Post-1989 natural regenerating forest	3.5	3.3	0.7	0.7	0.7	0.7	0.7	1.5	1.5	1.5	1.5	3.0	2.5	2.7	2.8	3.2
Cropland – annual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–
Cropland – perennial	0.2	0.2	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–	–	–	–	–
Grassland – high producing	54.7	54.7	7.5	7.5	7.5	7.5	7.5	12.3	12.3	12.3	12.3	12.1	12.0	12.0	12.0	12.0
Grassland – low producing	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Grassland – with woody biomass	3.3	3.3	1.9	1.9	1.9	1.9	1.9	0.4	0.4	0.4	0.5	0.3	0.3	0.3	0.3	0.3
Wetland – open water	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – vegetative non-forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Settlements	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other land	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0
From Grassland – with woody biomass, to...																
Natural forest	–	–	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	–	–	–	–	–
Pre-1990 planted forest	–	–	0.0	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	–	–	–	–	–
Post-1989 forest	0.5	0.4	0.7	1.2	1.5	2.7	2.6	1.0	0.8	0.8	0.3	1.4	1.0	3.3	5.8	7.7
Post-1989 natural regenerating forest	3.9	3.7	1.0	1.0	1.0	1.0	1.0	0.5	0.5	0.5	0.5	0.6	0.5	0.5	0.6	0.6
Cropland – annual	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cropland – perennial	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – high producing	1.1	1.1	1.4	1.4	1.4	1.4	1.4	0.8	0.8	0.9	0.8	0.6	0.5	0.5	0.5	0.5
Grassland – low producing	1.2	1.2	2.5	2.5	2.5	2.5	2.5	1.7	1.8	1.7	1.7	1.4	1.2	1.2	1.2	1.2
Grassland – with woody biomass	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–

From Pre-1990 natural forest, to...	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Wetland – open water	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – vegetative non-forest	0.0	0.0	–	–	–	–	–	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Settlements	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other land	0.0	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
From Wetland – open water, to...																
Natural forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Pre-1990 planted forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Post-1989 forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–	–	–	–	–
Post-1989 natural regenerating forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – annual	–	–	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–	–	–	–	–
Cropland – perennial	0.0	0.0	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Grassland – high producing	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – low producing	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – with woody biomass	0.0	0.0	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Wetland – open water	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Wetland – vegetative non-forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–	–	–	–	–
Settlements	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–
Other land	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–
From Wetland – vegetative non forest, to...																
Natural forest	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Pre-1990 planted forest	–	–	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–	–	–	–	–
Post-1989 forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Post-1989 natural regenerating forest	0.0	0.0	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – annual	0.0	0.0	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Cropland – perennial	0.0	0.0	–	–	–	–	–	–	–	–	–	–	–	–	–	–
Grassland – high producing	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Grassland – low producing	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – with woody biomass	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	–	–	–	–	–

From Pre-1990 natural forest, to...	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Wetland – open water	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – vegetative non-forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Settlements	0.0	0.0	-	-	-	-	-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other land	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
From Settlements, to...																
Natural forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pre-1990 planted forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Post-1989 forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-
Post-1989 natural regenerating forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-
Cropland – annual	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Cropland – perennial	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Grassland – high producing	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Grassland – low producing	-	-	-	-	-	-	-	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Grassland – with woody biomass	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Wetland – open water	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-
Wetland – vegetative non-forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Settlements	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Other land	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-
From Other land, to...																
Natural forest	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pre-1990 planted forest	-	-	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-	-	-	-	-
Post-1989 forest	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Post-1989 natural regenerating forest	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cropland – annual	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Cropland – perennial	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Grassland – high producing	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	-	-	-	-	-
Grassland – low producing	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grassland – with woody biomass	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

From Pre-1990 natural forest, to...	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Wetland – open water	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Wetland – vegetative non-forest	0.0	0.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Settlements	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other land	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

A3.2.4 Soils methodology

New Zealand uses a Tier 2 method to estimate soil carbon changes in mineral soils and follows the Tier 1 approach for organic soils.

Mineral soils

New Zealand's Tier 2 method for mineral soils involves estimating steady state soil organic carbon (SOC) stocks for each land use based on New Zealand soil data (described in more detail below). Changes in SOC stocks associated with land-use change are calculated according to the IPCC default method (IPCC, 2006a) using the equation:

$$\Delta C = [(SOC_0 - SOC_{(0-T)})/20] \times A \quad (A3.2.1)$$

Where: ΔC = change in carbon stocks (tonnes)
 SOC_0 = stable SOC stock in the inventory year (tonnes C ha⁻¹)
 $SOC_{(0-T)}$ = stable SOC stock T years prior to the inventory year (tonnes C ha⁻¹)
 A = land area of parcels with these SOC terms (hectares)
20 = IPCC default SOC stock transition period (year).

The SOC stock for each land use is characterised with country-specific data via the Soil Carbon Monitoring System (Soil CMS) model (McNeill and Barringer, unpublished; McNeill et al., 2014). The correct operation of the Soil CMS model involves fitting the model to the soil carbon data set and then using the coefficients for the different land use categories for each land use transition (equation A3.2.1). The interpretation of the different land use effects is informed by multi-comparison significance.

Characterising SOC stocks: New Zealand's Soil Carbon Monitoring System

Unbiased estimates of SOC stocks associated with each land use in New Zealand are calculated by using country-specific data in the Soil CMS model. The operation of the Soil CMS model involves applying a linear statistical model to predict SOC stocks from land use, climate and soil order, which together regulate net SOC storage. The model also includes an additional environmental factor consisting of the product of slope and rainfall (hereafter, slope × rainfall), a term used as a proxy for erosivity, the potential for surface soil erosion to occur (Giltrap et al., unpublished). This allows for the explanatory effect of the land use category on SOC stocks to be isolated from other factors that affect SOC.

Two main assumptions underpin the operation of the Soil CMS model: first, the SOC values in the sample data set represent equilibrium SOC values for each stratified soil, climate and land use cell, and erosivity index; and second, changes in land use are the key drivers of change in SOC at the decadal scale, while all other changes due to soil type, climate or erosivity are assumed to be constant (McNeill et al., 2014). The model allows for an explanatory effect by land use category, so that estimates grouped by land use are unbiased where a specific land use category has an effect significantly different from the pooled soil carbon value from all land use categories. Where a land use category is a significant explanatory variable of SOC, incorporating land use in the model reduces the overall residual standard error associated with soil carbon (McNeill and Barringer, unpublished).

Soil carbon linear parametric model

The generalised least squares model used for the Soil CMS is a minimum variance unbiased estimator (Draper and Smith, 1998). This approach is consistent with the physically based soil carbon model outlined in the literature (Baisden et al., unpublished(b); Kirschbaum et al., unpublished; Scott et al., 2002; Tate et al., 2005).

The generalised least squares regression model for soil carbon in the 0–30-centimetre layer uses explanatory variables of the soil–climate factor, the land use category and slope × rainfall. This model is represented as an equation for the soil carbon $C_{i,j}^{0-30cm}$ in land use category i and soil–climate class j as:

$$C_{i,j}^{0-30cm} = M + L_i + S_j + b.SR + \varepsilon \quad (A3.2.2)$$

Where: M = the mean soil carbon in the 0–30-centimetre layer for the combination of the reference level of land use (low producing grassland), the reference level for soil climate (MstTempHAC, i.e., ‘moist temperate high activity clay’), and level ground
 L_i = the effect of the i -th land use, specifying the difference in soil carbon relative to the reference land use (low producing grassland), in tonnes per hectare
 S_j = the effect of the j -th soil–climate class relative to the reference level
 $b.SR$ = the additional soil carbon for each unit of erosivity (slope × rainfall) (millidegree × 10^{-1})
 ε = the model uncertainty.

The quantities M , L_i , S_j , as well as the slope × rainfall coefficient $b.SR$, are obtained by fitting a statistical model to the Soil CMS calibration data set; all other quantities are obtained from other data sets or from separate analyses (McNeill and Barringer, unpublished). For example, the mean value of the slope × rainfall must be obtained from national statistics of rainfall and a terrain slope map, which has been calculated from geographic information system (GIS) layers (Giltrap et al., unpublished).

More elaborate alternatives to the model have been considered but were not found to be significantly better than the model given in equation A3.2.2 (McNeill et al., 2014).

Soil data sets

Soil data for the Soil CMS inventory model come from five sources.

Historic soils: This data set is derived primarily from the National Soils Database,⁸ with a small number of samples from various supplementary data sets; data from all sources were collected between 1935 and 2005. The National Soils Database represents soil profile data for over 1,500 soil pits scattered throughout New Zealand. These data contain the soil description following either the Soil Survey Method (Taylor and Pohlen, 1962) or *Soil Description Handbook* (Milne et al., 1995), as well as physical and chemical analyses from either the Landcare Research Environmental Chemistry Laboratory or the Department of Scientific and Industrial Research Soil Bureau. This data set was collated as the first stocktake of available soil data for national greenhouse gas reporting and, as such, underwent substantial quality-assurance and quality-control checks (Baisden et al., unpublished(b); Scott et al., 2002; Tate et al., 2005).

⁸ National Soils Database: <https://viewer-nsdr.landcareresearch.co.nz/search>.

Pre-1990 natural forest soils: This data set was gathered between 2002 and 2007 as part of the pre-1990 Natural Forest Survey, with soil subsampled on an 8-kilometre grid across the country (Garrett, unpublished; see section A3.2.5, 'National forest inventory' for more details of the 8-kilometre national grid system). The natural forest soils were important in the development of the Soil CMS model because they provided spatial balancing in areas of New Zealand not adequately covered by the historic soils data set.

Cropland data set: The third source of data originated as a set of intensively spatially sampled high producing grassland, annual cropland and perennial cropland records collected for other purposes, referred to as the cropland data set (Lawrence-Smith et al., 2010).

Wetlands: The fourth source of data comprises wetland soil data from a recent research effort to combine field data with analysis of the spatial distribution of current wetlands in New Zealand (Ausseil et al., 2015). This resulted in the addition of 21 wetland mineral soil samples to the Soil CMS data set (McNeill et al., 2014).


Post-1989 natural and planted forest data: This data set was added to the analysis in 2014. It contains data collected specifically for United Nations Framework Convention on Climate Change (Convention) reporting from 90 post-1989 forest sites across New Zealand (Basher et al., unpublished; Interpine Forestry Limited, unpublished).

Together, the five combined data sets cover most of New Zealand (see figure A3.2.5), including Stewart Island. Coverage does not extend to the Chatham Islands and other offshore islands. In addition to soil data, each record contains the site-specific climate, slope and rainfall attributes that are used in the analysis.

Due to a reliance on available data, coverage is dense in areas of agricultural activity, and the density of points varies widely between different regions (see figure A3.2.5). In addition, types of land use vary geographically: some are widespread (e.g., high producing grassland), whereas others are spatially constrained (e.g., cropland), so that the number of soil samples needed varies by land use (McNeill et al., 2014).

The number of records associated with the different land use categories and soil orders varies widely, with the largest land use category *Grassland* having 1,216 samples and the smallest (*Other land*) only three samples. While efforts to collect or obtain additional data in under-sampled land use categories have been made since LUCAS was established, helping to reduce uncertainties, the effect on uncertainty due to the considerable variability of sampling points among the different land use types remains.

Figure A3.2.5 Soil samples in the Soil CMS model calibration data set



Settlements and the open water component of *Wetlands* were not used in the model due to lack of soil carbon data. Both land uses are assigned the reference level carbon stock, which is the same as low producing grassland, because no data are available for these land uses. The basis for using the reference level for *Settlements* is supported by the land use definition used for the category because it includes not only impervious surfaces but also green spaces (urban park land, golf courses and other recreational areas). These areas are likely to have elevated carbon stock levels compared with low producing grassland due to the treatments they receive.

Ancillary data

In addition to the soil data, the following ancillary data are used in the Soil CMS Model.

S-map: S-map is a contemporary digital soil spatial information system for New Zealand (Lilburne et al., 2012), which provides the best-available knowledge of the classification of the soil order consistent with the *New Zealand Soil Classification* (Hewitt, 2010). S-map coverage is not available for all the land area, because its focus is on regions of intensive agricultural use.

Fundamental Soils Layer: Where data on soil order were unavailable in S-map, data from the Fundamental Soils Layer⁹ were used instead. The Fundamental Soils Layer provides GIS information on the expert-assessed classification of soil order and other soil or landscape attributes over New Zealand. It is generated from the NZLRI and National Soils Database.

Topographic information: Topographic slope information was estimated from a digital elevation model generated from Land Information New Zealand 1:50,000 scale topographic data layers including 20-metre contours, spot heights, lake shorelines and coastline.

Land use effects: Characterising soil carbon stocks

The 2014 version of the Soil CMS model used in this report builds on previous model versions (McNeill and Barringer, unpublished). The 'land use effect' (LUE) denotes the influence of land use on SOC stocks and corresponds to the model coefficients calculated for each land use. The LUE for a transition from low producing grassland to one of the other land uses can be obtained by using the coefficients of the soil carbon model (see table A3.2.5). Steady state SOC stocks for each land use (see table A3.2.6) are derived from the LUE coefficient in relation to the intercept (the reference of low producing grassland on high activity soils in a moist temperate climate, see table A3.2.5). These values are used in equation A3.2.3 (as SOC₀ and SOC_(0-T)) to calculate soil carbon changes due to land-use change.

Table A3.2.5 Land use effect coefficients with standard errors, t-values, and corresponding p-value significance estimates, extracted from full model results

Land use	Value	Standard error	t-value	p-value
Intercept: Low producing grassland	105.98	3.96	26.79	0.000
High producing grassland	-0.64	3.13	-0.21	0.8370
Grassland with woody biomass	-7.75	3.68	-2.11	0.0350
Perennial cropland	-17.54	6.37	-2.76	0.0059
Annual cropland	-16.21	4.45	-3.64	0.0003
Vegetated wetland	30.08	8.53	3.52	0.0004
Pre-1990 planted forest	-13.54	5.78	-2.34	0.0193
Post-1989 planted forest	-14.06	4.86	-2.90	0.0038
Pre-1990 natural forest	-13.73	3.70	-3.71	0.0002
Other land	-47.61	21.05	-2.26	0.0238

Source: McNeill and Barringer (unpublished)

Note: The model intercept (estimate for low producing grassland) is used for *Settlements* and *Wetlands – open water* land use categories due to lack of data.

Table A3.2.6 Steady state soil organic carbon stocks, with 95 per cent confidence intervals, calculated from Soil CMS model

Land use	Steady state carbon SOC stock (t C ha ⁻¹)	95% confidence intervals (CI)	
		2.5% CI SOC stock (t C ha ⁻¹)	97.5% CI SOC stock (t C ha ⁻¹)
Pre-1990 natural forest	92.25	84.99	99.51
Pre-1990 planted forest	92.44	81.12	103.77
Post-1989 planted forest	91.92	82.40	101.44
Post-1989 natural forest	91.92	82.40	101.44
Grassland with woody biomass	98.23	91.02	105.43
High producing grassland	105.34	99.21	111.47

⁹ Fundamental Soils Layer: <https://soils.landcareresearch.co.nz/tools/fsl/maps-fsl/>

Land use	Steady state carbon SOC stock (t C ha ⁻¹)	95% confidence intervals (CI)	
		2.5% CI SOC stock (t C ha ⁻¹)	97.5% CI SOC stock (t C ha ⁻¹)
Low producing grassland	105.98	98.23	113.73
Perennial cropland	88.44	75.96	100.92
Annual cropland	89.77	81.04	98.49
Wetlands – open water	105.98	98.23	113.73
Wetlands – vegetated	136.06	119.33	152.78
Settlements	105.98	98.23	113.73
Other land	58.37	17.12	99.62

Source: Calculated from McNeill and Barringer (unpublished)

An Akaike information criterion (AIC) model selection procedure was used for the Soil CMS model. AIC is used to select the model that is the best trade-off between the complexity of the model and the goodness of fit. The use of the AIC value as a model selection and comparison mechanism is widely supported in the literature in soil modelling (Burnham and Anderson, 2002; Elsgaard et al., 2012; Ogle et al., 2007).

The selected model residual standard error is 41.3 tonnes per hectare. The spatial autocorrelation scale distance is 18.1 kilometres, with a nugget of 0.47 (McNeill and Barringer, unpublished). A correction for spatial correlation is necessary to reduce the potential spatial bias in SOC stock values that may occur from multiple samples that are located close to one another. These values are consistent with earlier analyses (McNeill, unpublished(a), (b)).

The uncertainty of the LUE (the change in soil carbon, assuming the transition is stable) between two land use categories in isolation is conceptually straightforward: two estimates of LUE are more likely to be significantly separated if their point estimates are farther apart after taking account of the covariance between the two land use effects. The standard error $\sigma_{i,j}$ of the LUE change for a transition between two land use categories with effects L_i and L_j is then estimated from:

$$\sigma_{i,j} = \sqrt{\text{Var}(L_i) + \text{Var}(L_j) - 2 \cdot \text{Cov}(L_i, L_j)} \quad (\text{A3.2.3})$$

Where: $\text{Var}(L_i)$ = the variance of land use effect i

$\text{Cov}(L_i, L_j)$ = the covariance between land use effects L_i and L_j (McNeill and Barringer, unpublished; McNeill et al., 2014).


Although equation A3.2.3 provides a mathematically straightforward way to estimate the significance of a single transition from one land use category to another (a comparison-wise significance), it is often desirable to be able to determine whether a number of land use categories are likely to be significantly different or essentially the same as an ensemble. As more comparisons are made between many different land use types, it becomes more likely that at least one of the LUE changes will be different as a result of random chance alone, resulting in an increase in the Type 1 error. Thus, the significance of all possible land use transitions must be calculated as a family of simultaneous comparisons (multiple comparison significance), rather than one at a time (McNeill et al., 2014).

To control the Type 1 error rate in multiple comparison significance testing for the soil carbon change model, all possible combinations of the land use categories were tested for equality (a two-sided test) simultaneously. For the Soil CMS model (McNeill et al., 2014), a closed-

testing procedure described by Marcus et al. (1976) was used; this procedure is a general method for performing a number of hypothesis tests simultaneously implemented in the multi-comparison package in R (Bretz et al., 2010).

The closed-testing procedure described by Marcus et al. (1976) yielded point estimates and confidence intervals of a test statistic for each distinct combination of land use transitions, and the critical test is whether the confidence intervals include zero. All land use transition pairs were significant, except those involving *Other land* (see figure A3.2.6).

Figure A3.2.6 Result of applying the Marcus multi-comparison test to the adopted model



Source: McNeill et al. (2014)

Note: The marker is the estimated value for the specified transition to indicate significance, and the error bars represent the 95 per cent confidence interval of the test statistic. Land use transitions with point estimates and confidence intervals marked with a grey square are considered highly significant differences within the set of all possible land use transitions.

As the model results show (see figure A3.2.6), all transitions are significant in the multi-comparison sense, except those involving *Other land*. Land use transitions involving *Other land* contribute relatively little to the carbon change estimates, because they make up around 0.6 per cent of all land-use change detected between 1990 and 2021.

It is important to note that this interpretation of significance does not alter the method of calculation of the soil carbon change as a result of land use transition. In particular, it would not be correct to substitute a value of zero for the effect of a land use transition where the transition itself is not significant in the multi-comparison sense, because, if such a substitution were to be carried out, the calculation of the soil carbon would no longer be unbiased. Avoiding the bias in this manner also reduces the residual uncertainty of the soil carbon estimates. For this reason, the effect of all land use transitions ought to be included in calculations of soil carbon change (McNeill and Barringer, unpublished; McNeill et al., 2014).

Uncertainties in mineral soils

For the most part, uncertainties associated with the model coefficients (see table A3.2.6) are substantially reduced from the Tier 1 default value of 95 per cent. Land uses with higher uncertainties are those with few data points, such as *Other land*, or are dominant land uses in the country and, thus, occur across a range of environmental conditions, such as low producing grassland.

Uncertainties also arise from lack of soil carbon data for some soil, climate and land use combinations (Scott et al., 2002), and from variations in site selection, sample collection and laboratory analysis with data from different sources and time periods (Baisden et al., unpublished(b)). Other uncertainties in the Soil CMS model include: the assumption that soil carbon reaches steady state in all land uses and that there is a 20-year linear transition period to reach steady state; lack of soil carbon data and soil carbon change estimates below 0.3 metres; potential carbon losses from mass-movement erosion; and a possible interaction between land use and the soil–climate classification (Tate et al., 2004, 2005).

The inclusion of additional samples collected across a wider distribution has led to a reduction in the uncertainties for the land use effects, meaning all land use transitions, except for those involving *Other land*, are now significant in the multi-comparison sense (McNeill and Barringer, unpublished).

Source-specific quality control, quality assurance and verification

Quality-control and quality-assurance procedures have been adopted for all data collection and data analyses, to be consistent with 2006 IPCC Guidelines (IPCC, 2006a) and New Zealand's inventory quality-control and quality-assurance plan:

- details of the quality-management system for data collection, laboratory analyses and database management of the National Soils Database are given in Wilde (2003)
- recent data collection, analyses and management methods are subject to the soils quality-control and quality-assurance plan
- the consolidated soils data set used within the Soil CMS model has been subject to further quality-assurance procedures (Fraser et al., unpublished).

The Soil CMS model has been subject to various forms of testing, validation and recalibration. Testing of the Soil CMS model was completed to evaluate its ability to predict SOC stocks at regional and local scales. The results from the Soil CMS have been compared against independent, stratified soil sampling for South Island low producing grassland (Scott et al., 2002) and for an area of the South Island containing a range of land-cover and soil-climate categories (Tate et al., 2003a, 2003b). A regional-scale validation exercise has also been performed using the largest climate–soil–land use combination cell, moist temperate and volcanic × high producing grassland, within dependent random sampling of 12 profiles taken on a fixed grid over a large area (2,000 square kilometres). Mean values derived from the random sampling were well within the 95 per cent confidence limits of the database values (Tate et al., 2005; Wilde et al., 2004). A second study validated the Soil CMS model for a different cell, dry temperate – high-activity clay – low producing grassland, finding no significant differences among field data, calibration data and model estimates (Hedley et al., 2012). Overall, tests have indicated that the Soil CMS model estimates SOC stocks reasonably well at a range of scales (Tate et al., 2005).

The system has also been validated for its ability to predict soil carbon changes between land uses at steady state for New Zealand's main land-use change, grassland converted to planted forest. This was done by comparing the Soil CMS results with estimates based on paired sites (Baisden et al., unpublished(a); Tate et al., 2003a). This validation approach compares two nearby sites that have reasonably uniform morphological properties and were previously under a single land use, for which one site has changed to a different land use and sufficient time has elapsed for it to reach steady state values for soil carbon (Baisden et al., unpublished(a), unpublished(b)). This removes the influence that differing soil types, differing climatic conditions and previous land use regimes may have on soil carbon. Therefore, any resulting changes in soil carbon can be attributed to the most recent change in land use. In one study, results indicated that, once a weighting for forest species type was applied to the paired-site data set (to remove potential bias because *Pinus radiata* was under-represented in the analysis), the predictions of mean soil carbon from the Soil CMS model and paired sites were in agreement within 95 per cent confidence intervals (Baisden et al., unpublished(a), unpublished(b)). In a more recent study comparing low producing grassland and pre-1990 planted forests (Hewitt et al., 2012), the measured decrease in SOC under pre-1990 planted forest (-17.4 tonnes ha^{-1}) matched that determined by the Soil CMS model (McNeill et al., unpublished). This supported the Soil CMS model estimate (both in magnitude and direction) that forests planted pre-1990 have significantly lower SOC stocks than the low producing grassland and that the sampling depth of 0.3 metres was adequate for the estimation of SOC stock change.

The carbon stock estimates produced by the Soil CMS model reflect the type of soils in New Zealand (over 50 per cent being high activity clay soils) and the history of land use (fairly recent human settlement and forest clearance when compared with many other countries). As a comparison, when New Zealand reported using the Tier 1 default methodology (as in the 2011 submission), low producing grassland had the second highest SOC stock of all land uses (the highest being high producing grassland). The SOC stock for low producing grassland was also higher than for pre-1990 natural forests in that analysis.

Organic soils

Organic soils occupy a small proportion of New Zealand's total land area (1.0 per cent), and the area of organic soils subject to land-use change is around 0.7 per cent of New Zealand's total land area. New Zealand uses a Tier 1 method to estimate SOC stock change in organic soils.

The definition of organic soils is derived from the *New Zealand Soil Classification* (Hewitt, 2010), which defines organic soils as those soils with at least 18 per cent organic carbon in horizons at least 30 centimetres thick and within 60 centimetres of the soil surface. New Zealand-specific climate and soil data are used to estimate the areas of organic soil found in each climate zone. Climate data are based on the temperature data layer of the Land Environments New Zealand classification (Leathwick et al., 2002). Soil-type data are based on the Fundamental Soils Layer associated with the NZLRI (Newsome et al., 2008) and converted to the IPCC classification (Daly and Wilde, unpublished). These data layers have been analysed in a GIS system to determine the areas of organic soils in warm and cold climatic zones. These areas are compared with the land use to determine the area of organic soils in each.

The Land Use, Land-Use Change and Forestry (LULUCF) organic soils definition is the same as that used for reporting under the Agriculture sector (Dresser et al., 2011).

New Zealand has used IPCC default emission factors for organic soils under the *Forest land, Grassland, Cropland, Wetlands* and *Settlements* categories (IPCC, 2006a) to estimate organic soil emissions (see table A3.2.7). IPCC guidance for organic soils under forest is limited to

estimates associated with the drainage of organic soils in managed forests. In New Zealand, the drainage of pre-1990 natural forests does not occur, because the land is assumed to be in its natural state, and therefore no emissions are estimated from organic soils under natural forest. It is assumed that all planted forests on organic soils are drained before forest establishment. The temperate default emission factor for forest land is applied to the area of organic soils under planted forests to estimate emissions. The warm temperate and cold temperate default emission factors for the *Grassland*, *Cropland* and *Settlements* categories are applied in proportion to the area of land in New Zealand where the mean annual temperature is above or below 10°C respectively. New Zealand applies IPCC default emission factors for organic soils in the *Wetlands* category for areas under peat extraction. There are no default emission factors for organic soils under *Other land*; therefore, emissions from organic soils under this land use category are not estimated.

Table A3.2.7 New Zealand emission factors for organic soils

Land use	Climatic temperature regime	IPCC Tier 1 default emission factor applied and ranges (t C ha ⁻¹ yr ⁻¹)	Reference
Pre-1990 natural forest	Temperate	NA	IPCC guidance applies only to drained forest organic soils, which do not occur in natural forests in New Zealand (IPCC, 2006a, section 4.2.3.2).
Pre-1990 and post-1989 planted and natural forest	Temperate	0.68 (range 0.41–1.91)	IPCC (2006a, section 4.2.3.2, table 4.6)
Cropland	Cold temperate	5.0 ± 90%	IPCC (2006a, section 5.2.3.2, table 5.6)
	Warm temperate	10.0 ± 90%	
Grassland	Cold temperate	0.25 ± 90%	IPCC (2006a, section 6.2.3.2, table 6.3)
	Warm temperate	2.5 ± 90%	
Wetlands	NA	0.2 ± 90%	IPCC guidance applies to managed peatlands and flooded lands to which separate methodologies apply for soils. See IPCC, 2006a, chapter 7.
Settlements	Cold temperate	5.0 ± 90%	Cropland emission factors used (IPCC, 2006a, section 8.2.3.2)
	Warm temperate	10.0 ± 90%	
Other land	NA	NE	No IPCC guidance is available (IPCC, 2006a, chapter 9.3.3)

Note: NA = not applicable; NE = not estimated.

Uncertainties in organic soils

New Zealand uses the IPCC Tier 1 default value for uncertainty of organic soils under the categories *Forest land*, *Grassland*, *Cropland*, *Wetlands* and *Settlements*, as given in the 2006 IPCC Guidelines (2006a, tables 4.6, 5.6, 6.3 and 7.4). These values vary from 40 per cent for managed forests to 90 per cent for the other land uses.

Further detail on uncertainty for each land use is discussed in the appropriate category sections. The same method is used for all years of reporting to ensure time-series consistency.

A3.2.5 Forest land methodologies

Calculation of harvest area

Total destocking area (all harvesting and deforestation) for each year is first calculated for all planted forests. This total destocking area is then partitioned into deforestation and harvesting areas for pre-1990 and post-1989 planted forests using the following steps.

Total destocking area

1. Total destocking area between 1990 and 2012 is based on the harvested area reported in the NEFD (Ministry for Primary Industries, 2021) and adjusted to calendar years, plus the mapped deforestation area of post-1989 forest. The mapped deforestation of post-1989 planted forest is added because the NEFD is thought to underestimate destocking of forests belonging to small forest growers (Ministry for Primary Industries, 2021), which comprise a higher proportion of the post-1989 planted forest estate.
2. Total destocking area between 2013 and 2021 is calculated by combining planted forest yield tables, the destocking age profile (under 'Calculation of harvest area by age and forest age profile' below) and estimated roundwood volume removed from planted forests (Ministry for Primary Industries, 2022). This approach estimates the total destocking area required to achieve the Ministry for Primary Industries roundwood volume estimate, based on the average volume per hectare removed on harvest (calculated from the harvest age profile combined with LUCAS yield tables). This approach provides greater consistency with roundwood volume estimates and carbon inputs in the *Harvested wood product* category from 2013 to 2021 (figure A3.2.10).
3. The change in approach from 2013 onwards is due to concerns about the completeness of the NEFD survey, which shows an increasing mismatch in total harvest volume estimates for recent years compared with Ministry for Primary Industries roundwood removal statistics (Ministry for Primary Industries, 2022).

Deforestation area

1. Deforestation area from 1990 to 2019, for pre-1990 and post-1989 planted forest, is estimated from mapping data and supplementary statistics.
2. Deforestation area from 2020 onwards, for pre-1990 and post-1989 planted forest, is extrapolated from mapped years (see section A.3.2.2).

Harvest area

1. The harvest area of post-1989 forest from 2005 to 2007 is based on personal communication with industry experts.
2. The harvest area of post-1989 forest from 2008 to 2019 is based on mapped harvest area data.
3. From 2020 onwards, a harvest fraction approach is used to estimate the total destocking area in post-1989 and pre-1990 planted forest combined. This approach applies the overall harvest age profile (table A3.2.8) to the forest age profile in both forest types, to partition the area available to be harvested in each type. This provides an estimate of the destocking area to occur at each age in both forest types, as a proportion of the total destocking area.
4. Post-1989 harvest area from 2020 to 2021 is calculated as post-1989 total destocking area minus post-1989 deforestation area. A gross stocked to net stocked area adjustment is then applied. This reduces the estimated post-1989 harvest area but does not impact the total destocking estimate.
5. The harvest area of pre-1990 planted forest is then calculated for the whole time series (1990 to 2021) as total destocking area minus deforestation area (for both pre-1990 and post-1989 planted forest) and post-1989 harvest area.

Calculation of harvest area by age and forest age profile

Harvest and deforestation area by age

The harvest and deforestation area for pre-1990 and post-1989 planted forest is apportioned to an estimate of area by age (harvest age profile). This is because harvesting at a single age (28) years is not considered to reflect the actual harvesting that occurs and can lead to the harvest area exceeding the forest age available for harvest in some years. Estimating harvest area by age maintains the integrity of the forest age profile, limiting over-mature stands from growing on unharvested. The harvest or deforestation area by age is then combined with a yield table look-up value (see section A3.2.5, 'Planted forest yield tables' and tables A3.2.17–A3.2.19) to determine carbon losses.

A total destocking age profile is first calculated, which represents the percentage of total destocking (harvest and deforestation area) at each age class across all planted forest. The destocking age profile is derived from the loss of forest area in each age class with each annual update to the NEFD forest age profile (Ministry for Primary Industries, 2021). With each update, the losses in forest area for all age classes are combined to create an average destocking age profile, as a percentage of total destocking area. The average destocking age profile is then fitted to the average harvest age for each year, to capture the impact of the change in harvest age through time. Table A3.2.8 demonstrates the destocking age profile for all planted forest expressed as a percentage for the ages 15–45 years. Note that there is a small proportion of destocking that occurs before the age of 15 years (not shown in table A3.2.8). This relates to post-1989 destocking that predominately occurs prior to 2008.

An average harvest age of 28 years is assumed for 1990 to 1995. For subsequent years, the average harvest age is sourced from annual NEFD publications (Ministry for Primary Industries, 2021). The average harvest age is converted to calendar years and a three-year moving average is applied to smooth out any year-to-year fluctuations. The average harvesting age is considered to represent the entire planted forest estate (i.e., both pre-1990 and post-1989 forests). The average harvesting age is considered to represent the entire planted forest estate (i.e., both pre-1990 and post-1989 forests).

The destocking age profile is then combined with the annual harvest and deforestation area in pre-1990 and post-1989 planted forest. This gives an estimate of harvest and deforestation area by age for each forest type. The final harvest area by age profile in 2021 is demonstrated in figure A3.2.7 for pre-1990 planted forest and in figure A3.2.8 for post-1989 planted forest.

Both pre-1990 and post-1989 planted forests share the same underlying destocking and harvest age profiles. Therefore, as the rate of harvest increases in the post-1989 estate (with increasing harvest of younger trees due to the forest age profile of post-1989 forests), the proportion of harvesting of pre-1990 forest in these ages is reduced and the average harvest age of pre-1990 planted forest increases. This ensures that the average harvest age and harvest age profile are representative of the entire planted forest estate.

Forest age profile

Post-1989 planted forest

The forest age profile in post-1989 planted forest is driven by the area of new planting from 1990 onwards (see section A.3.2.2), adjusted for any harvesting or deforestation area.

Pre-1990 planted forest

The forest age profile in pre-1990 planted forest is driven by annual harvest area for all stands planted after 1990. A one-year lag between harvesting and replanting is assumed. This means an estimated harvest area of 18,789 hectares in 1990 will result in replanting of the same area in 1991.

Annual planting area before 1990 is established to meet the required harvest and deforestation area by age estimates from 1990 onwards. This means an estimated 3,000 hectares of forest harvested at age 30 in 2010 would require that same area to have been planted in 1980. The planting area by year required is then apportioned into new planting, based on the area converted to pre-1990 planted forest from 1962 to 1989 (see section A.3.2.2), or assigned harvest and replanting events.

The forest age profile for the remaining forest area that is not subject to harvest or deforestation after 1990 is estimated from the NEFD forest age profile. The forest age profile in the most recent reporting year for all forest planted before 1990 is estimated by multiplying the area of this forest by the proportion of forest in each age from the NEFD. This results in the area in each age group being slightly higher than the NEFD estimate, this difference can be seen in forest aged over 30 years in figure A3.2.11. The forest area by age in the most recent report year is then assigned a corresponding plant date.

Table A3.2.8 Proportion of total destocking area by age across all planted forest, 1990–2021

Year	Destocking age by year (%)																														
	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45
2021	0.4	1.0	0.9	0.9	0.6	1.4	0.8	1.7	0.8	2.6	4.1	6.7	10.6	12.2	12.2	7.1	8.9	6.9	5.4	3.9	2.8	2.1	1.6	1.3	0.8	0.6	0.5	0.4	0.3	0.2	0.3
2020	0.4	1.0	0.8	0.9	0.6	1.4	0.8	1.7	0.8	2.5	4.1	6.7	10.7	12.3	8.4	10.6	9.0	7.0	5.5	3.9	2.9	2.1	1.6	1.3	0.7	0.7	0.5	0.4	0.3	0.2	0.3
2019	0.1	0.2	0.2	0.2	0.2	1.4	0.8	1.7	0.8	2.6	4.0	6.6	10.7	9.5	12.6	10.9	9.2	7.2	5.6	4.1	2.9	2.1	1.7	1.3	0.8	0.7	0.5	0.4	0.3	0.2	0.3
2018	0.1	0.4	0.3	0.3	0.5	1.1	1.2	0.9	1.8	1.8	5.0	10.0	9.4	14.2	11.8	9.9	7.9	6.3	4.6	3.1	2.3	2.0	1.4	0.9	0.7	0.5	0.5	0.3	0.3	0.2	0.2
2017	0.1	0.4	0.3	0.3	0.4	1.2	1.3	0.9	1.8	1.8	5.1	7.2	12.2	15.2	11.5	9.7	7.7	6.4	4.3	3.0	2.2	1.9	1.5	0.9	0.6	0.6	0.5	0.3	0.2	0.2	0.2
2016	0.2	0.4	0.3	0.4	0.5	1.3	1.1	1.3	1.6	2.4	4.1	9.3	11.5	15.3	11.2	9.5	7.9	5.7	4.1	2.9	2.1	1.9	1.5	0.8	0.7	0.6	0.4	0.3	0.2	0.2	0.2
2015	0.2	0.4	0.4	0.4	0.7	1.2	1.2	1.2	1.5	1.6	4.7	8.5	11.2	14.3	11.2	10.5	7.8	6.1	4.5	2.8	2.4	2.2	1.3	1.0	0.8	0.6	0.4	0.3	0.2	0.2	0.2
2014	0.5	0.5	0.6	0.6	0.6	1.3	0.8	1.5	1.0	2.7	7.4	9.4	14.2	10.9	12.2	8.7	7.0	5.3	3.6	2.5	2.5	1.5	1.2	0.9	0.6	0.5	0.3	0.2	0.2	0.2	0.3
2013	0.5	0.7	0.6	0.6	0.6	1.4	1.0	1.1	1.3	2.8	6.2	10.5	11.4	13.3	11.0	9.0	7.3	5.5	3.2	3.4	1.8	1.9	1.5	0.8	0.6	0.6	0.3	0.3	0.3	0.2	0.3
2012	0.6	0.7	0.6	0.8	0.8	1.3	0.6	1.4	1.6	2.9	7.1	10.7	12.6	12.6	10.6	8.6	6.9	4.9	4.0	2.5	2.0	2.0	1.1	0.8	0.6	0.5	0.4	0.3	0.3	0.2	0.2
2011	1.0	0.6	0.5	0.7	0.7	0.9	0.9	1.4	1.3	2.4	5.9	9.5	11.7	13.0	11.2	9.2	7.5	5.7	4.1	2.9	2.3	1.9	1.3	0.9	0.6	0.6	0.4	0.3	0.2	0.2	0.2
2010	0.5	0.7	0.6	0.8	0.7	1.2	1.0	1.4	1.3	2.5	6.4	9.9	12.0	13.0	11.0	9.0	7.3	5.5	3.9	2.8	2.2	1.8	1.2	0.7	0.7	0.6	0.4	0.3	0.2	0.2	0.2
2009	0.8	0.8	0.7	0.8	1.0	1.3	0.8	1.4	1.4	3.1	7.2	10.4	12.0	12.7	10.6	8.6	6.9	5.2	3.6	2.6	2.1	1.7	1.0	0.8	0.6	0.6	0.3	0.3	0.2	0.2	0.3
2008	0.8	0.8	0.6	1.0	0.2	1.2	0.9	1.5	1.6	3.6	7.8	11.0	12.4	12.3	10.4	8.4	6.7	5.0	3.5	2.5	2.1	1.6	1.1	0.7	0.6	0.5	0.3	0.3	0.2	0.2	0.3
2007	2.2	0.9	0.8	0.5	0.4	1.1	0.8	1.3	1.5	3.4	7.2	10.1	11.2	12.5	10.9	8.7	6.9	5.1	3.6	2.6	1.4	1.8	1.7	0.8	0.6	0.5	0.3	0.3	0.2	0.2	0.3
2006	0.7	0.8	0.4	0.4	0.4	1.2	0.9	1.5	1.6	3.7	7.8	10.9	12.2	12.3	10.7	8.5	6.8	5.1	3.6	1.9	2.2	2.2	1.1	0.8	0.6	0.5	0.3	0.3	0.2	0.2	0.2
2005	0.9	0.3	0.3	0.3	0.3	1.1	1.0	1.4	1.9	4.4	8.4	11.2	12.1	12.2	10.5	8.4	6.6	4.9	2.7	3.3	2.1	1.6	1.0	0.7	0.6	0.5	0.3	0.3	0.2	0.2	0.3
2004	0.6	0.8	0.1	0.1	0.2	0.9	1.2	1.4	2.3	5.7	9.4	11.7	11.8	13.7	9.6	7.7	5.9	4.3	3.0	2.4	1.9	1.3	0.9	0.7	0.6	0.4	0.3	0.3	0.2	0.2	0.3
2003	0.1	0.8	1.1	0.1	0.1	0.8	1.4	1.3	2.6	6.4	10.0	12.1	11.7	13.3	9.2	7.4	5.7	4.0	2.8	2.3	1.9	1.2	0.8	0.6	0.6	0.4	0.3	0.2	0.2	0.2	0.3
2002	0.1	0.2	0.8	1.0	0.0	0.7	1.6	1.3	3.0	7.5	11.1	12.9	12.0	12.5	8.5	6.9	5.2	3.6	2.6	2.1	1.8	1.1	0.8	0.6	0.5	0.3	0.2	0.2	0.2	0.3	0.3
2001	0.0	0.3	0.2	1.0	0.9	0.6	1.7	1.3	3.2	8.3	11.8	13.3	11.9	12.4	8.1	6.5	4.8	3.3	2.3	2.0	1.7	0.9	0.7	0.6	0.5	0.3	0.2	0.2	0.2	0.3	0.3
2000	0.0	0.0	0.3	0.2	0.8	0.6	1.8	1.3	3.4	8.9	12.3	13.6	12.0	12.5	8.0	6.4	4.7	3.2	2.3	2.0	1.6	0.9	0.7	0.5	0.5	0.3	0.2	0.2	0.2	0.3	0.3
1999	0.0	0.0	0.0	0.2	0.2	0.9	1.5	1.4	2.8	7.1	11.0	13.1	12.7	11.4	9.1	7.3	5.6	4.0	2.8	2.3	1.9	1.2	0.8	0.6	0.6	0.4	0.3	0.3	0.2	0.3	0.3

Year	Destocking age by year (%)																														
	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45
1998	0.0	0.0	0.0	0.0	0.4	1.1	1.2	1.5	2.3	5.5	9.6	12.4	12.8	12.2	9.7	7.8	6.1	4.5	3.1	2.4	2.0	1.4	0.9	0.7	0.6	0.4	0.3	0.3	0.2	0.2	0.3
1997	0.0	0.0	0.0	0.0	0.0	1.1	1.2	1.5	2.1	5.1	9.3	12.3	13.0	12.2	9.9	8.1	6.3	4.6	3.3	2.4	2.0	1.5	1.0	0.7	0.6	0.5	0.3	0.3	0.2	0.2	0.3
1996	0.4	0.0	0.0	0.0	0.0	1.1	1.1	1.5	2.0	4.9	9.1	12.1	12.9	12.5	9.9	8.0	6.3	4.6	3.3	2.4	2.0	1.5	1.0	0.7	0.6	0.5	0.3	0.3	0.2	0.2	0.3
1995	0.0	0.5	0.0	0.0	0.0	1.2	1.1	1.5	2.0	4.7	9.0	12.1	13.0	12.2	10.0	8.1	6.4	4.7	3.3	2.5	2.0	1.5	1.0	0.7	0.6	0.5	0.3	0.3	0.2	0.2	0.3
1994	0.2	0.2	0.4	0.0	0.0	1.2	1.0	1.5	1.8	4.3	8.6	11.8	13.0	12.5	10.2	8.2	6.6	4.9	3.4	2.5	2.1	1.6	1.0	0.7	0.6	0.5	0.3	0.3	0.2	0.2	0.3
1993	0.7	0.4	0.3	0.8	0.0	1.2	1.0	1.5	1.8	4.2	8.4	11.6	12.7	12.6	10.0	8.1	6.4	4.8	3.3	2.5	2.0	1.6	1.0	0.7	0.6	0.5	0.3	0.3	0.2	0.2	0.3
1992	0.7	0.8	0.3	0.3	0.8	1.2	1.0	1.5	1.8	4.1	8.4	11.5	12.6	12.6	9.9	8.0	6.4	4.7	3.3	2.4	2.0	1.6	1.0	0.7	0.6	0.5	0.3	0.2	0.2	0.2	0.3
1991	0.0	0.8	0.8	0.6	0.5	1.2	1.0	1.5	1.8	4.1	8.4	11.5	12.7	12.8	9.9	8.0	6.4	4.7	3.3	2.4	2.0	1.6	1.0	0.7	0.6	0.5	0.3	0.2	0.2	0.2	0.3
1990	0.3	0.0	0.9	1.1	1.0	1.2	1.0	1.5	1.7	4.0	8.2	11.2	12.4	14.1	9.7	7.8	6.2	4.6	3.2	2.4	2.0	1.5	1.0	0.7	0.6	0.5	0.3	0.2	0.2	0.2	0.3

Figure A3.2.7 Harvest area by age for pre-1990 planted forest in 2021





Figure A3.2.8 Harvest area by age for post-1989 planted forest in 2021



National forest inventory

New Zealand has established a sampling framework for forest inventory purposes based on an 8-kilometre national grid system (8-kilometres north–south by 8–kilometres east–west). The grid has a randomly selected origin and provides an unbiased framework for establishing plots for field and/or Light Detection and Ranging (LiDAR) measurements. The network is further subdivided into a 4-kilometre grid for measurement of post-1989 forest. Forest monitoring plots are established and measured where a grid point falls in the land use to be sampled.

Pre-1990 natural forest

A national monitoring programme designed to enable unbiased estimates of carbon stock and change for New Zealand’s natural forests was developed between 1998 and 2001 (Coomes et al., 2002). Permanent circular sample plots of 0.13 hectares (i.e., 20 metre radius) were installed systematically on the 8-kilometre grid across New Zealand’s natural forests and these were first measured (t_1) over five years between 2002 and 2007.

The plots were sampled using vegetation monitoring methods designed specifically for the purpose of calculating carbon stocks (Payton et al., 2004). A 0.04 hectare plot (20 × 20 square metres) sits nested at the centre of each circular plot where all live stems with diameter at breast height (1.35 metres) greater than or equal to 2.5 centimetres are measured. Stems greater than 60 centimetres diameter at breast height are sampled on the 0.13 hectare circular plot.

Re-measurement of the plot network provides repeat data suitable for calculating carbon stock change in natural forest. The first re-measurement of the plot network was completed between 2009 and 2014 (t_2) following a revised methodology for re-measurement purposes (Ministry for the Environment, unpublished). For the third round of measurement, the programme is continuing at a reduced rate, with plots being measured on a 10-year cycle. Measurement of plots for this round began in 2014 and is scheduled for completion in 2024. Fieldwork generally takes place between October and March each year. Data collection on the natural forest plot network has recently transitioned (2020–21 field season) to an electronic data capture system. It had previously relied on a paper-based system. This will improve data quality and reduce the time between field collection and analysis.

At each plot, data are collected to calculate the volumes of trees, shrubs and dead organic matter present. These measurements are then used to estimate the carbon stocks for the biomass pools of:

- living biomass (comprising above-ground biomass and below-ground biomass)
- dead organic matter (comprising dead wood and litter).

Table A3.2.9 summarises the method used to calculate the carbon stock in each biomass pool from the information collected at each plot.

Table A3.2.9 Summary of methods used to calculate New Zealand’s natural forest biomass carbon stock from plot data

Pool		Method	Source
Living biomass	Above-ground biomass	Plot measurements; allometric equations	Paul et al., 2021
	Below-ground biomass	Estimated as the ratio of below-ground biomass to above-ground biomass	Paul et al., 2021; Easdale et al., 2019
Dead organic matter	Dead wood	Modelled from plot measurements; allometric equations	Garrett et al., 2019; Paul et al., 2021; Kimberley et al., 2019
	Litter	Plot samples; laboratory analysis of samples collected at plots	Paul et al., 2021; Garrett, unpublished

Living biomass

Living biomass is separated into two carbon pools.

1. Above-ground biomass. The carbon content of individual trees and shrubs is calculated using species-specific allometric relationships between diameter, height and wood density (for trees), a non-specific conversion factor with diameter and height (for tree ferns) or volume and biomass (for shrubs) (Beets et al., 2012b; Paul et al., 2021). Shrub volumes are converted to carbon stocks using species- and/or site-specific conversion factors determined from the destructive harvesting of reference samples. Carbon fractions of 0.51 for gymnosperms and 0.48 for broadleaf species (IPCC, 2006a).
2. Below-ground biomass. The below-ground biomass for each individual tree is calculated based on an estimate of the root:shoot ratio for that species (the ratio of the below-ground biomass to above-ground biomass). Applying the root:shoot ratios as

published in Easdale et al. (2019) has been included to address the expert review team recommendation L.4, 2019 (FCCC/ARR/2019/NZL, UNFCCC, 2020). Tree and shrub species in different taxonomic groups were assigned different root:shoot ratios, as outlined in (Paul et al., 2021) and are summarised in table A3.2.10.

Table A3.2.10 Summary of root:shoot ratios applied to the different taxonomic groups in pre-1990 natural forest

Taxonomic group	Root:shoot ratio
Angiosperm trees (> 5 cm diameter at breast height)	0.234
Monocots (palms and cabbage trees)	0.194
Gymnosperms and shrubs	0.235

Dead organic matter

Dead organic matter is separated into two carbon pools.

1. Dead wood. The carbon content of dead standing trees is determined in the same way as live trees but excludes branch and foliage biomass calculations. The carbon content of the fallen wood and stumps is derived from the volume of the piece of wood, its species (if it can be identified) and what stage of decay it is at. Dead wood comprises woody debris with a diameter greater than 10 centimetres. The dead wood pool is difficult to measure in the field (particularly for wood that is in an advanced state of decay) and is currently being underestimated by the monitoring programme (Kimberley et al., 2019). An adjustment factor, derived by an approach developed by Kimberley et al. (2019), was applied to correct for this (Paul et al., 2021). Deadwood is measured on all new plots and modelled for re-measured plots using initial measurements, inputs from mortality and known decay rates (Paul et al., 2021).
2. Litter. The carbon content of the fine debris is calculated by laboratory analysis of sampled material. The samples are bulked by sampling depth (0–10 centimetres, 10–20 centimetres, 20–30 centimetres) and the total fine earth mass is measured and then analysed for carbon content. Litter comprises fine woody debris (dead wood from 2.5 centimetres to 10.0 centimetres in diameter), the litter (all material less than 2.5 centimetres in diameter) and the fermented humic horizons. Samples were taken at around one-third of the natural forest plots.

Carbon stock change

Carbon stock change in the living biomass pool is calculated using the methods described in Paul et al. (2021). In this method, carbon stock change for each plot is calculated by summing the stock change for each individual live stem and subtracting the summed carbon at t_1 for individual stems that died in the period between t_1 and t_2 . To account for ingrowth (stems that have reached the 2.5 centimetre diameter at breast height threshold since the last plot measurement) and missing measurements, the diameter of trees measured at t_2 that were not measured at t_1 were predicted and used in the calculation of stock change, provided that the diameter at t_2 was above the threshold for field measurement (e.g., 2.5 centimetres for the embedded 0.04 hectare square plot, and 60 centimetres for the 0.13 hectare circular plot). The total summed carbon is calculated for each plot, and the mean change across all plots measured twice is used as the national average.

New Zealand has inventoried its pre-1990 natural forest at two points in time: 2002 to 2007 and 2009 to 2014 (the third round of measurements is under way and due for completion in 2024). The average measurement date of the first measurement period is 2004 and average

measurement date of the second measurement period is 2011. Pre-1990 natural forest was classified into tall and regenerating subcategories using the 2008 land cover mapped in Land Cover Database version 5.0. Carbon stock change was then calculated separately for both subcategories.

Between 2002 and 2007 and 2009 and 2014, the regenerating forest component of New Zealand's pre-1990 estate had a rate of carbon stock change of 0.43 ± 0.51 tonnes C ha⁻¹ yr⁻¹ (estimated from Paul et al., 2021). The tall forest component changed very little over the same period (-0.01 ± 0.19 tonnes C ha⁻¹ yr⁻¹). The data for both components are extrapolated back to 1990 and forward to the current inventory year to calculate stock changes for all years. The combined overall net change across all pre-1990 natural forest was indistinguishable from zero (0.03 ± 0.18 tonnes C ha⁻¹ yr⁻¹; estimated from Paul et al., 2021). Carbon stock change in regenerating forest was driven primarily by an increase in live above-ground biomass of 0.36 ± 0.26 tonnes C ha⁻¹ yr⁻¹ (Paul et al., 2021). Carbon stock change in tall forest was driven primarily by a decrease in live above-ground biomass of -0.01 ± 0.15 tonnes C ha⁻¹ yr⁻¹.

In an effort to reduce sampling uncertainty and fulfil the practical recommendations made by Holdaway et al. (2014) and the related expert review team recommendation L.1, 2019 (FCCC/ARR/2019/NZL, UNFCCC, 2020), several improvements have been implemented in the management of the natural forest plot measurement programme and analysis of data over time. First, the number and size of plots included in carbon stock and stock change analyses have increased through time. A total of 874 plots were included in Holdaway et al. (2017). This has increased to 1,030 plots for updated carbon stock calculations and 908 plots for updated carbon stock change calculations in Paul et al. (2021). Paul et al. (2021) included stems from a larger plot area (0.13 hectares) than in previous analyses, which only included stems from the nested 20 × 20 metre (0.04 hectare) plot (Holdaway et al., 2017).

Second, changes to the approach for estimating dead organic matter (i.e., adjusting for under-estimation of field measurements) represents an improvement in stock and stock change estimates. Third, the stem-level carbon stock change methods used by Paul et al. (2021) and described above account for ingrowth stems and missed stems. This reduces bias in the carbon stock change estimate and represents an improvement on previous methods (Holdaway et al., 2017) where a simple stock change approach was used. The effect of some of these improvements to methodologies has been outlined and quantified in table 8 of Paul et al. (2021).

Additional changes were made for the 2022 submission to align the plot-based carbon stock analysis with the mapped areas of each land cover sub-class. Pre-1990 tall and regenerating forest plots were previously classified using species composition (Wiser, 2016) to estimate the annual carbon stock change of each forest type. However, the tall and regenerating forest areas were estimated from mapped land cover (LCDBv3). This created a mismatch where carbon stock change estimates were not consistent with the forest area they were intended to represent. This was corrected for the 2022 submission by using the land cover approach (LCDBv5) to classify both the plots and the mapped areas of tall and regenerating forest. This change resulted in a difference in emissions in 2019, increasing the estimate by approximately 1,000 kt CO₂-e in the pre-1990 natural regenerating forest category.

Post-1989 natural forest

Estimates of carbon stock and stock change in post-1989 natural forest are calculated using measurements taken from the field inventory. The inventory samples post-1989 natural forest using 0.13 hectare permanent sample plots on the systematic 4-kilometre grid. Twenty plots in post-1989 natural forests were established and measured for the first time in 2012. A second

round of measurements, on 25 plots (of which 13 were also measured in 2012), was conducted in 2019. A yield table was generated from the plot measurements to provide estimates of carbon stock change (Paul et al. unpublished(b)). The plot network design is described in Beets et al. (2012a, 2014b), and detailed methods for plot measurement are given in the data collection manual (Ministry for the Environment, unpublished).

Living biomass and dead organic matter

At permanent sample plots within post-1989 natural forest, measurements are taken of standing and fallen, live and dead plants. Destructive biomass samples have also been taken outside of the plots and are used to create plot-specific allometric equations, which are then applied to these measurements to calculate above-ground live biomass.

The biomass of standing dead wood (woody debris with a diameter greater than 10 centimetres) and litter (woody debris with a diameter of less than 10 centimetres) is measured and calculated using the same methods as used in pre-1990 natural forest described above.

Biomass sampling on post-1989 natural forest plots includes the determination of plant age, which enables the back-casting of biomass through time. Back-cast estimates of biomass are used to calculate carbon stock change. The method used to do this was developed and validated using plots for which multiple measurements in time had been obtained and for which carbon stock change was able to be measured directly (Beets et al., 2014a). Full methods for the calculation of carbon stock and stock change in post-1989 natural forest are described in Beets et al. (2014b) and Paul et al. (unpublished(b)).

Carbon stock change in the living biomass pool is calculated using the methods described in Paul et al. (unpublished (b)). In addition, a post-1989 natural forest yield table is included and is used in conversions from *Grassland with woody biomass* (see table A3.2.16). The yield table starts at the same carbon stock as *Grassland with woody biomass* resulting in no emissions from biomass in the first year of conversion because this conversion represents ecological succession.

The carbon stock estimate for post-1989 natural forest was 38.55 ± 10.23 tonnes C ha⁻¹ (at the 95 per cent confidence interval) as of 31 December 2019 (Paul et al. unpublished(b)). The average rate of carbon sequestration in post-1989 natural forest between 2012 and 2019 was 2.48 tonnes C ha⁻¹ yr⁻¹ (calculated from Paul et al., unpublished(b)). This rate is slightly higher than previously reported rates of carbon sequestration in regenerating forest in New Zealand (Carswell et al., 2012; Trotter and MacKay, unpublished). This possibly reflects differences in the composition of species that were targeted in these studies (Paul et al. unpublished(b)).

Planted forest

The planted forest inventory consists of 749 circular 0.06-hectare plots established on the systematic 8-kilometre grid and nested 4-kilometre grid as described above (339 in pre-1990 planted forest and 410 in post-1989 planted forest). These plots are ground measured using procedures described in Herries et al. (unpublished). Stand records and ground measurements are recorded between June and October at each plot. Measurements include tree age; stocking (stems per hectare); stem diameters at breast height of live and dead trees; a sample of tree total heights for each tree species; pruned heights; and the timing of pruning and thinning activities. Ground plot centres were located using a 12-channel differential global positioning system (GPS) for accurate LiDAR co-location and relocation for future measurements (Beets et al., 2011a; 2012a).

Living biomass and dead organic matter

The crop tree plot data collected from the planted forest inventories are modelled using a forest carbon modelling system (the Forest Carbon Predictor, version 4.12; Beets and Garrett, 2018; Beets et al., 2018a, 2018b; Paul et al., unpublished(a); Paul and Wakelin, unpublished) developed for the two most common plantation tree species in New Zealand: *Pinus radiata* and *Pseudotsuga menziesii*. To enable predictions of carbon stocks and changes in New Zealand's planted forests, this system integrates:

- the 300 Index growth model (Kimberley and Dean, 2006) for *Pinus radiata*
- the 500 Index growth model for Douglas fir (Knowles, 2005)
- a wood density model (Beets et al., 2007)
- a stand tending model (Beets and Kimberley, unpublished)
- the C_Change carbon allocation model (Beets et al., 1999).

The individual components of the Forest Carbon Predictor are explained below and illustrated in figure A3.2.9.

The 300 Index and 500 Index growth models produce a productivity index for forest plots derived from stand parameters. These stand parameters include stand age, mean top height, basal area, stocking and stand silvicultural history. Plot latitude and altitude are also required to run the models. The growth models use these parameters to predict stem volume under bark over a full rotation (planting to harvest). A specific productivity index is produced for each plot, which is then used to estimate the total live and dead stem volume by annual increment. The growth models account for past and future silvicultural treatments using plot data, information on past silvicultural treatments and assumptions of future management events based on plot observations and standard regimes (Beets and Kimberley, unpublished).

The wood density model within the Forest Carbon Predictor uses site mean annual temperature, soil nitrogen fertility, ring age and stocking to determine the mean density of stem wood growth sheaths produced annually in *Pinus radiata*. Wood density is an important variable in the estimation of carbon. Of the parameters entered into the wood density model, temperature and stand age have the greatest influence on wood density, followed by site fertility and stocking. The combined result of these individual effects can be substantial, as shown in table A3.2.11 (Beets et al., 2007).

Table A3.2.11 Influence of individual site and management factors on predicted wood density for New Zealand planted forest

Factor affecting wood density	Range in predicted density	
	(kg m ⁻³)	(% difference)
Temperature: 8°C versus 16°C	359–439	22
Age: 10-year-old versus 30-year-old	380–446	17
C:N ratio: 12 versus 25	384–418	9
Stocking: 200 versus 500 stems ha ⁻¹	395–411	4

Note: C:N = carbon:nitrogen.

The stand tending model: New Zealand's plantation forests are intensively managed; therefore, pruning and thinning provide the majority of the inputs to the dead wood and litter pools. The Forest Carbon Predictor requires silvicultural history inputs to predict changes between biomass pools over time. The information required includes initial stocking, the timing of management events, stocking following each thinning operation and the pruned height and number of stems pruned for each pruning lift. Information on silvicultural events

before the plot measurement date is normally gathered from forest owners but sometimes these data are incomplete. A history module has been incorporated into the Forest Carbon Predictor that makes use of existing data to identify potential gaps in the stand history. Within the history module, assumptions are made to complete the stand history based on field observations, standard management regimes and known silviculture to date (Beets and Kimberley, unpublished). The history module enables reasonable estimates of stand history and, therefore, biomass transfers between pools resulting from past silvicultural events.

The C_Change carbon allocation model is designed to apportion carbon to needles, branches, stems, roots and reproductive parts via growth partitioning functions and is integrated into the Forest Carbon Predictor. Dead wood and litter pools are estimated by accounting for losses to the live pools from natural mortality, disease effects on needle retention, branch and crown mortality and silvicultural management activities, for example, pruning and thinning. Component-specific and temperature-dependent decay functions are used to estimate losses of carbon to the atmosphere (Beets et al., 1999). The Forest Carbon Predictor also takes into account biomass removals during production thinning.

The individual plot yield curves generated by the Forest Carbon Predictor are combined into estimates of above-ground live biomass, below-ground live biomass, dead wood and litter in an area-weighted and age-based carbon yield table for the productive area of each type of planted forest. Plots that are located outside the productive area within the mapped forest boundary are used to provide emission factors for unstocked areas in both post-1989 forest and pre-1990 planted forest (Paul et al., unpublished(f)).


Below-ground biomass is derived from the above-ground biomass estimates. For plantation crop trees, below-ground biomass is assumed to be 15 per cent to 20 per cent of total production, depending on stand age (Beets et al., 1999). The ratio for non-crop trees and shrubs is 25 per cent (Coomes et al., 2002).

The carbon content of the dead wood pool within a rotation is estimated using the Forest Carbon Predictor model as described above. Immediately following harvesting, 30 per cent of the above-ground biomass pool is transferred to the dead wood pool; the other 70 per cent is instantaneously emitted. All material in the dead wood and litter pools is decayed using an empirically derived, temperature-dependent decay profile as described in Garrett et al. (2010).

Yield tables: Mean yield tables are derived from individual plot tables, described below using plot area-weighted averages. Yield values based on the backcast values from the first measurements are used from year 0 to the year of the first measurement. A straight interpolation is used between the first and the second measurement and any subsequent measurements.

From the last measurement onwards the forecasts are based on the most recent measurement to predict the stand yield until age 60 for both post-1989 planted forest and pre-1990 planted forest plots. A further adjustment is made using an imputation method to account for forecasting and backcasting errors. This method applies a greater weighting at yield table ages close to the plot measurement age. The planted forest yield tables used in this submission are given in section A3.2.5, 'Planted forest yield tables'.

Figure A3.2.9 New Zealand's planted forest inventory modelling process



Note: CRF = common reporting format; LiDAR = Light Detection and Ranging.

For shrubs and non-crop tree species measured within the planted forest plot network, the carbon content is estimated using species-specific allometric equations. These equations estimate carbon content from diameter and height measurements, and wood density by species (Beets et al., 2012a).

For the 2022 submission, the period-specific yield tables used for pre-1990 planted forest were updated to improve accuracy. In earlier submissions (before 2022), three yield tables were applied to the pre-1990 planted forest estate, representing stands planted before 1990, between 1990 and 2009, and from 2010 onwards. However, the yield table for stands planted from 2010 onwards was found to have high uncertainty and tended to overestimate carbon stock per hectare relative to the field measurements. During the review of the 2021 submission, it was suggested that submissions revert back to two yield tables until further data are available for the yield table representing stands from 2010 onwards. Therefore, in this submission two yield tables have been applied: one for stands planted before 1990 and one for stands planted from 1990 onwards.

For the 2022 submission, the post-1989 planted forest yield table was revised to include plots measured in the 2020 forest inventory. The revised yield table is based on two full measurement inventories of permanent sample plots in post-1989 planted forest carried out in 2008/09 and 2011/12, and a partial re-measure annually between 2016 and 2020 (due to the transition to the continuous five-year inventory cycle).

Pre-1990 planted forest

Stock change in the productive area of pre-1990 planted forests is estimated using forest type-specific national yield tables. Plots that are located outside the productive area within the mapped forest boundary are used to provide emission factors for unstocked areas of pre-1990 planted forests (Paul et al., unpublished(a)).

A stratification approach has been developed to stratify the data, allowing the modelling of period-dependent yield tables, creating historic and current yield tables based on reporting periods for pre-1990 planted forests (Paul and Wakelin, unpublished). These yield tables better reflect the conditions and productivity during the past. Using the plot measurements described above under the pre-1990 planted forest inventory, a single yield table per plot was developed using:

- the earlier measurement for ages below the first measurement age
- the later measurement for ages above the later measurement age
- an interpolated estimate for the ages between the earlier and later measurements.

For plots that have been measured once, a ratio estimator derived from plots that have been measured twice is applied to the predicted stocks at the missing measurement date (assuming that the correction for possible bias was the same in both strata) (Paul et al., unpublished(a)).

Post-1989 planted forest

In the post-1989 planted forest inventory, circular 0.06 hectare permanent sample plots have been established within forests on a systematic 4-kilometre grid coincident with that used for the pre-1990 natural forest and pre-1990 planted forest inventories (Moore and Goulding, unpublished). Permanent sample plots were selected over temporary sample plots because change over time is more easily analysed when there are multiple measurements of the same plot set (Beets et al., 2011a).

The initial post-1989 planted forest inventory carried out during the winters of 2007 and 2008 at 246 sites consisted of up to four sample plots per site in a cluster arrangement. The plots were sampled using the methods as described in Payton et al. (unpublished). A second inventory was carried out during the winters of 2011 and 2012 where the centre plot of the earlier established cluster plots was re-measured and additional new plots were established. In total, 342 plots were ground measured from the mapped area of post-1989 planted forest in the second inventory. Importantly, the additional plots in the second inventory addressed a bias in the earlier estimates caused by incomplete sampling of the forest area. This was due to the initial field inventory beginning before the completion of the 2008 land use map. The planted forest inventory shifted from a periodic to a continuous inventory in 2016. The continuous inventory measures around 140 permanent sample plots annually over a five-year re-measurement cycle. The continuous inventory provides annual data on forest management (e.g., harvest age and thinning), natural disturbance and growth that can be incorporated into planted forest carbon stock estimates.

The ground measurements in the post-1989 planted forest inventory are the same as those used in the pre-1990 planted forest inventory described above.

Stock change in the productive area of post-1989 planted forest is estimated using a forest type-specific national yield table approach similar to that described above within pre-1990 planted forest. Plots that are located outside the productive area within the mapped forest boundary are used to provide carbon stock estimates for unstocked areas of post-1989 planted

forests (Paul et al., unpublished(d)). It has been demonstrated in the development of the post-1989 forest yield table that forests planted on grassland are more productive than those planted on forest land (Paul et al., unpublished(c)).

To use all plot measurements described above, a single yield table per plot was developed using the estimated carbon stock at each measurement date. An interpolated estimate is used to provide carbon stock at all ages between the measurement dates. The advantage of the interpolation method is that it maintains the actual carbon stock values at individual measurement dates. Individual yield tables are combined as weighted means in a national yield table for the productive area of post-1989 planted forest (Paul et al., unpublished(d)).

New Zealand plantation forests are actively managed, with thinning and pruning activities undertaken early in the rotation. Most of these activities are completed before trees reach the age of 13 years. Thus, the dead wood and litter pools from these management practices gradually increase leading up to this age. After the age of 13 years, when pruning and thinning cease and decay exceeds inputs, these pools decline. Due to the age-class structure of post-1989 forest in New Zealand, this can be seen as a rapid increase in the dead wood and litter pools over consecutive years.

Quality assurance and quality control

Quality-assurance and quality-control activities were conducted throughout the pre-1990 and post-1989 planted and natural forest data capture and processing steps. These activities were associated with the following: inventory design (Beets et al., 2014b; Brack, unpublished; Moore and Goulding, unpublished); acquisition of raw LiDAR data and LiDAR processing; checking eligibility of plots; independent audits of field plot measurements (Beets and Holt, unpublished); auditing data entry; data processing and modelling; regression analysis and double-sampling procedures (Woollens, unpublished); and investigating LiDAR and ground plot collocation (Brack and Broadley, unpublished). These activities are described in detail below.

Pre-1990 natural forest

During the initial measurement of the natural forest plot network (2002–07), 5 per cent of plots measured in the first field season were randomly selected for audit (Beets and Payton, unpublished). In all subsequent field seasons, data collection followed quality-assurance and quality-control processes, as described in Payton et al. (unpublished). This included on-site quality-control checks of field data and review by senior ecologists. Data were collected in the field and recorded by hand on paper field-sheets. The electronic entry of these data has been subject to ongoing quality assurance and quality control, including line-by-line checking of the transcription of all data used in carbon calculations.

During the re-measurement of the plot network from 2009 to 2014, 10 per cent of plots measured were subject to independent audit. For the current re-measurement of the plot network, this has been reduced to 5 per cent of plots measured. This audit involves a partial re-measure of randomly selected plots, and the assessment of measurements against data quality standards as described in the data collection manual (Ministry for the Environment, unpublished). Up until 2020, entry of data into the electronic database from paper-based plot sheets was subject to quality assurance by the Ministry for the Environment. Line-by-line checks were conducted for 10 per cent of all plots, data are now collected electronically so bypass the need for manual data entry. The data are also subject to further checking for measurement and data entry errors before analysis (Paul et al., 2021).

Post-1989 natural forest

As for pre-1990 natural forest, quality control and quality assurance were undertaken at the data collection, entry and analysis stages.

During field data collection in 2012, 10 per cent of plots were subject to an independent field audit. The audit involved randomly selected sites being re-measured by an audit field team, and the assessment of differences between inventory and audit measurements against set data quality standards as set out in Ministry for the Environment (unpublished). Audit results are described in Beets and Holt (unpublished). The same audit process was conducted for the re-measurement of the post-1989 natural forest plot network in 2019. These results are described in Paul and Dowling (unpublished). Similarly to pre-1990 natural forest, entry of data into the electronic database from paper-based plot sheets is subject to quality assurance by the Ministry for the Environment. Line-by-line checks are conducted for 10 per cent of all plots. Further checks for data entry and measurement were also undertaken before the data analysis stage, as described in Beets et al. (unpublished) and Paul et al. (unpublished(b)).

Pre-1990 planted forest and post-1989 planted forest

Of the planted forest inventory plots, 7.5 per cent are randomly audited without the prior knowledge of the field teams. Plots are fully and partially re-measured, with feedback supplied no later than one month after measurement, to ensure prompt identification of any data collection errors and/or procedural issues. Differences between the inventory and audit measurements are objectively and quantitatively scored. Measurements that exceed predefined tolerances incur incremental demerit points. Demerit severity depends on the size of error and the type of measurement. Special attention is given to the most influential measurements; for example, tree diameter, tree height and the number of trees in a plot. Plots that fail quality control would have to be re-measured (Beets et al., 2011a, 2012a). Following each inventory season, the data collection manual (Herries et al., unpublished) is revised to clarify any potential sources of error or ambiguity.

The inventory data are pre-processed using Scion's Permanent Sample Plot (PSP) system. The PSP system has been programmed to check for erroneous values over a wide range of attributes. The system automatically identifies fields that do not meet predetermined validation rules so these can be repaired manually before plot data are modelled by the Forest Carbon Predictor. The PSP data validation system and the Forest Carbon Predictor model were independently reviewed by Woollens (unpublished). The Forest Carbon Predictor has been validated in Beets et al. (2011b).

Forest land model validations

LUCAS harvest losses versus Ministry for Primary Industries roundwood statistics

The above-ground biomass estimated to be removed from all planted forest destocking (all harvest and deforestation) was compared to the estimated carbon stored in annual Ministry for Primary Industries roundwood removal statistics (Ministry for Primary Industries, 2022; figure A3.2.10). The Ministry for Primary Industries' roundwood volume was converted to tonnes of carbon based on the carbon fractions used in the harvested wood products model (0.21 t C m⁻³ for coniferous and 0.25 t C m⁻³ for non-coniferous timber).


The results show alignment between the two data sources between 2013 and 2020. This is because roundwood volume is now used to estimate the total destocking area over this period. However, the two data sources deviate from 1990 to 2012. The LUCAS above-ground

biomass losses from harvest are greater than those estimated from roundwood volume statistics from 1996 to 2012 and are slightly lower from 1990 to 1994. The LUCAS above-ground biomass estimate may differ from the Ministry for Primary Industries roundwood removal estimate because the latter uses conversion factors for each forestry product to estimate the corresponding roundwood input required to produce the total forest product output. In addition, the Ministry for Primary Industries reports roundwood removal under-bark while the LUCAS estimate is over-bark.

Further work is planned to improve the consistency of forest carbon losses due to harvest and carbon inputs into the harvested wood products pool within the LUCAS model. For this, adjustments might need to be made to some of the input parameters to the planted forest model, such as:

1. the harvest area
2. the average harvest age
3. the assumed proportion of above-ground biomass removed as merchantable volume on harvest
4. the yield tables.

Figure A3.2.10 Comparison of the LUCAS estimate for above-ground biomass removed on planted forest destocking and carbon stored in roundwood production from 1990 to 2021



Note: Roundwood production data sourced from Ministry for Primary Industries, 2022. LUCAS = Land Use and Carbon Analysis System; MPI = Ministry for Primary Industries.

Planted forest age profile

The LUCAS net stocked area planted forest age profile was compared to the NEFD planted forest age profile (Ministry for Primary Industries, 2021; figure A3.2.11). Despite the LUCAS forest age profile using NEFD data as an input, notable differences are evident between the two data sources. This is primarily because the NEFD is based on a survey of forest owners, while the LUCAS planted forest age profile at 2021 is based on a modelled simulation of forest activity from 1990 onwards. The LUCAS forest age profile is also influenced by mapped areas of post-1989 and pre-1990 forest and harvest activity from 1990 to 2021. Details on discrepancies between these two data sources are outlined below.

1. The total mapped area of post-1989 forest is smaller than the area of new planting from 1990 onwards that is reported in the NEFD. As a result, the forest area by age for this type of forest is scaled down relative to the NEFD estimate. This also explains why the LUCAS area is lower from ages 18 to 30.
2. The LUCAS pre-1990 planted forest age profile from ages 0 to 30 is driven by the reported areas harvested and replanted in the Calculation and Reporting Application (CRA) simulation model (with a one-year lag on replanting after harvest). The LUCAS estimates of harvest and replanting are therefore not consistent with the NEFD replanting estimates, resulting in a difference in the forest age profiles.
3. The LUCAS forest age profile from ages 30 onwards represents forests planted before 1990. The LUCAS age profile of these forests follows the same pattern as the NEFD data (which they are based on) but reports a higher area for each age. This is because the LUCAS area for these age groups is greater than the areas reported in the NEFD and is scaled to the total mapped area of pre-1990 planted forest net stocked area.

Figure A3.2.11 Comparison of the planted forest age profile as estimated from LUCAS net stocked area for all planted forest (post-1989 and pre-1990) and the NEFD age profile as at 2021



Note: The age profile starts age 1 and does not include areas of forest that were planted or harvested and awaiting replanting in 2021. LUCAS = Land Use and Carbon Analysis System; NEFD = National Exotic Forest Description.

LUCAS planted forest inventory plot measurements versus yield table values

The yield tables generated by the Forest Carbon Predictor were validated in Beets et al. (2011b). The results indicated a good match between carbon stock and stock change predicted from the Forest Carbon Predictor with plot measurements.

A validation was carried out by LUCAS in 2021 on the carbon stock per hectare values of the yield tables used in this and the previous inventory submission, as suggested by the expert review team during the review of the 2021 submission. A comparison was made between the yield table carbon stocks and the measured plot values from the 2016 to 2020 forest inventory used to generate them. The yield table values were adjusted down by half a year to be more consistent with the period that each plot was measured. The yield tables were then fitted to the ages of the measured plots, to provide a comparison of carbon stock per hectare estimates of the yield tables and the measured plots.


The results of this comparison are shown in table A3.2.12 and figures A3.1.12 to A3.1.13. An area weighted average carbon stock per hectare for each forest type was calculated from plots measured in the 2016 to 2020 planted forest inventory and from adjusted yield table values fitted to the ages of these plots (table A3.2.12). Note that the average carbon stock per hectare of measured plots differs from the results of Paul et al. (unpublished(e)), because these estimates only include plots used to generate the yield tables.

Table A3.2.12 Comparison of the average biomass carbon stock per hectare for each forest type, calculated as the area weighted average of plot measurements and yield table values

Forest type	Plots measured in 2016–20 forest inventory			Yield table values fitted to plot age		Difference t C ha ⁻¹
	t C ha ⁻¹	95% CI	Number of plots	t C ha ⁻¹	95% CI	
Post-1989	168.3	9.5	315	173.4	7.2	-5.1
Pre-1990 – after 1990	96.7	11.0	221	108.7	5.0	-12.0
Pre-1990 – before 1990	252.5	43.9	23	282.3	22.4	-29.8


The average carbon stock per hectare of measured plots in post-1989 forest that were used to generate the yield table was 168.3 ± 9.3 tonnes C ha⁻¹. When fitting the adjusted yield table carbon values to the measured plot ages, the average carbon stock per hectare is 173.4 ± 7.2 tonnes C ha⁻¹. On average for post-1989 forests, the yield table estimates carbon stock per hectare to be 5.1 tonnes C ha⁻¹ higher than the measured plot values. This suggests a relatively good fit and is within the average confidence interval (7.2 tonnes C ha) of the yield table.

Figure A3.2.12 Post-1989 planted forest carbon stock by age estimated from the yield table and plots measured in the 2016 to 2020 forest inventory



The average carbon stock per hectare of measured plots in pre-1990 forest planted after 1990 is 96.7 ± 11 tonnes C ha⁻¹. When fitting the adjusted yield table carbon values to the measured plot ages, the average carbon stock per hectare is 108.7 ± 5 tonnes C ha⁻¹. On average, the yield table estimates carbon stock per hectare to be 12 tonnes C ha⁻¹ higher than the measured plot values for these forests. This suggests the yield table for pre-1990 forest planted after 1990 could be overestimating carbon stocks in this forest type. Figure A3.2.13 indicates this could be partially driven by several low-yield plots aged between 16 and 25 that drag the average carbon stock per hectare down.

Figure A3.2.13 Pre-1990 planted forest – planted after 1990 – carbon stock by age estimated from the yield table and plots measured in the 2016 to 2020 forest inventory




The average carbon stock per hectare of measured plots in pre-1990 forest planted before 1990 is 252.5 ± 44 tonnes C ha⁻¹. When fitting the adjusted yield table carbon values to the measured plot ages, the average carbon stock per hectare is 282.3 ± 22.4 tonnes C ha⁻¹. On average, the yield table estimates carbon stock per hectare to be 29.8 tonnes C ha⁻¹ higher than the measured plot values. This suggests the yield table for pre-1990 forest planted after 1990 could also be overestimating the current carbon stocks in this forest type.

However, the large difference between the average yield table values and plots measured in the most recent five-year inventory (2016 to 2020) is likely because the period specific yield tables were estimated from plots that have since been harvested. It is likely that higher yielding plots have been prioritised in harvest and are no longer represented in the 2016 to 2020 forest inventory. Thus, the plots remaining in the 2016 to 2020 forest inventory may be expected to have lower carbon values on average than the yield table that represents all plots planted before 1990.

Analysis and validation of plot measurements to yield table carbon stock values will continue to be undertaken as more data becomes available.

Figure A3.2.15 Post-1989 planted forest above-ground biomass carbon per hectare estimated from the forest inventory and LUCAS Calculation and Reporting Application model




In pre-1990 planted forests, the average above-ground biomass carbon stock per hectare in the 2016 to 2020 forest inventory was 62.3 ± 19.5 tonnes C ha⁻¹ (derived from Paul et al., unpublished(e)). This is much lower than the estimated carbon stock per hectare in 2018 generated from the LUCAS CRA model used for this submission (111.7 tonnes C ha⁻¹; figure A3.2.16). This discrepancy raises questions around how well the LUCAS CRA model simulation of yield tables, combined with planting and harvest data, estimates carbon stock and stock change in pre-1990 planted forests.

Several factors may contribute to this discrepancy.

1. The estimates from the forest inventory (Paul et al., unpublished(e)) assume all plots that were too young to measure had a carbon stock value of zero. In contrast, the yield tables have estimated carbon stocks for these younger stands. Because the pre-1990 planted forest has a large area of forest that is recently harvested, this is likely to contribute to higher carbon stock per hectare in estimates in the LUCAS model.
2. A comparison of yield table carbon stocks to plot measurements (table A3.2.12) indicates that the yield tables tend to predict higher carbon stocks than measured plots in pre-1990 planted forest. The difference between yield table and plot measurements is more pronounced in older plots (figures A3.2.13 and A3.2.14).
3. A relatively large area of forest considered to be older than 40 years is captured in the NEFD and thus also in the LUCAS age profile. Additionally, the approach used to calculate the forest age profile results in this area of older forest being scaled up from the NEFD estimate, to meet the total net stocked area estimate (figure A3.2.11). In comparison, a lower proportion of plots in this older age group are detected in the planted forest inventory. It is possible that the area of forest in this age range is overestimated, resulting in higher carbon stock per hectare estimates in the LUCAS model.

Further analysis and validation between plot measurements, yield table values and the CRA model output is planned in future inventory submissions, to ensure reliable estimates of carbon stock and stock change from the LUCAS model.

Figure A3.2.16 Pre-1990 planted forest above-ground biomass carbon per hectare estimated from the forest inventory and LUCAS Calculation and Reporting Application model



Natural forest carbon stock change estimates and yield tables

This section contains the natural forest carbon stock change estimates and yield tables used for this submission.

Table A3.2.13 Pre-1990 natural forest – tall forest carbon stocks by year (tonnes C ha⁻¹)¹

Year	Above-ground biomass	Below-ground biomass	Dead wood	Litter	Total biomass
1990	145.1	34	44	22.2	245.4
1991	145.1	34	44	22.2	245.4
1992	145.1	34	44	22.2	245.4
1993	145.1	34	44	22.2	245.4
1994	145.1	34	44.1	22.2	245.4
1995	145.1	34	44.1	22.2	245.4
1996	145	34	44.1	22.2	245.4
1997	145	34	44.1	22.2	245.4
1998	145	34	44.1	22.2	245.4
1999	145	34	44.1	22.2	245.3
2000	145	34	44.1	22.2	245.3
2001	145	34	44.1	22.2	245.3
2002	145	34	44.1	22.2	245.3
2003	144.9	34	44.1	22.2	245.3
2004	144.9	34	44.1	22.2	245.3
2005	144.9	34	44.1	22.2	245.3
2006	144.9	34	44.1	22.2	245.3
2007	144.9	34	44.2	22.2	245.3
2008	144.9	34	44.2	22.2	245.2
2009	144.9	34	44.2	22.2	245.2
2010	144.8	34	44.2	22.2	245.2
2011	144.8	34	44.2	22.2	245.2
2012	144.8	34	44.2	22.2	245.2
2013	144.8	33.9	44.2	22.2	245.2
2014	144.8	33.9	44.2	22.2	245.2

Year	Above-ground biomass	Below-ground biomass	Dead wood	Litter	Total biomass
2015	144.8	33.9	44.2	22.2	245.2
2016	144.8	33.9	44.2	22.2	245.1
2017	144.7	33.9	44.2	22.2	245.1
2018	144.7	33.9	44.2	22.2	245.1
2019	144.7	33.9	44.2	22.2	245.1
2020	144.7	33.9	44.2	22.2	245.1
2021	144.7	33.9	44.3	22.2	245.1

¹ Data derived from Paul et al., 2021.

Table A3.2.14 Pre-1990 natural forest – regenerating forest carbon stocks by year (tonnes C ha⁻¹)¹

Year	Above-ground biomass	Below-ground biomass	Dead wood	Litter	Total biomass
1990	34.1	8.2	10.4	9.7	62.3
1991	34.5	8.3	10.3	9.7	62.8
1992	34.8	8.4	10.3	9.7	63.2
1993	35.2	8.5	10.3	9.7	63.6
1994	35.5	8.5	10.3	9.7	64.1
1995	35.9	8.6	10.3	9.7	64.5
1996	36.2	8.7	10.3	9.7	64.9
1997	36.6	8.8	10.3	9.7	65.3
1998	37	8.9	10.3	9.7	65.8
1999	37.3	8.9	10.2	9.7	66.2
2000	37.7	9	10.2	9.7	66.6
2001	38	9.1	10.2	9.7	67
2002	38.4	9.2	10.2	9.7	67.5
2003	38.7	9.3	10.2	9.7	67.9
2004	39.1	9.4	10.2	9.7	68.3
2005	39.5	9.4	10.2	9.7	68.8
2006	39.8	9.5	10.2	9.7	69.2
2007	40.2	9.6	10.2	9.7	69.6
2008	40.5	9.7	10.1	9.7	70
2009	40.9	9.8	10.1	9.7	70.5
2010	41.2	9.9	10.1	9.7	70.9
2011	41.6	9.9	10.1	9.7	71.3
2012	41.9	10	10.1	9.7	71.7
2013	42.3	10.1	10.1	9.7	72.2
2014	42.7	10.2	10.1	9.7	72.6
2015	43	10.3	10.1	9.7	73
2016	43.4	10.4	10	9.7	73.5
2017	43.7	10.4	10	9.7	73.9
2018	44.1	10.5	10	9.7	74.3
2019	44.4	10.6	10	9.7	74.7
2020	44.8	10.7	10	9.7	75.2
2021	45.2	10.8	10	9.1	75.6

¹ Data derived from Paul, et al., 2021.

Table A3.2.15 Post-1989 natural forest yield table (tonnes C ha⁻¹)¹

Age	Above-ground biomass	Below-ground biomass	Dead wood	Litter	Total biomass
0	1.6	0.4	0.0	0.0	2.0
1	2.5	0.6	0.0	0.0	3.2
2	3.7	0.9	0.0	0.1	4.7
3	5.1	1.3	0.0	0.1	6.5
4	6.6	1.7	0.0	0.1	8.4
5	8.3	2.1	0.0	0.1	10.6
6	10.2	2.6	0.1	0.2	13.0
7	12.2	3.1	0.1	0.2	15.5
8	14.4	3.6	0.1	0.2	18.2
9	16.6	4.2	0.1	0.2	21.1
10	18.9	4.7	0.1	0.3	24.0
11	21.4	5.3	0.1	0.3	27.1
12	23.9	6.0	0.1	0.3	30.2
13	26.4	6.6	0.1	0.4	33.5
14	29.0	7.2	0.1	0.4	36.7
15	31.6	7.9	0.1	0.4	40.0
16	34.2	8.6	0.1	0.4	43.3
17	36.8	9.2	0.1	0.5	46.7
18	39.4	9.9	0.1	0.5	49.9
19	42.0	10.5	0.1	0.5	53.2
20	44.5	11.1	0.1	0.6	56.4
21	47.0	11.8	0.2	0.6	59.5
22	49.4	12.3	0.2	0.6	62.5
23	51.7	12.9	0.2	0.6	65.4
24	53.9	13.5	0.2	0.7	68.2
25	56.0	14.0	0.2	0.7	70.9
26	57.9	14.5	0.2	0.7	73.3
27	59.7	14.9	0.2	0.8	75.6
28	61.4	15.3	0.2	0.8	77.7
29	62.9	15.7	0.2	0.8	79.6
30	64.1	16.0	0.2	0.9	81.3
31	65.2	16.3	0.2	0.9	82.7

¹ Yield table source Paul, et al., unpublished(b).

Table A3.2.16 Post-1989 natural forest yield table (tonnes C ha⁻¹)¹ (for transitions from grassland with woody biomass)

Age	Above-ground biomass	Below-ground biomass	Dead wood	Litter	Total biomass
0	9.4	3.1	0.1	0.6	13.1
1	10.2	3.3	0.1	0.6	14.1
2	11.2	3.5	0.1	0.6	15.4
3	12.4	3.8	0.1	0.6	16.9
4	13.8	4.1	0.1	0.6	18.6
5	15.3	4.5	0.1	0.6	20.5
6	16.9	4.8	0.1	0.6	22.5
7	18.7	5.3	0.1	0.6	24.7
8	20.6	5.7	0.1	0.6	27

Age	Above-ground biomass	Below-ground biomass	Dead wood	Litter	Total biomass
9	22.5	6.2	0.1	0.6	29.5
10	24.6	6.7	0.1	0.6	32
11	26.7	7.2	0.1	0.7	34.7
12	28.9	7.7	0.1	0.7	37.4
13	31.1	8.2	0.1	0.7	40.1
14	33.4	8.7	0.2	0.7	43
15	35.7	9.3	0.2	0.7	45.8
16	37.9	9.8	0.2	0.7	48.6
17	40.2	10.4	0.2	0.7	51.5
18	42.5	10.9	0.2	0.7	54.3
19	44.8	11.4	0.2	0.7	57.1
20	47	12	0.2	0.8	59.9
21	49.1	12.5	0.2	0.8	62.5
22	51.2	13	0.2	0.8	65.1
23	53.2	13.5	0.2	0.8	67.6
24	55.2	13.9	0.2	0.8	70
25	57	14.3	0.2	0.8	72.3
26	58.7	14.7	0.2	0.8	74.4
27	60.3	15.1	0.2	0.8	76.4
28	61.7	15.5	0.2	0.8	78.2
29	63	15.8	0.2	0.9	79.8
30	64.1	16	0.2	0.9	81.3
31	65.1	16.3	0.2	0.9	82.5

¹ Yield table source derived from Paul, et al., unpublished(b).

Planted forest yield tables

This section contains the planted forest yield tables used for this submission.

Table A3.2.17 Pre-1990 'planted before 1990' planted forest yield table (tonnes C ha⁻¹)¹

Age	Above-ground biomass	Below-ground biomass	Dead wood	Litter	Total biomass
0	3.3	0.8	0.26	0	4.36
1	3.44	0.85	0.26	0.01	4.55
2	3.85	0.98	0.26	0.03	5.12
3	5.12	1.4	0.26	0.14	6.92
4	7.89	2.18	0.27	0.45	10.78
5	12.09	3.2	0.35	1.25	16.89
6	17.35	4.41	0.62	2.59	24.96
7	22.58	5.51	1.61	4.5	34.2
8	27.07	6.44	3.36	6.74	43.59
9	32.65	7.62	4.24	8.07	52.58
10	39.45	9.04	4.53	8.65	61.67
11	45.43	10.26	5.97	9.47	71.13
12	52.11	11.62	6.9	9.85	80.48
13	60.3	13.3	6.64	9.65	89.89
14	68.44	14.96	6.78	9.56	99.74
15	76.49	16.61	7.2	9.53	109.83
16	83.65	18.05	8.61	9.66	119.97

Age	Above-ground biomass	Below-ground biomass	Dead wood	Litter	Total biomass
17	91.24	19.59	9.46	9.6	129.89
18	99.67	21.32	9.01	9.35	139.35
19	108.05	23.04	8.66	9.14	148.89
20	116.99	24.9	8.17	8.8	158.86
21	125.8	26.75	7.89	8.53	168.97
22	134.58	28.61	7.69	8.3	179.18
23	143.44	30.5	7.45	8.06	189.45
24	152.08	32.35	7.42	7.89	199.74
25	160.44	34.16	7.51	7.77	209.88
26	168.57	35.93	7.6	7.64	219.74
27	176.58	37.67	7.63	7.53	229.41
28	184.15	39.31	7.94	7.5	238.9
29	191.65	40.96	8.22	7.46	248.29
30	199.64	42.71	8.04	7.34	257.73
31	207.7	44.5	7.87	7.21	267.28
32	215.59	46.24	7.72	7.1	276.65
33	223.37	47.97	7.58	7	285.92
34	231.15	49.73	7.48	6.89	295.25
35	238.78	51.45	7.46	6.79	304.48
36	246.22	53.16	7.52	6.7	313.6
37	253.54	54.85	7.63	6.61	322.63
38	260.76	56.53	7.77	6.52	331.58
39	267.87	58.2	7.94	6.44	340.45
40	274.87	59.86	8.14	6.37	349.24
41	281.75	61.49	8.36	6.3	357.9
42	288.44	63.08	8.59	6.23	366.34
43	294.92	64.64	8.84	6.16	374.56
44	301.19	66.17	9.1	6.09	382.55
45	307.27	67.66	9.37	6.01	390.31
46	313.19	69.13	9.64	5.93	397.89
47	319	70.58	9.92	5.85	405.35
48	324.69	72.01	10.19	5.76	412.65
49	330.24	73.42	10.47	5.68	419.81
50	335.66	74.8	10.74	5.6	426.8
51	340.96	76.17	11.02	5.52	433.67
52	346.13	77.51	11.28	5.44	440.36
53	351.19	78.83	11.54	5.36	446.92
54	356.14	80.13	11.8	5.28	453.35
55	360.99	81.42	12.05	5.21	459.67
56	365.74	82.68	12.29	5.14	465.85
57	370.4	83.93	12.53	5.07	471.93
58	374.97	85.17	12.75	5	477.89
59	379.46	86.39	12.97	4.93	483.75
60	383.91	87.6	13.19	4.87	489.56

¹ Yield table source derived from Paul et al., unpublished(e).

Table A3.2.18 Pre-1990 'planted 1990 onwards' planted forest yield table (tonnes C ha⁻¹)¹

Age	Above-ground biomass	Below-ground biomass	Dead wood	Litter	Total biomass
0	2.47	0.61	0.17	0	3.25
1	2.62	0.67	0.17	0.01	3.47
2	3.05	0.8	0.17	0.04	4.06
3	4.4	1.24	0.17	0.14	5.95
4	7.46	2.1	0.18	0.45	10.19
5	12.31	3.27	0.23	1.19	17
6	18.72	4.71	0.37	2.42	26.22
7	26.21	6.3	0.68	4.07	37.26
8	34.04	7.91	1.38	6.01	49.34
9	42.02	9.53	2.34	7.82	61.71
10	50.69	11.3	2.88	9.05	73.92
11	59.78	13.14	3.33	9.84	86.09
12	69.08	15.03	3.84	10.36	98.31
13	78.86	17.04	4.09	10.58	110.57
14	88.71	19.06	4.51	10.69	122.97
15	98.9	21.17	4.75	10.65	135.47
16	109.56	23.39	4.64	10.44	148.03
17	120.22	25.62	4.63	10.23	160.7
18	130.68	27.83	4.77	10.03	173.31
19	140.96	30.01	4.98	9.83	185.78
20	151.16	32.21	5.21	9.62	198.2
21	161.27	34.43	5.52	9.43	210.65
22	171.31	36.66	5.91	9.25	223.13
23	181.04	38.84	6.62	9.14	235.64
24	190.62	41.01	7.36	9.05	248.04
25	200.3	43.22	7.86	8.91	260.29
26	209.84	45.42	8.41	8.79	272.46
27	219.24	47.61	9	8.68	284.53
28	228.45	49.78	9.64	8.58	296.45
29	237.5	51.93	10.31	8.49	308.23
30	246.35	54.05	11	8.4	319.8
31	254.97	56.14	11.72	8.32	331.15
32	263.37	58.2	12.45	8.24	342.26
33	271.55	60.23	13.19	8.16	353.13
34	279.51	62.24	13.94	8.07	363.76
35	287.28	64.22	14.69	7.98	374.17
36	294.85	66.17	15.43	7.89	384.34
37	302.24	68.09	16.17	7.81	394.31
38	309.47	70	16.89	7.72	404.08
39	316.55	71.88	17.59	7.63	413.65
40	323.5	73.75	18.29	7.55	423.09
41	330.27	75.58	18.96	7.48	432.29
42	336.83	77.36	19.62	7.4	441.21
43	343.18	79.12	20.25	7.32	449.87
44	349.34	80.85	20.86	7.24	458.29
45	355.32	82.54	21.43	7.16	466.45

Age	Above-ground biomass	Below-ground biomass	Dead wood	Litter	Total biomass
46	361.12	84.2	21.98	7.07	474.37
47	366.77	85.83	22.5	6.97	482.07
48	372.26	87.44	22.99	6.88	489.57
49	377.6	89.02	23.46	6.79	496.87
50	382.81	90.58	23.89	6.69	503.97
51	387.89	92.11	24.3	6.6	510.9
52	392.86	93.62	24.68	6.51	517.67
53	397.71	95.1	25.03	6.42	524.26
54	402.46	96.57	25.35	6.34	530.72
55	407.12	98.02	25.65	6.25	537.04
56	411.7	99.45	25.93	6.18	543.26
57	416.19	100.87	26.18	6.1	549.34
58	420.6	102.27	26.41	6.03	555.31
59	424.95	103.65	26.62	5.96	561.18
60	429.27	105.03	26.82	5.89	566.98

¹ Yield table source derived from Paul et al., unpublished(e).

Table A3.2.19 Post-1989 planted forest yield table (tonnes C ha⁻¹)¹

Age	Above-ground biomass	Below-ground biomass	Dead wood	Litter	Total biomass
0	2.26	0.56	0.21	0	3.03
1	2.44	0.63	0.21	0.01	3.29
2	3.08	0.79	0.21	0.04	4.12
3	4.12	1.32	0.22	0.15	5.81
4	6.47	2.48	0.23	0.41	9.59
5	9.12	3.82	0.25	1.14	14.33
6	17.19	5.16	1.3	2.81	26.46
7	27.41	6.52	2.42	5.22	41.57
8	32.93	7.61	3.7	7.75	51.99
9	37.35	8.47	6.41	10.08	62.31
10	41.67	9.34	8.93	11.65	71.59
11	47.59	10.58	10.06	12.12	80.35
12	55.24	12.17	10.05	11.95	89.41
13	63.91	13.95	9.71	11.61	99.18
14	73	15.8	9.39	11.29	109.48
15	82.28	17.68	9.05	10.96	119.97
16	91.66	19.58	8.74	10.65	130.63
17	101.16	21.52	8.47	10.34	141.49
18	110.76	23.5	8.19	10.03	152.48
19	120.24	25.45	8.02	9.74	163.45
20	129.61	27.4	7.89	9.46	174.36
21	138.87	29.36	7.64	9.16	185.03
22	148.16	31.34	7.28	8.89	195.67
23	157.58	33.36	6.91	8.66	206.51
24	167.01	35.38	6.3	8.46	217.15
25	176.15	37.36	7.13	8.35	228.99
26	185.14	39.33	8.68	8.29	241.44
27	194.19	41.33	7.13	8.1	250.75

Age	Above-ground biomass	Below-ground biomass	Dead wood	Litter	Total biomass
28	203.14	43.32	4.84	7.82	259.12
29	211.97	45.3	4.72	7.57	269.56
30	220.61	47.25	4.96	7.39	280.21
31	229.06	49.18	4.88	7.25	290.37
32	237.3	51.08	4.82	7.12	300.32
33	245.35	52.96	6.33	7.03	311.67
34	253.22	54.81	8.11	6.99	323.13
35	260.89	56.63	8.57	6.95	333.04
36	268.39	58.43	8.99	6.87	342.68
37	275.72	60.2	9.41	6.79	352.12
38	282.89	61.95	9.83	6.7	361.37
39	289.92	63.69	10.26	6.62	370.49
40	296.83	65.41	10.7	6.53	379.47
41	303.6	67.1	11.13	6.46	388.29
42	310.19	68.75	11.56	6.38	396.88
43	316.6	70.38	11.99	6.31	405.28
44	322.83	71.97	12.41	6.23	413.44
45	328.89	73.54	12.83	6.15	421.41
46	334.77	75.08	13.24	6.07	429.16
47	340.5	76.59	13.64	5.99	436.72
48	346.08	78.08	14.02	5.91	444.09
49	351.52	79.54	14.4	5.82	451.28
50	356.83	80.98	14.76	5.74	458.31
51	362.02	82.4	15.11	5.66	465.19
52	367.09	83.79	15.44	5.58	471.9
53	372.06	85.17	15.76	5.5	478.49
54	376.93	86.53	16.07	5.43	484.96
55	381.71	87.87	16.37	5.35	491.3
56	386.39	89.19	16.65	5.28	497.51
57	390.99	90.5	16.92	5.22	503.63
58	395.52	91.8	17.18	5.15	509.65
59	399.97	93.08	17.42	5.09	515.56
60	404.39	94.35	17.66	5.03	521.43

¹ Yield table source derived from Paul et al., unpublished(e).

A3.2.6 Harvested wood products – exported raw material methodologies

Export market activity data and half-lives

The weighted half-lives applied to sawnwood, wood panels and paper are calculated from the sub-product lifetimes reported by the Ministry for Primary Industries (2016) and Wakelin and Kimberley (unpublished). Sub-product half-lives and their proportions for each export market are summarised in tables A3.2.20 to A3.2.22 based on data collected in 2015.

Table A3.2.20 Harvested wood product type, waste and fuel product type, exported volume in 2015 and assumed half-lives for China

Product	Sub-product	Waste/fuel product	Volume (million m ³)	Half-life
PANEL	Appearance plywood		0.1039604	25
PANEL	Construction plywood	Panel (recycled)	1.1435644	2.5
PANEL		Burned	1.1435644	0.5
PANEL	Packaging plywood		0.2079208	3
	Plymill residue	Burned	0.0519802	0
PAPER		Pulp	0.1559406	2
PANEL		Particle board	0.1559406	25
PANEL		MDF	0.1559406	25
SAWN	Plymill core		0.2079208	2
SAWN	Appearance lumber	Remanufactured	0.9356436	35
SAWN	Construction lumber	Panel (recycled)	1.2475248	2.5
		Burned	1.2475248	0.5
SAWN	Packaging lumber		1.1435644	3
	Slabwood	Burned	0.2079208	0
PAPER		Pulp	1.1435644	2
PANEL		Particle board	0.2079208	25
PANEL		MDF	0.2079208	25
	Sawdust	Burned	0.1559406	0
		Pellets	0.519802	0
PANEL		Particle board	0.1559406	25

Table A3.2.21 Harvested wood product type, waste and fuel product type, exported volume in 2015 and assumed half-lives for South Korea

Product	Sub-product	Waste/fuel product	Volume (million m ³)	Half-life
PANEL	Construction plywood	Panel (recycled)	0.1841	25.5
PANEL		Burned	0.0526	0.5
PANEL	Appearance plywood		0.1052	25
PANEL	Plymill residue	MDF	0.2104	25
SAWN	Appearance lumber		0.0263	35
SAWN	Construction lumber	Particle board	0.6838	25.5
SAWN		Burned	0.1841	0.5
SAWN	Packaging lumber		0.3682	3
PANEL	Slabwood	MDF	0.526	25
	Sawdust	Agriculture	0.1841	0
		Burned	0.0526	0
PANEL	MDF		0.0526	25

Table A3.2.22 Harvested wood product type, waste and fuel product type, exported volume in 2015 and assumed half-lives for India

Product	Sub-product	Waste/fuel product	Volume (million m ³)	Half-life
SAWN	Construction lumber		0.432	0.5
SAWN	Packaging lumber	Export	0.352	3
SAWN		Domestic	0.144	0.5
PANEL	Blockboard		0.208	7
	Slabwood	Fuel	0.224	0
PANEL	Sawdust	Particleboard	0.048	25
		Fuel	0.192	0

A3.2.7 Biomass burning detailed methodology

Wildfire

Wildfires induced by natural disturbances (e.g., lightning) are estimated to account for only 0.1 per cent of burning in the *Grassland* and *Forest land* categories in New Zealand (Doherty et al., unpublished; Wakelin, unpublished(b)). No distinction is made between data collected on anthropogenic and natural wildfire events. Given the small incidence of natural-disturbance-induced wildfires in New Zealand, this is not regarded as a significant source of error.

A single weighted biomass density is used to estimate non-CO₂ emissions from wildfire in the *Forest land remaining forest land* category. Wildfire activity data were attributed to each category by the proportion of forest type estimated to be burned over the time series until 2007. For 2007 to 2016, area data available in the wildfire database were split into natural and planted forests (Wakelin, unpublished(d)). The split before 2007 assumed 87.5 per cent to be planted forest and the remainder to be natural forest (Wakelin, unpublished(g)). The planted forest activity data were further split into pre-1990 forest and post-1989 forest by the proportion of area each forest type makes up of the total planted forest area. In planted forest, it is assumed that the carbon stock affected by wildfire is equivalent to the carbon stock at the average stand age in each forest type (Wakelin, unpublished(d)). The individual forest type estimates that make up the single weighted figure are derived from the national forest plot network described in section A3.2.5, 'National forest inventory'.

From 2017 onwards, Fire and Emergency New Zealand has supplied mapped wildfire activity data. This has enabled the use of the land use map to infer the type of land cover and category of forest burnt. Consequently, since 2017 areas of pre-1990 and post-1989 planted and natural forest affected by wildfire are now derived from the land use map. It is still assumed that the carbon stock affected by wildfire is equivalent to the carbon stock at the average stand age in each forest type.

In planted forest, it is assumed that the carbon stock affected by wildfire is equivalent to the carbon stock at the average stand age in each forest type (Wakelin, unpublished(d)). The individual forest type estimates that are used to calculate the single weighted figure are derived from the national forest plot network described in section A3.2.5, 'National forest inventory'.

An estimate for wildfire in *Land converted to grassland* is provided in the inventory. The activity data for wildfire in *Grassland* are attributed to the *Land converted to* and *Land remaining* categories by the proportion each category makes up of the total area.

Controlled burning

Activity data (area of land-use change) for controlled burning for *Forest land* is estimated based on a survey carried out in 2011. Activity data for *Grassland with woody biomass* converted to forest are based on annual land-use changes and an estimate of area burned from the survey of forest owners.

The survey also provided data on the burning of post-harvest slash before restocking. This activity was found to occur mainly as a training exercise for wildfire control or for the clearing of slash heaps on skid sites. The data indicated that 0.8 per cent of restocked area was burned each year in recent years. This estimate was combined with two earlier estimates of controlled burning in planted forest (Forest Industry Training and Education Council, 2005; Robertson,

1998) to provide activity data throughout the time series. It is assumed that 1.6 per cent of restocked area was burned from 1990 to 1997. From 1997, the area burned declines linearly to 0.8 per cent, which is used from 2005 onwards (Wakelin, unpublished(e)).

Activity data are combined with an emission factor derived from the pre-1990 planted forest (planted before 1990) carbon-yield table to estimate emissions from the burning of post-harvest slash (harvest residue) on *Forest land*. The harvest residue is calculated by subtracting the amount of above-ground biomass that is taken off site as logs (70 per cent) from the total above-ground biomass predicted at the age of 28 years (the average harvest age in New Zealand). Below-ground biomass is assumed not to burn. The IPCC default combustion proportion for the burning of harvest residue in non-eucalypt temperate forest (0.62) is applied to estimate emissions from this activity (table 2.6, IPCC, 2006a).

An estimate is provided for burning of post-harvest residues associated with deforestation in the National Inventory Report. No information is available on the extent of burning associated with deforestation in New Zealand. Therefore, it is assumed that 30 per cent of conversions involve burning to clear residues. The IPCC default combustion proportion for the burning of harvest residue in non-eucalypt temperate forest (0.62) is applied to category-specific emission factors to estimate emissions from this activity. The emission factor excludes the proportion of logs taken off site (70 per cent of above-ground biomass) and is taken from the plot-network-derived yield tables by forest type at the average age of harvest in New Zealand.

Carbon dioxide emissions from controlled burning in planted forests are captured at the time of conversion or harvest.

The burning of tussock (*Chionochloa* spp.) grassland occurs in the South Island of New Zealand for pasture renewal and weed control. The amount of burning has been decreasing steadily over the past 50 years, as a result of changes in lease tenure and a reduction in grazing pressure. The tussock burning data are sourced from consents under the Resource Management Act 1991 for activities that occurred between 1990 and 2004. Stats NZ provides these data from 2005 because burning became a permitted activity under the Act in some regions (Thomas et al., 2011).

Current practice in New Zealand is to burn in damp spring conditions, reducing the amount of biomass consumed by fire. To reflect this, a country-specific combustion factor of 0.619 is applied (spring burn carbon fractions averaged across two sites (Payton and Pearce, 2009)) to a country-specific biomass density of 28 (t dm ha⁻¹). The ratio of biomass density to carbon lost upon burning is 0.45 (as cited in Thomas et al., 2011).

An estimate for controlled burning in *Grassland remaining grassland (Grassland with woody biomass)* is provided in the inventory. The activity data are sourced from Stats NZ's Agricultural Production survey. The activity data are combined with an emission factor derived from the national forest plot network to estimate non-CO₂ emissions from burning associated with the clearing of vegetation for pasture regeneration. Below-ground biomass is assumed not to burn. The New Zealand-specific default combustion proportion for the burning of shrublands of 0.7 (Wakelin, unpublished(a),(h)) is then applied to estimate emissions from this activity (table 2.6, IPCC, 2006a).

Different emission factors derived from the LUCAS plot network are used for wildfire and controlled burning on *Grassland with woody biomass* in the inventory. The differences are due to the vegetation that is typically converted to forest, which is generally of a lesser stature when compared with other shrubland (Wakelin and Beets, unpublished).

A3.2.8 Uncertainty analysis for the LULUCF sector

All uncertainties associated with activity and emission factors are combined to provide an overall uncertainty estimate for total LULUCF emissions and for each subcategory. For the LULUCF sector, all uncertainties are combined using approach 1 in the IPCC 2006 General Guidance and Reporting: the propagation of error (IPCC, 2006b).

Methods used to calculate uncertainty in Forest land

Uncertainty in net CO₂ emissions from *Forest land* are calculated using several inputs, including uncertainty in mapping, uncertainty in carbon stocks and uncertainty in carbon stock change.

Mineral SOC stocks have an estimated uncertainty of ± 7.9 per cent in pre-1990 natural forest, ± 12.5 per cent in pre-1990 planted forest and ± 10.4 per cent for post-1989 natural and planted forest, as calculated from the Tier 2 method estimates of SOC. Uncertainties in soil carbon stock change are calculated for each specific land use transition (see section A3.2.4, 'Mineral soils').

The uncertainty associated with biomass losses on conversion to *Forest land* is calculated from the carbon stocks in the *from land use* category. Details on the uncertainty associated with biomass gains on conversion to *Forest land* and biomass losses associated with measured carbon stock change losses due to land-use change events are outlined for each forest subcategory below.

Pre-1990 natural forest

The estimates for carbon stock and carbon stock change in pre-1990 natural forests were adapted from Paul et al. (2021). Carbon stocks in 2021 are estimated to be 75.6 ± 14.0 tonnes C ha⁻¹ in regenerating natural forest and 245.1 ± 15.3 tonnes C ha⁻¹ in tall natural forest, with an associated uncertainty at the 95 per cent confidence interval of ± 27.3 per cent and ± 20.9 per cent, respectively. The uncertainty associated with carbon stock estimates for the current reporting year were propagated through time using equation 3.2 from IPCC General Guidance and Reporting (IPCC, 2006b). These estimates of carbon stock per hectare are used as emission factors to calculate emissions for land converted from pre-1990 natural forest.

It is possible that the average carbon stock per hectare estimates for tall and regenerating forest, across the entire pre-1990 natural forest estate, are not representative of the forest that has actually been deforested. Consequently, there is additional uncertainty in the estimate of carbon losses from the deforestation of pre-1990 natural forest, due to a potential lack of representativeness in the data. To account for this potential lack of representativeness in the data, expert judgement was made to add an additional component of 20.0 per cent uncertainty to the carbon stocks, to provide an overall uncertainty for carbon losses on deforestation.

Carbon stock change was estimated to be 0.43 ± 0.51 tonnes C ha⁻¹ yr⁻¹ in regenerating natural forest and -0.01 ± 0.19 tonnes C ha⁻¹ yr⁻¹ for tall natural forest, with an associated uncertainty at the 95 per cent confidence interval of ± 119.5 per cent and $\pm 1,678.6$ per cent, respectively. The uncertainty in carbon stock change is applied to carbon gains or losses within the pre-1990 natural forest category. Further information on the inputs used to calculate uncertainty associated with pre-1990 natural forest is outlined in table A3.2.23.

Table A3.2.23 Uncertainty in New Zealand’s 2021 carbon estimates from pre-1990 natural forest (including land in transition)

Variable	Uncertainty at a 95% confidence interval (%)
Activity data	
Uncertainty in land area	±5.0
Emission factors	
Uncertainty in tall forest biomass carbon stocks	±20.9
Uncertainty in regenerating forest biomass carbon stocks	±27.3
Uncertainty in tall forest biomass carbon change	±1,678.6
Uncertainty in regenerating forest biomass carbon change	±119.5
Uncertainty in soil carbon stocks	±7.9
Uncertainty introduced into net emissions for LULUCF	±17.1

Note: Land area includes land in transition in 2021. The activity data and combined emission factor uncertainty are weighted values and have been calculated using equations 3.1 and 3.2 from IPCC General Guidance and Reporting (IPCC, 2006b).

Post-1989 natural forest

The average carbon stock in post-1989 natural forest is estimated to be 38.55 ± 10.23 tonnes C ha⁻¹ in 2021 with an associated uncertainty of ±27.0 per cent. The average carbon stock change in post-1989 natural forest is estimated to be 2.48 ± 1.1 tonnes C ha⁻¹ yr⁻¹, with an associated uncertainty of ±44.8 per cent. The uncertainty in carbon stocks is applied to losses from deforestation, while the uncertainty in carbon stock change is applied to carbon gains from forest growth. The uncertainty in the estimates of post-1989 natural forest for the 2023 submission is provided in table A3.2.24.

Table A3.2.24 Uncertainty in New Zealand’s 2021 carbon estimates from post-1989 natural forest (including land in transition)

Variable	Uncertainty at a 95% confidence interval (%)
Activity data	
Uncertainty in land area	±8.0
Emission factors	
Uncertainty in biomass carbon stocks (losses)	±27.0
Uncertainty in biomass carbon stock change (gains)	±44.8
Uncertainty in soil carbon stocks	±10.4
Uncertainty introduced into net emissions for LULUCF	±1.2

Note: Land area includes land in transition in 2021. The activity data and combined emission factor uncertainty are weighted values and have been calculated using equations 3.1 and 3.2 from IPCC General Guidance and Reporting (IPCC, 2006b).

Pre-1990 planted forest

The uncertainty in carbon losses applied to New Zealand’s pre-1990 forest biomass carbon stocks is ±21.2 per cent at the 95 per cent confidence interval, while the uncertainty in carbon stock change (carbon gains) is ±11.3 per cent (see table A3.2.25). The uncertainty in carbon stocks is applied to carbon losses that occur from harvesting and deforestation and the uncertainty in carbon stock change applies to carbon gains from forest growth. These uncertainty estimates take into account the area-weighted uncertainty in carbon stocks for each age in the yield table (Paul and Wakelin, unpublished) and the associated uncertainty in estimating the forest age profile and harvest age profile.

The uncertainty in the carbon estimates of pre-1990 planted forest for the 2020 Inventory submission is provided in table A3.2.25.

Table A3.2.25 Uncertainty in New Zealand’s 2021 carbon estimates from pre-1990 planted forest (including land in transition)

Variable	Uncertainty at a 95% confidence interval (%)
Activity data	
Uncertainty in land area	±5.0
Emission factors	
Uncertainty in planted forest biomass carbon stocks (losses)	±21.2
Uncertainty in planted forest biomass carbon stock change (gains)	±11.3
Uncertainty in unstocked forest biomass carbon stocks	±146.0
Uncertainty in riparian forest biomass carbon stocks	±75.0
Uncertainty in soil carbon stocks	±12.3
Uncertainty introduced into net emissions for LULUCF	±55.5

Note: Land area includes land in transition in 2021. The activity data and combined emission factor uncertainty are weighted values and have been calculated using equations 3.1 and 3.2 from IPCC General Guidance and Reporting (IPCC, 2006b).

Post-1989 planted forest

The uncertainty in carbon losses applied to New Zealand’s post-1989 planted forest biomass carbon stocks is ±20.5 per cent, while the uncertainty in carbon stock change (carbon gains) is ±8.9 per cent (see table A3.2.26). The uncertainty in carbon stocks is applied to carbon losses from harvesting and deforestation and the uncertainty in carbon stock change applies to carbon gains from forest growth. These uncertainty estimates take into account the area weighted uncertainty in carbon stocks for each age in the yield table (Paul et al., unpublished(d)) and the associated uncertainty in estimating the forest age profile and harvest age profile.

The uncertainty in the estimates of post-1989 planted forest for the 2021 Inventory submission is provided in table A3.2.26.

Table A3.2.26 Uncertainty in New Zealand’s 2021 carbon estimates from post-1989 planted forest (including land in transition)

Variable	Uncertainty at a 95% confidence interval (%)
Activity data	
Uncertainty in land area	±8.0
Emission factors	
Uncertainty in planted forest biomass carbon stocks (losses)	±20.5
Uncertainty in planted forest biomass carbon stock change (gains)	±8.9
Uncertainty in unstocked forest biomass carbon stocks	±72.0
Uncertainty in riparian forest biomass carbon stocks	±75.0
Uncertainty in soil carbon stocks	±10.4
Uncertainty introduced into net emissions for LULUCF	±19.5

Note: Land area includes land in transition in 2021. The activity data and combined emission factor uncertainty are weighted values and have been calculated using equations 3.1 and 3.2 from IPCC General Guidance and Reporting (IPCC, 2006b).

Methods used to calculate uncertainty in Cropland

The uncertainty in mapping *Cropland* is ± 8.0 per cent (see table A3.2.27).

New Zealand uses IPCC default values for biomass accumulation in annual cropland. For perennial cropland, a New Zealand-specific emission factor is used (Davis and Wakelin, unpublished). Because the perennial and annual cropland emission factors are based on only a limited number of biomass studies, the uncertainty in these figures is estimated as ± 75.0 per cent (table 5.9, IPCC, 2006a). The uncertainty associated with biomass losses on conversion to *Cropland* is calculated from the carbon stocks in the *from land use* category. Mineral soil organic carbon stocks have an estimated uncertainty of ± 9.7 per cent in annual cropland and ± 14.1 per cent in perennial cropland, as calculated from the Tier 2 method estimates of SOC (see table A3.2.27). Uncertainties in soil carbon stock change are calculated for each specific land use transition (section A3.2.4, 'Mineral soils').

For organic soils, New Zealand uses IPCC default values for annual and perennial cropland. The uncertainty associated with the IPCC default values is 90 per cent (based on table 2.3, IPCC, 2006a).

As shown in table A3.2.27, while uncertainty in activity data is low, the uncertainty in the IPCC default variables dominates the overall uncertainty in the estimate provided by New Zealand.

Table A3.2.27 Uncertainty in New Zealand's 2020 Cropland carbon estimates (including land in transition)

	Uncertainty at a 95% confidence interval	
	Annual cropland (%)	Perennial cropland (%)
Activity data		
Uncertainty in land area	± 8.0	± 8.0
Emission factors		
Uncertainty in biomass carbon stocks	± 75.0	± 75.0
Uncertainty in mineral soil carbon stocks	± 9.7	± 14.1
Uncertainty introduced into net emissions for LULUCF	± 1.2	± 0.3

Methods used to calculate uncertainty in Grassland

The uncertainty in mapping *Grassland* is ± 8.0 per cent for *High and Low producing grassland* and ± 83.0 per cent for *Grassland with woody biomass* (table A3.2.28).

New Zealand uses IPCC default values for biomass accumulation in high producing and low producing grassland. The uncertainty in these figures is given as ± 75.0 per cent (table 6.4, IPCC, 2006a). A New Zealand-specific value derived from the LUCAS national forest plot network is used for biomass accumulation in *Grassland with woody biomass*. Due to the uncertainty in this estimate, the IPCC default value of ± 75.0 is also applied to *Grassland with woody biomass*. The uncertainty associated with biomass losses on conversion to *Grassland* is calculated from the carbon stocks in the *from land use* category.

Mineral SOC stocks have an estimated uncertainty of ± 5.8 per cent in high producing grassland and ± 7.3 per cent for both low producing grassland and *Grassland with woody biomass*, as calculated from the Tier 2 method estimates of SOC (see table A3.2.28). Uncertainties in soil carbon stock change are calculated for each specific land use transition (section A3.2.4, 'Mineral soils').

For organic soils, New Zealand uses IPCC default values for annual and perennial cropland. The uncertainty associated with the IPCC default values is ± 90.0 per cent (table 2.3, IPCC, 2006a).

Table A3.2.28 Uncertainty in New Zealand’s 2021 carbon estimates for the *Grassland* category (including land in transition)

Land use	Uncertainty at a 95% confidence interval		
	High producing (%)	Low producing (%)	With woody biomass (%)
Activity data			
Uncertainty in land area	±8.0	±8.0	±83.0
Emission factors			
Uncertainty in biomass carbon stocks	±75.0	±75.0	±75.0
Uncertainty in soil carbon stocks	±5.8	±7.3	±7.3
Uncertainty introduced into net emissions for LULUCF	±0.6	±5.3	±0.9

Note: Uncertainty in biomass carbon stocks for *Grassland with woody biomass* is estimated using the IPCC default uncertainty value because an independent estimate of uncertainty for this category is not available.

Methods used to calculate uncertainty in Wetlands

The uncertainty in mapping *Wetlands* is ±33.0 per cent (see table A3.2.29).

The uncertainty associated with biomass losses on conversion to *Wetlands* is calculated from the carbon stocks in the *from land use* category. A New Zealand-specific value derived from a recent literature review (Easdale et al., unpublished) is used for biomass accumulation in *Wetlands – vegetated*. There is assumed to be no gain in carbon biomass on conversion to *Wetlands – open water*.

The uncertainty for mineral SOC stocks in vegetated wetlands is ±12.3 per cent. An estimated uncertainty of ±90 per cent is used for mineral SOC stocks in open water wetlands. Uncertainties in soil carbon stock change on conversion to and from vegetated wetland are calculated for each specific land use transition (section A3.2.4, ‘Mineral soils’). An estimated uncertainty of ±100.0 per cent is applied to all land use conversion to and from open water wetlands (apart from *Other land*, which applies a higher uncertainty, see section A3.2.4, ‘Mineral soils’).

The uncertainty in the emission factor for peat extracted for horticultural use is ±90.0 per cent, the default IPCC value provided in the 2006 IPCC Guidelines (IPCC, 2006a).

Because emissions from *Wetlands* are very small, the uncertainty introduced into the total net emissions for LULUCF is also very small.

Table A3.2.29 Uncertainty in New Zealand’s 2021 carbon estimates for the *Wetlands* category (including land in transition)

Variable Land use	Uncertainty at a 95% confidence interval	
	Wetlands – vegetated (%)	Wetlands – open water (%)
Activity data		
Uncertainty in land area	±33.0	±33.0
Emission factors		
Uncertainty in biomass carbon stocks	±150	NA
Uncertainty in mineral soil carbon stocks	±12.3	±90.0
Uncertainty in organic soil carbon stocks (on-site CO ₂ emissions from peat extraction)	±90.0	NA
Uncertainty introduced into net emissions for LULUCF	±0.0	±0.0

Note: NA = not applicable. The activity data and combined emission factor uncertainty are weighted values and have been calculated using equation 3.2 from IPCC General Guidance and Reporting (IPCC, 2006b).

Methods used to calculate uncertainty in Settlements

The uncertainty in mapping *Settlements* is ± 22.0 per cent (see table A3.2.30).

The uncertainty associated with biomass losses on conversion to *Settlements* is calculated from the carbon stocks in the *from land use* category. There is assumed to be no gain in carbon biomass on conversion to *Settlements*.

Soil organic carbon stocks have an estimated uncertainty of ± 95.0 per cent, with a soil carbon stock change from all conversions to and from settlements having an uncertainty of ± 100.0 per cent (apart from *Other land*, which applies a higher uncertainty, section A3.2.4, 'Mineral soils').

Table A3.2.30 Uncertainty in New Zealand's 2021 carbon estimates for the *Settlements* category (including land in transition)

Variable	Uncertainty at a 95% confidence interval (%)
Activity data	
Uncertainty in land area	± 22.0
Emission factors	
Uncertainty in biomass carbon stocks	NA
Uncertainty in soil carbon stocks	± 95.0
Uncertainty introduced into net emissions for LULUCF	± 0.4

Note: NA = not applicable. The activity data and combined emission factor uncertainty are weighted values and have been calculated using equation 3.2 from IPCC General Guidance and Reporting (IPCC, 2006b).

Methods used to calculate uncertainty in Other land

The uncertainty associated with biomass losses on conversion to *Other land* is calculated from the carbon stocks in the *from land use* category. There is assumed to be no gain in carbon biomass on conversion to *Other land*.

Soil mineral organic carbon stocks have an uncertainty ± 70.7 , as calculated from the Tier 2 method estimates of SOC (see table A3.2.31). Uncertainties in soil carbon stock change on conversion to and from *Other land* are calculated for each specific land use transition (section A3.2.4, 'Mineral soils').

Table A3.2.31 Uncertainty in New Zealand's 2021 carbon estimates for the land use category *Other land* (including land in transition)

Variable	Uncertainty at a 95% confidence interval (%)
Activity data	
Uncertainty in land area	± 22.0
Emission factors	
Uncertainty in biomass carbon stocks	NA
Uncertainty in soil carbon stocks	± 70.7
Uncertainty introduced into net emissions for LULUCF	± 0.3

Note: NA = not applicable. The activity data and combined emission factor uncertainty are weighted values and have been calculated using equation 3.2 from IPCC General Guidance and Reporting (IPCC, 2006b).

Methods used to calculate uncertainty in Harvested wood products

Uncertainty in the *Harvested wood products* estimates is introduced by activity data, conversion factors and decay parameters.

Additions to the *Harvested wood products* carbon pool are calculated by multiplying wood product production volume or weight by product-specific wood density and carbon fractions. Uncertainties for these factors can be combined using approach 1 for combining uncertainties (IPCC, 2006b).

Losses from the *Harvested wood products* pool are estimated using first order decay functions, based on k factors (discard rates) derived from each product's assumed half-life. The same rule for combining uncertainties cannot be used because the k factor is not multiplied by the other factors.

For *Harvested wood product* exports, the following parameters are considered in the uncertainty calculation:

- uncertainty in export log production
- uncertainty in allocation to export market
- uncertainty in mill conversion to products
- uncertainty in wood density
- uncertainty in carbon content.

The *Harvested wood products* category provides the second-greatest contribution to uncertainty in the LULUCF sector. This is driven by large removals because carbon in harvested timber is transferred to this pool and the high uncertainty associated with the end-use and discard rates of New Zealand wood. Uncertainty limits for *Harvested wood products* data and parameters are given in table A3.2.32. Uncertainty in New Zealand's 2021 carbon estimates from emissions associated with *Harvested wood products* is provided in table A3.2.33.

Table A3.2.32 Uncertainty in *Harvested wood products* data and parameters

Parameter	% uncertainty	Origin
Harvested wood products production, import and export data	±15.0	IPCC default (table 12.6, IPCC, 2006a)
Product volume to weight factors	±10.0	Country specific (Wakelin et al., 2020)
Oven dry product weight to carbon weight	±5.0	Country specific (Wakelin et al., 2020)
Discard rate, domestic	±50.0	Country specific (Wakelin et al., 2020)
Discard rate, export	±90.0	Country specific (Wakelin, unpublished(f))

Table A3.2.33 Uncertainty in New Zealand's 2021 carbon estimates from emissions associated with *Harvested wood products*

Variable	Uncertainty at a 95% confidence interval (%)
Activity data	
Uncertainty in activity data	±15.0
Emission factors	
Domestic production	±51.3
Export raw materials	±91.9
Total domestically milled and exported products uncertainty	±68.2
Uncertainty introduced into net emissions for LULUCF	±28.0

Methods used to calculate uncertainty for nitrous oxide emissions from soils

Uncertainties for direct N₂O emission factors associated with drainage, as well as indirect N₂O emissions from leaching and runoff, are sourced from chapter 11 of the IPCC 2006 Guidelines (IPCC, 2006a). Tables 11.1 and 11.3 of the 2006 IPCC Guidelines give an uncertainty range. The relative uncertainty is then adjusted using the approach for dealing with asymmetric uncertainties, as described by equations 3.3 and 3.4, chapter 3 of the IPCC General Guidance and Reporting (IPCC, 2006b). For N₂O emissions associated with nitrogen mineralisation, an uncertainty of ±80.0 per cent is applied. Uncertainty associated with the variable used to calculate N₂O emissions from land-use change is summarised in table A3.2.34.

Table A3.2.34 Uncertainty in New Zealand’s 2020 estimates from nitrous oxide (N₂O) emissions associated with land-use change

Source	Uncertainty at a 95% confidence interval (%)
Direct N₂O emissions from nitrogen mineralisation	
Activity data	±8.0
Soil carbon	±30.0
C:N ratio	±15.0
N ₂ O emission factor	±80.0
Direct N₂O emissions from drainage	
Activity data	±33.0
N ₂ O emission factor	±80.0
Indirect N₂O emissions from leaching and runoff	
Activity data	±8.0
N ₂ O emission factor	±75.0
Fraction of leaching	±56.0
Uncertainty introduced into net emissions for LULUCF	±0.6

Note: C:N = carbon:nitrogen.

Disaggregated uncertainty analysis for the LULUCF sector

This section contains the disaggregated uncertainty analysis for the LULUCF sector. This additional information has been provided as a result of the review of New Zealand’s 2010 inventory (2012 submission). One of the recommendations of the review was that New Zealand provides “a detailed disaggregated assessment of uncertainty, as well as the aggregated uncertainty associated with the LULUCF sector, consistent with the Intergovernmental Panel on Climate Change (IPCC) good practice guidance for LULUCF”. This information is provided in table A3.2.35.

Table A3.2.35 Uncertainty analysis for the LULUCF sector

IPCC category	Gas	1990 emissions or removals (kt CO ₂ -e)	2021 emissions or removals (kt CO ₂ -e)	Activity data uncertainty (%)	Emission factor / estimation parameter uncertainty (biomass) (%)	Combined uncertainty (%)	Contribution to variance by category in 2021	Type A sensitivity (%)	Type B sensitivity (%)	Uncertainty in trend in LULUCF emissions introduced by emission factor/ estimation parameter uncertainty (%)	Uncertainty in trend in LULUCF emissions introduced by activity data uncertainty (%)	Uncertainty introduced into the trend in total LULUCF emissions (%)
Pre-1990 natural forest	CO ₂	-1,373.5	-1,369.9	0	263.4	263.4	0.0293	0.3	6.8	0.9	0.0	0.0
Pre-1990 planted forest	CO ₂	-18,059.9	-7,822.5	0	149.6	149.6	0.3079	54.4	38.8	81.4	0.0	66.2
Post-1989 planted forest	CO ₂	153.8	-6,083.9	0	67.7	67.7	0.0382	31.0	30.2	21.0	0.0	4.4
Post-1989 natural forest	CO ₂	3.8	-718.5	0	35.4	35.4	0.0001	3.6	3.6	1.3	0.0	0.0
Cropland perennial	CO ₂	126.1	65.4	0	141.9	141.9	0.0000	0.3	0.3	0.5	0.0	0.0
Cropland annual	CO ₂	342.9	306.8	0	81.0	81.0	0.0001	0.3	1.5	0.2	0.0	0.0
Grassland low producing	CO ₂	133.9	1,784.3	0	63.1	63.1	0.0029	8.2	8.8	5.1	0.0	0.3
Grassland high producing	CO ₂	439.4	583.6	0	21.9	21.9	0.0000	0.6	2.9	0.1	0.0	0.0
Grassland with woody biomass	CO ₂	68.7	340.2	0	57.7	57.7	0.0001	1.3	1.7	0.8	0.0	0.0
Wetlands – open water	CO ₂	-18.0	-2.2	0	211.2	211.2	0.0000	0.1	0.0	0.2	0.0	0.0
Wetlands – vegetive non-forest	CO ₂	0.1	-4.5	0	53.6	53.6	0.0000	0.0	0.0	0.0	0.0	0.0
Wetlands – vegetive non-forest – peat extraction	CO ₂	9.2	0.0	0	18.9	18.9	0.0000	0.0	0.0	0.0	0.0	0.0
Settlements	CO ₂	75.7	122.2	0	63.4	63.4	0.0000	0.2	0.6	0.1	0.0	0.0
Other land	CO ₂	13.7	81.6	0	29.4	29.4	0.0000	0.3	0.4	0.1	0.0	0.0
Harvested wood products	CO ₂	-2,481.2	-8,658.8	0	68.2	68.2	0.0784	30.0	42.9	20.5	0.0	4.2
Direct N ₂ O emissions from N mineralisation/immobilisation	N ₂ O	181.4	82.1	8	86.5	86.9	0.0000	0.5	0.4	0.5	0.1	0.0
Direct N ₂ O emissions from drainage and rewetting	N ₂ O	77.7	124.6	33	80.0	86.5	0.0000	0.2	0.6	0.2	0.1	0.0
Indirect N ₂ O emissions from leaching and runoff	N ₂ O	40.8	18.5	8	127.6	127.8	0.0000	0.1	0.1	0.2	0.0	0.0

IPCC category	Gas	1990 emissions or removals (kt CO ₂ -e)	2021 emissions or removals (kt CO ₂ -e)	Activity data uncertainty (%)	Emission factor / estimation parameter uncertainty (biomass) (%)	Combined uncertainty (%)	Contribution to variance by category in 2021	Type A sensitivity (%)	Type B sensitivity (%)	Uncertainty in trend in LULUCF emissions introduced by emission factor/ estimation parameter uncertainty (%)	Uncertainty in trend in LULUCF emissions introduced by activity data uncertainty (%)	Uncertainty introduced into the trend in total LULUCF emissions (%)
N ₂ O emissions from biomass burning	N ₂ O	25.6	22.2	30	41.6	51.3	0.0000	0.0	0.1	0.0	0.0	0.0
CH ₄ emissions from biomass burning	CH ₄	68.4	32.6	30	41.6	51.3	0.0000	0.2	0.2	0.1	0.1	0.0
Total		-20,171.2	-21,078.2				Total uncertainty (%)	67.7			Total uncertainty in trend (%)	86.7

A3.2.9 LUCAS data management system

The LUCAS data management system stores, manages and archives data for international greenhouse gas reporting for the LULUCF sector. This system is used for managing the land use spatial databases, plot and reference data, and for combining the two sets of data to calculate the numbers required for reporting under the Convention and the Kyoto Protocol (see figure A3.2.17).

The data collected are stored and manipulated within three systems: the Geospatial System, the Gateway and the CRA.


The main objectives of these systems are to:

- provide a transparent system for data storage and carbon calculations
- provide a repository for the versioning and validation of plot measurements and land use data
- calculate carbon stocks, emissions and removals per hectare for land uses and carbon pools based on the plot and spatial data collected
- calculate biomass burning emissions by land use based on area and emission factors stored in the Gateway
- produce the outputs required for the LULUCF sector reporting under the UNFCCC and the Kyoto Protocol
- archive all inputs and outputs used in reporting.

The module 'joint calculations' refers to the process New Zealand uses to estimate national average carbon values by carbon pool for each land use category and subcategory.

The joint calculation process is performed within the CRA. Within the joint calculations interface, the user selects the appropriate area data and emission factors. The results of the calculations are carbon gains, losses and net change for all land use subcategories (whether in a conversion state or land remaining land), by year and by carbon pool.

Figure A3.2.17 New Zealand's LUCAS data management system




Note: IPCC = Intergovernmental Panel on Climate Change; LULUCF = Land Use, Land-Use Change and Forestry; LUM = land use map; QA/QC = quality assurance/quality control. Joint calculations are described below.

Geospatial System

The Geospatial System consists of applications and virtualised infrastructure customised to meet LULUCF reporting requirements. The infrastructure comprises a variety of file/object storage, relational databases, and servers supporting spatial data storage, management, versioning and running of web services for mapping. The core components of the Geospatial System are outlined in figure A3.2.18.

Figure A3.2.18 New Zealand's Geospatial System components



Land use mapping functionality

The LUCAS NZ Land Use Map (LUM) is made up of several spatial data sets including the Chatham Islands and the other offshore islands within New Zealand's Exclusive Economic Zone. For accuracy, these islands are mapped and stored in their own separate NZGD2000 map projections but are combined statistically for reporting purposes.

The LUM is designed to represent land use and confirmed land-use change at each of the mapping dates of 1990, 2008, 2012 and 2016. It is also necessary to track unconfirmed change that is detected in parallel maps (e.g., forest destocking) until the land use intention is confirmed and the change can be integrated into the LUM.

A variety of coded scripts (functioning as tools) is used to harness the Esri ArcGIS suite to manage and validate the integrity of these data sets. As the version of the ArcGIS suite is upgraded, the scripts are recompiled as required. Where coded scripts are not present, carefully documented manual processes are used to structure and manage the data sets.

Manual editing of the maps is typically carried out via branch-versioned Esri REST web feature services with coded calculation and constraint rules active. These rules ensure that attribution is consistent and complete, which includes ensuring that the sequences of land-use change mapped are valid.

The system allows multiple editors to have concurrent access from anywhere with internet, providing resilience during map production and the ability to scale mapping teams as required.

To accommodate scripted edits, these maps are exported from their respective feature services to Esri file geodatabase(s) with rules disabled, are processed, and then are scanned for topology and attribute validity before being re-imported into child versions within the services.


Production spatial data sets are stored in Esri ArcGIS Enterprise geodatabases. Some databases are used for reference and others for supporting web services.

The archive of remote sensing data, from which the LUCAS land-use change detection is derived, is structured and maintained in cloud-based object storage. Its metadata are managed centrally in ArcGIS Enterprise geodatabases.

LUCAS Management Studio

The LUCAS Management Studio (see figure A3.2.19) is the package of applications used to store activity data and calculate and report New Zealand's emissions and removals for LULUCF. The LUCAS Gateway is a data warehouse with the purpose of storing, versioning and validating activity data and emission factors. The CRA sources all data from the Gateway. It then calculates and outputs New Zealand's emissions and removals for LULUCF for land remaining land and land converted to another land use by pool and year.

Figure A3.2.19 LUCAS Management Studio



LUCAS Gateway


The LUCAS Gateway enables the storage of activity data such as field plot data, land use area, biomass burning and other data needed by the CRA, such as IPCC defaults.

The LUCAS Gateway provides a viewing, querying and editing interface to the source (plot, land use area, carbon and non-carbon) data. It also stores any published or saved results from running the CRA.

All activity data and emission factors are stored within the Gateway database (see figure A3.2.20). It contains the following main components.

- A data and results layer containing all activity data (natural, planted forest, soils, default carbon, non-carbon, land use areas, land-use change and reference tables). The user has the ability to create a 'snapshot' in time (a data set archiving system) of the data held in the Gateway. This enables users of the CRA to select from a range of data snapshots and ensures past results can be replicated over time.
- A validation layer allows users to judge the suitability of data for use in the CRA calculations, subsequent to passing primary validation. Where records are deemed not acceptable for use within published reports, they are tagged as 'invalid' in the LUCAS Gateway database.
- An audit trail provides a history of any changes to the database tables within the Gateway.
- Versioning at a number of levels ensures any changes to data, schema or the database itself are logged and versioned, while providing the user with the ability to track what changes have been applied and roll back to a previous version if required. The results of saved or published reports within the CRA are also stored within the Gateway for repeatability and reference.
- Primary data validation, both during data capture and during import of the data into the Gateway, ensures only data that have passed acceptability criteria are available for a publishable CRA run.
- Hosting and application support provides hosting services, system security, backup and restore, daily maintenance and monitoring for the Gateway and CRA.

Figure A3.2.20 LUCAS Gateway database



Calculation and Reporting Application

The CRA enables users to import carbon and non-carbon data from the Gateway and, by running the various modules, determine emissions and removals by New Zealand's *Forest land, Cropland, Grassland* and other land use types. This information, combined with land area data, enables New Zealand to meet its reporting requirements under the Convention and the Kyoto Protocol.

The CRA allows for the inclusion of other data sets, models and calculations without the complete redesign of the applications. All models, data and results are versioned, and the CRA allows the user to alter specific key values within a model or calculation (parameters) without the intervention of a programmer or technical support officer. The CRA is deployed as a client-based application that sources the required data from the Gateway.

The CRA comprises four modules: natural forest, soils, non-carbon and joint calculations. Any of these modules can be run independently or as a group. The results are provided as 'views' to the user at the completion of the run.

To activate a module, the user selects the module to run within the CRA, the version of the data set to be used, the model version and other calculation parameters. The natural forest and soil carbon modules use R statistical language as the base program language, while the non-carbon module and joint calculations module are developed in the programming language C Sharp (C#).

Within the joint calculations module, the user has the option of using the carbon results from running the modules or using default carbon estimates (based on published reports) stored within the Gateway. The joint calculations module combines the carbon estimates with the land use area to calculate carbon stock and change following the methodology set out in section 2.3 of volume 4 of the 2006 IPCC Guidelines (IPCC, 2006a). The results represent carbon stock and change for every 'from' and 'to' land use combination outlined by the IPCC since 1990.

On completion of running a module, the results can be saved or published back to the Gateway. This provides a versioned and auditable record of the results used for reporting. If the results are saved or published, other information is also saved for tracking and audit control, such as the time created, the user’s identification and the module-particular parameters that were used.


The CRA is maintained and supported by Interpine Innovation, a New Zealand-based company that specialises in forestry inventories and related information technology development. Interpine Innovation also provides support services, such as database and application backups, day-to-day issue resolution and enhancement projects to the Gateway or CRA as required.

Any changes to the data or table structure within the Gateway, or to the people accessing the Gateway or CRA, are tracked via audit logs. For any changes to the data within the Gateway, the person making the change, the date, the reason for change and the version are logged and reports are made available to users for review.

Quality control management for implementing planned improvements

In 2020, further quality control processes were introduced and formalised for introducing improvements to the National Inventory Report. This was done to help manage the large number of improvements to the LULUCF sector that were made for the 2021 submission and to improve the quality control procedures for implementation of future improvements. The quality control process is described in figure A3.2.21.

Figure A3.2.21 Improvements implementation quality control procedure



Document management

All reference material, including scientific reports containing information on methodologies or emission factors used in the production of the LULUCF and Kyoto Protocol estimates, is archived on the Ministry for the Environment’s document management store, Te Puna.

The emission factors and area estimates for published runs are also archived within the Gateway and can be accessed via the Gateway or the CRA. Information is not directly accessible by the public but can be supplied upon request.

Annex 3: References

Some references may be downloaded directly from the following webpage: www.mpi.govt.nz/news-and-resources/statistics-and-forecasting/greenhouse-gas-reporting/agriculture-greenhouse-gas-inventory-reports.

The Ministry for Primary Industries is progressively making reports used for the inventory available on this page, provided copyright permits.

Ausseil A, Jamali H, Clarkson B, Golubiewski N. 2015. Soil carbon stocks in wetlands of New Zealand and impact of land conversion since European settlement. *Wetlands Ecology and Management* 23(5): 947–961.

Baisden WT, Beets PN, Davis M, Wilde RH, Arnold G, Trotter CM. Unpublished(a). Changes in New Zealand's Soil Carbon Stocks Following Afforestation of Pastures. Contract report LC0506/105 prepared for the Ministry for the Environment by Landcare Research in 2006.

Baisden WT, Wilde RH, Arnold G, Trotter CM. Unpublished(b). Operating the New Zealand Soil Carbon Monitoring System. Contract report LC0506/107 prepared for the Ministry for the Environment by Landcare Research in 2006.

Basher L, Burrows L, Hough S, Lambie S, Ross C, Thornburrow D, Garrett L, Paul T, Evanson T, Oliver G, Osorio R, Pearce S, Hill R. Unpublished. Soil Data Collection and Analysis of Post-1989 Planted Forest Plot Network. Contract report prepared for the Ministry for the Environment by Landcare Research, New Zealand Forest Research Institute Limited (trading as Scion) and LandSystems.

Beets PN, Brandon AM, Goulding CJ, Kimberley MO, Paul TSH, Searles NB. 2011a. The inventory of carbon stock in New Zealand's post-1989 planted forest. *Forest Ecology and Management* 262: 1119–1130.

Beets PN, Brandon AM, Goulding CJ, Kimberley MO, Paul TSH, Searles N. 2012a. The national inventory of carbon stock in New Zealand's pre-1990 planted forest using a LiDAR incomplete-transect approach. *Forest Ecology and Management* 280: 187–197.

Beets PN, Garrett, LG. 2018. Carbon fraction of *Pinus radiata* biomass components within New Zealand. *New Zealand Journal of Forestry Science* 48: 14.

Beets PN, Holt L. Unpublished. Audit of the LUCAS Post-1989 Natural Forest Plot Inventory 2012–2013. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion) in 2014.

Beets PN, Kimberley MO. Unpublished. Improvements Contained in the Forest Carbon Predictor Version 3. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion).

Beets PN, Kimberley MO, McKinley RB. 2007. Predicting wood density of *Pinus radiata* annual growth increments. *New Zealand Journal of Forestry Science* 37: 241–266.

Beets PN, Kimberley MO, Oliver GR, Pearce SH. 2018a. Predicting wood density of growth increments of Douglas-fir stands in New Zealand. *New Zealand Journal of Forestry Science* 48: 8.

Beets PN, Kimberley MO, Oliver GR, Pearce SH, Graham JD. 2014a. The application of stem analysis methods to estimate carbon sequestration in arboreal shrubs from a single measurement of field plots. *Forests* 5: 919–935.

Beets PN, Kimberley MO, Oliver GR, Pearce SH, Graham JD. Unpublished. Post-1989 Natural Forest Carbon Stocks and Changes. Contract report 20093 prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion) in 2013.

Beets PN, Kimberley MO, Oliver GR, Pearce, SH, Graham JD, Brandon A. 2012b. Allometric equations for estimating carbon stocks in natural forest in New Zealand. *Forests* 3: 818–839.

- Beets PN, Kimberley MO, Oliver GR, Pearce SH, Graham JD, Henley D, Meason DF. 2018b. Plantation species-specific adjustment functions for the Forest Carbon Predictor in New Zealand. *New Zealand Journal of Forestry Science* 48: 20.
- Beets PN, Kimberley MO, Paul TSH, Garrett LG. 2011b. Planted Forest Carbon Monitoring System – Forest carbon model validation study for *Pinus radiata*. *New Zealand Journal of Forestry Science* 41: 177–189.
- Beets PN, Kimberley MO, Paul TSH, Oliver GR, Pearce SH, Buswell JM. 2014b. The inventory of carbon stocks in New Zealand’s post-1989 natural forest for reporting under the Kyoto Protocol. *Forests* 5(9): 2230–2252.
- Beets PN, Payton IJ. Unpublished. CMS Indigenous Forest and Shrubland QA/QC in Implementation Year 2002. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd and Landcare Research New Zealand Ltd.
- Beets PN, Robertson KA, Ford-Robertson JB, Gordon J, Maclaren JP. 1999. Description and validation of C_{change}: A model for simulating carbon content in managed *Pinus radiata* stands. *New Zealand Journal of Forestry Science* 29(3): 409–427.
- Brack C. Unpublished. Planted Forest Inventory Design: Post-1989 exotic. Contract report prepared for the Ministry for the Environment.
- Brack C, Broadley J. Unpublished. Co-location of LiDAR and Post-1989 Forest Plots. Contract report prepared for the Ministry for the Environment.
- Bretz F, Hothorn T, Westfall P. 2010. *Multiple Comparisons Using R*. Boca Raton, FL: CRC Press.
- Burnham KP, Anderson DR. 2002. *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach*. 2nd edn. New York: Springer.
- Carran RA, Theobald PW, Evans JP. 1995. Emissions of nitrous oxide from some grazed pasture soils. *New Zealand and Australian Journal of Soil Research* 33: 341–352.
- Carswell FE, Burrows LE, Hall GMJ, Mason NWH, Allen RB. 2012. Carbon and plant diversity gain during 200 years of woody succession in lowland. *New Zealand Journal of Ecology* 36(2): 191–202.
- Coomes DA, Allen RB, Scott NA, Goulding CJ, Beets PN. 2002. Designing systems to monitor carbon stocks in forests and shrublands. *Forest Ecology and Management* 164: 89–108.
- Daly BK, Wilde RH. Unpublished. Contribution of Soil Carbon to New Zealand’s CO₂ Emissions: I Reclassification of New Zealand Soil Series to IPCC Categories. Contract report LC9697/096 prepared for the Ministry for the Environment by Landcare Research in 1997.
- Davis MR, Wakelin SJ. Unpublished. Perennial Cropland Biomass: Sampling Requirements. Contract report 11407 prepared for the Ministry for the Environment by New Zealand Forest Research Institute Limited (trading as Scion).
- de Klein CAM, Barton L, Sherlock RR, Li Z, Littlejohn RP. 2003. Estimating a nitrous oxide emission factor for animal urine from some New Zealand pastoral soils. *Australian Journal of Soil Research* 41: 381–399.
- Doherty JJ, Anderson SA, Pearce HG. Unpublished. An Analysis of Wildfire Records in New Zealand: 1991–2007. Contract report 12796 prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion).
- Draper NR, Smith H. 1998. *Applied Regression Analysis*. New York: Wiley.
- Dresser M, Hewitt A, Willoughby J, Bellis S. 2011. *Area of Organic Soils*. Report prepared for the Ministry of Agriculture and Forestry by Landcare Research. Wellington: Ministry of Agriculture and Forestry.
- Dymond JR, Shepherd JD. 2004. The spatial distribution of indigenous forest and its composition in the Wellington region, New Zealand, from ETM+ satellite imagery. *Remote Sensing of Environment* 90: 116–125.
- Dymond JR, Shepherd JD, Qi J. 2001. A simple physical model of vegetation reflectance for standardising optical satellite imagery. *Remote Sensing of Environment* 37: 230–239.

- Easdale TA, Richardson SJ, Marden M, England JR, Gayoso-Aguilar J, Guerra-Carcamo JE, Brandon AM. 2019. Root biomass allocation in southern temperate forests. *Forest Ecology and Management* 453: 117542.
- Easdale TA, Burge OR, Wiser S, Richardson SJ. Unpublished. Carbon Assessment of Wetland Vegetation. Contract report LC4173 prepared for the Ministry for the Environment by Landcare Research New Zealand Ltd in 2022.
- Elsgaard L, Görresa CM, Hoffmann CC, Blicher-Mathiesen G, Schelde K, Petersen SO. 2012. Net ecosystem exchange of CO₂ and carbon balance for eight temperate organic soils under agricultural management. *Agriculture, Ecosystems & Environment* 162: 52–67.
- Eyles GO. 1977. NZLRI worksheets and their applications to rural planning. *Town Planning Quarterly* 47: 38–44.
- Fick J, Saggat S, Hill J, Giltrap D. 2011. *Poultry Management in New Zealand: Production, Manure Management and Emissions Estimations for the Commercial Chicken, Turkey, Duck and Layer Industries within New Zealand*. Report prepared for the Ministry of Agriculture and Forestry by Poultry Industry Association, Egg Producers Association, Landcare Research and Massey University. Wellington: Ministry of Agriculture and Forestry.
- Forest Industry Training and Education Council. 2005. *Best Practice Guidelines for Land Preparation*. Revised edn. Rotorua: Forest Industry Training and Education Council.
- Fraser S, Wilde H, Payton I, Scott J. Unpublished. Historic Soils Dataset – Land Use Reclassification. Contract report LC0809/131 prepared for the Ministry for the Environment by Landcare Research New Zealand Ltd.
- Garrett LG. Unpublished. Natural Forests Soils – Data Checking and Carbon Content of the Mineral Soil. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion).
- Garrett LG, Kimberley MO, Oliver GR, Parks M, Pearce SH, Beets PN, Paul TSH. 2019. Decay rates of above- and below-ground coarse woody debris of common tree species in New Zealand's natural forest. *Forest Ecology and Management* 438: 96–102.
- Garrett LG, Kimberley MO, Oliver GR, Pearce SH, Paul TSH. 2010. Decomposition of woody debris in managed *Pinus radiata*. *Forest Ecology and Management* 260: 1389–1398.
- Giltrap DJ, Betts H, Wilde RH, Oliver G, Tate KR, Baisden WT. Unpublished. Contribution of Soil Carbon to New Zealand's CO₂ Emissions. XIII: Integrate general linear model and digital elevation model. Landcare Research, Forest Research Joint Contract Report: JNT 9899/136.
- Hedley CB, Payton IJ, Lynn IH, Carrick ST, Webb TH, McNeill S. 2012. Random sampling of stony and non-stony soils for testing a national soil carbon monitoring system. *Soil Research* 50(1): 18–29.
- Herries D, Paul TSH, Beets PN, Chikono C, Thompson R, Searles N. Unpublished. Land Use and Carbon Analysis System: Planted Forest Data Collection Manual Version 6.4. Wellington: Ministry for the Environment released in 2019.
- Hewitt A. 2010. *New Zealand Soil Classification*. 3rd edn. Landcare Research Science Series No. 1. Lincoln: Manaaki Whenua Press.
- Hewitt A, Forrester G, Fraser S, Hedley C, Lynn I, Payton I. 2012. Afforestation effects on soil carbon stocks of low productivity grassland in New Zealand. *Soil Use and Management* 28(4): 508–516.
- Hill J. 2012. *Recalculate Pork Industry Emissions Inventory*. Report prepared for the Ministry of Agriculture and Forestry by Massey University and the New Zealand Pork Industry Board. Wellington: Ministry of Agriculture and Forestry.
- Holdaway RJ, McNeill SJ, Mason NW, Carswell F. 2014. Propagating uncertainty in plot-based estimates of forest carbon stock and carbon stock change. *Ecosystems* 17(4): 627–640.
- Holdaway RJ, Easdale TA, Carswell FE, Richardson SJ, Peltzer DA, Maon NW, Brandon AM, Coomes DA. 2017. Nationally representative plot network reveals contrasting drivers of net biomass change in secondary and old-growth forests. *Ecosystems* 20: 944–959.

- Hunter G, McNeill S. Unpublished. Review of LUCAS Land use Backcasting 1962–1989. Contract report LC70 prepared for the Ministry for the Environment by Landcare Research New Zealand Ltd.
- Indufor Asia Pacific. 2018. *New Zealand Deforestation Mapping 2015 and 2016 – Final Report*. Contract report prepared for the Ministry for the Environment. Wellington: Ministry for the Environment.
- Interpine Forestry Limited. Unpublished. LUCAS Post-1989 Planted Forest Soil Survey. Report prepared for the Ministry for the Environment in 2014.
- IPCC. 1996. Houghton JT, Meira Filho LG, Lim B, Treanton K, Mamaty I, Bonduki Y, Griggs DJ, Callender BA (eds). *IPCC/OECD/IEA. Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories*. Bracknell: United Kingdom Meteorological Office.
- IPCC. 2000. Penman J, Kruger D, Galbally I, Hiraishi T, Nyenzi B, Emmanuel S, Buendia L, Hoppaus R, Martinsen T, Meijer J, Miwa K, Tanabe K (eds). *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. IPCC National Greenhouse Gas Inventories Programme. Japan: Published for the IPCC by the Institute for Global Environmental Strategies.
- IPCC. 2006a. Eggleston HS, Buendia L, Miwa K, Ngara T, Tanabe K (eds). *2006 IPCC Guidelines for National Greenhouse Gas Inventories. Volume 4. Agriculture, Forestry and Other Land Use*. IPCC National Greenhouse Gas Inventories Programme. Japan: Published for the IPCC by the Institute for Global Environmental Strategies.
- IPCC. 2006b. Eggleston HS, Buendia L, Miwa K, Ngara T, Tanabe K (eds). *General Guidance and Reporting. Volume 1*. Japan: Published for the IPCC by the Institute for Global Environmental Strategies.
- Joyce K. Unpublished. Ministry for the Environment LUCAS 1990–2008 Change Detection Quality Assessment Procedure, GNS Science Consultancy Report 2009/45.
- Kelliher FM, de Klein CAM. Unpublished. Review of New Zealand’s Fertiliser Nitrous Oxide Emission Factor (EF₁) Data. Report prepared for the Ministry for the Environment by Landcare Research and AgResearch in 2006.
- Kimberley MO, Beets PN, Paul TSH. 2019. Comparison of measured and modelled change in coarse woody debris carbon stocks in New Zealand’s natural forest. *Forest Ecology and Management* 434 (2019): 18–28.
- Kimberley MO, Dean MG. 2006. *A Validation of the 300 Index Growth Model*. Report prepared for the Plantation Management Cooperative.
- Kirschbaum M, Trotter C, Wakelin S, Baisden T, Curtin D, Dymond J, Ghani A, Jones HS, Deurer M, Arnold G, Beets PN, Davis MR, Hedley C, Peltzer D, Ross C, Schipper L, Sutherland A, Wang H, Beare M, Clothier B, Mason N, Ward M. Unpublished. Carbon Stocks and Changes in New Zealand’s Soils and Forests, and Implications of Post-2012 Accounting Options for Land-based Emissions Offsets and Mitigation Opportunities. Contract report prepared for the Ministry for the Environment.
- Knowles RL. 2005. Development of a productivity index for Douglas-fir. *New Zealand Journal of Forestry* 50(2): 19–22.
- Lassey K. 2011. *Methane Emissions and Nitrogen Excretion Rates for New Zealand Goats*. Report for the Ministry of Agriculture and Forestry, National Institute of Water and Atmospheric Research. Wellington: National Institute of Water and Atmospheric Research.
- Lawrence-Smith EJ, Tregurtha CS, Beare MH. 2010. *Land Management Index Data for use in New Zealand’s Soil Carbon Monitoring System*. Contract report prepared by Plant and Food Research for the Ministry for the Environment. Contract report number: SPTS 4612. Wellington: Ministry for the Environment.
- Leathwick J, Morgan F, Wilson G, Rutledge D, McLeod M, Johnston K. 2002. *Land Environments of New Zealand: A Technical Guide*. Wellington: Ministry for the Environment.
- Lilburne LR, Hewitt AE, Webb TW. 2012. Soil and informatics science combine to develop S-map: A new generation soil information system for New Zealand. *Geoderma* 170: 232–238.
- Lynker Analytics Consortium. 2020. Deforestation Mapping 2017 and 2018 Technical Report. Prepared for the Ministry for the Environment by Lynker Analytics Consortium. Wellington: Ministry for the Environment.

- Manderson A, Hoogendoorn C, Newsome P. 2018. *Grassland improvement mapping using Innovative Data Analysis (IDA) techniques*. Contract report prepared for the Ministry for the Environment by Landcare Research New Zealand Ltd. Lincoln: Manaaki Whenua Landcare Research New Zealand Ltd.
- Manley B. 2009. *2008 Deforestation Intentions Survey*. Contract report prepared for the Ministry of Agriculture and Forestry by New Zealand School of Forestry, University of Canterbury. Wellington: Ministry of Agriculture and Forestry.
- Manley B. 2019. *Deforestation Intentions Survey 2018*. Wellington: Ministry for Primary Industries.
- Manley B. 2022. *Afforestation and Deforestation Intentions Survey 2021*. Wellington: Ministry for Primary Industries.
- Marcus R, Peritz E, Gabriel KR. 1976. On closed testing procedures with special reference to ordered analysis of variance. *Biometrika* 63: 655–660.
- McNeill SJ. Unpublished(a). LC93 Soil CMS Model Recalibration and Uncertainty Analysis. Contract report prepared for the Ministry for the Environment by Landcare Research New Zealand Ltd.
- McNeill SJ. Unpublished(b). LC975 Respecification and Reclassification of the MfE Soil CMS model. Contract report prepared for the Ministry for the Environment by Landcare Research New Zealand Ltd.
- McNeill SJ, Barringer JRF. Unpublished. Respecification and Reclassification of the 2013 MfE Soil CMS Model. Prepared for the Ministry for the Environment by Landcare Research New Zealand Ltd.
- McNeill SJ, Barringer JRF, Forrester GJ. Unpublished. LC1650 Development, Refinement and Calibration of the MfE Soil CMS Model. Prepared for the Ministry for the Environment by Landcare Research New Zealand Ltd.
- McNeill SJ, Golubiewski NE, Barringer J. 2014. Development and calibration of a soil carbon inventory model for New Zealand. *Soil Research* 52: 789–804.
- Milne JDG, Clayden B, Singleton PL, Wilson AD. 1995. *Soil Description Handbook*. Lincoln: Manaaki Whenua Press.
- Ministry for Primary Industries. 2016. *Material Flow and End-use of Harvested Wood Products Produced from New Zealand Log Exports*. Wellington: Ministry for Primary Industries.
- Ministry for Primary Industries. 2021. *National Exotic Forest Description as at 1 April 2021*. Wellington: Ministry for Primary Industries. Retrieved from www.canopy.govt.nz/forestry-data-research/national-exotic-forest-description-2021 (15 October 2022).
- Ministry for Primary Industries. 2022. *Quarterly Production Roundwood Removals from NZ Forests at 30 June 2022*. Wellington: Ministry for Primary Industries. Retrieved from www.mpi.govt.nz/forestry/forest-industry-and-workforce/forestry-wood-processing-data/wood-processing-data/ (27 September 2022).
- Ministry for the Environment. 2012. *Land Use and Carbon Analysis System: Satellite imagery interpretation guide for land use classes*. 2nd edn. Wellington: Ministry for the Environment.
- Ministry for the Environment. Unpublished. Land Use Carbon Analysis System Natural Forest Data Collection Manual Version 2.9 (released in 2019). Wellington: Ministry for the Environment.
- Moore JR, Goulding CJ. Unpublished. Sampling Methods and Associated Levels of Precision for a National Carbon Inventory in Planted Forests. Contract report prepared for Ministry for the Environment.
- Muller C, Sherlock RR, Williams PH. 1995. Direct field measurements of nitrous oxide emissions from urine-affected and urine-unaffected pasture in Canterbury. In: *Proceedings of the Workshop on Fertilizer Requirements of Grazed Pasture and Field Crops: Macro and Micronutrients*. Currie LD, Loganathan P (eds). Occasional Report No. 8. Palmerston North: Massey University. pp 243–234.
- Newsome P, Shepherd J, Pairman D. 2013. *Establishing New Zealand's LUCAS Land Use and Land Use-Change and Forestry 2012 Map*. Contract report prepared for the Ministry for the Environment by Landcare Research New Zealand Ltd. Lincoln: Manaaki Whenua Landcare Research New Zealand Ltd.

- Newsome P, Shepherd J, Pairman D, Bellis S, Manderson A. 2018. Establishing New Zealand's LUCAS 2016 Land Use Map. Contract report prepared for the Ministry for the Environment by Landcare Research New Zealand Ltd. Lincoln: Manaaki Whenua Landcare Research New Zealand Ltd.
- Newsome PF, Wilde RH, Willoughby EJ. 2008. *Land Resource Information System Spatial Data Layers: Data Dictionary*. Palmerston North: Landcare Research New Zealand Ltd.
- Ogle SM, Breidt FJ, Easter M, Williams S, Paustian K. 2007. An empirically based approach for estimating uncertainty associated with modelling carbon sequestration in soils. *Ecological Modelling* 205: 453–463.
- Pacheco D, Waghorn G, Rollo M. Unpublished. Methodology for Splitting Nitrogen between Livestock Dung and Urine. Report prepared for the Ministry for Primary Industries by AgResearch in 2018.
- Paul TSH, Andersen C, Kimberley MO, Dash J, Beets PN. Unpublished(a). Carbon Stock Changes in New Zealand's Pre-1990 Planted Forests Based on a Periodic Ground Inventory. Contract report prepared for the Ministry for the Environment by the New Zealand Forest Research Institute Ltd (trading as Scion) in 2016.
- Paul TSH, Beets PN, Kimberley MO. Unpublished(b). Carbon stocks in New Zealand's Post-1989 Natural Forest – Analysis of the 2018/2019 forest inventory data. Contract report prepared for the Ministry for the Environment by the New Zealand Forest Research Institute Ltd (trading as Scion) in 2020.
- Paul TSH, Dowling LJ. Unpublished. Audit of the LUCAS Natural Forest Plot Inventory 2018/2019. Contract report prepared for the Ministry for the Environment by the New Zealand Forest Research Institute Ltd (trading as Scion) in 2019.
- Paul TSH, Kimberley MO, Beets PN. 2021. Natural Forests in New Zealand: A large terrestrial carbon pool in a national state of equilibrium. *Forest Ecosystems* 8(1): 1–21.
- Paul TSH, Kimberley MO, Beets PN. Unpublished(c). Post-1989 and Pre-1990 Planted Forest Carbon Yield Tables and Stock Changes. Contract report prepared for the Ministry for the Environment by the New Zealand Forest Research Institute Ltd (trading as Scion) in 2013.
- Paul TSH, Kimberley MO, Dodunski C. Unpublished(d). The 2019 NFI Plot Analysis Yield Tables and Carbon Stocks at Measurement. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion) in 2020.
- Paul TSH, Wakelin SJ. Unpublished. Yield Tables and Approach for Estimating Historic Carbon Stocks in Pre-1990 Planted Forests for Greenhouse Gas Inventory Reporting. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion) in 2020.
- Paul TSH, Wakelin SJ, Dodunski, C. Unpublished(e). The NFI 2016–2020 Analysis: Yield Tables and Carbon Stocks in Planted Forests in New Zealand Based on a Five-year Inventory Cycle. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion) in 2021.
- Paul TSH, Wakelin SJ, Kimberley MO, Beets PN. Unpublished(f). Stocked Plantation Area in Mapped Post-1989 and Pre-1990 Forests, and Associated Crop Yield Tables and CRA Simulation Input Data. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion) in 2014.
- Payton IJ, Beets PN, Wilde H, Beadel S. Unpublished. CMS: Progress Report on Fieldwork Contract (03/04-0226-L) for the Period to 15 February 2004. Contract report prepared for the Ministry for the Environment by Landcare Research New Zealand Limited and New Zealand Forest Research Ltd.
- Payton IJ, Newell CL, Beets PN. 2004. *New Zealand Carbon Monitoring System Indigenous Forest and Shrubland Data Collection Manual*. Christchurch: Caxton Press.
- Payton IJ, Pearce HG. 2009. Fire-induced changes to the vegetation of tall-tussock (*Chionochloa rigida*) grassland ecosystems. *Science for Conservation* 290. Wellington: Department of Conservation.
- Pickering A, Fick JM. 2015. *Detailed methodologies for agricultural greenhouse gas emission calculation*. Version 3. Wellington: Ministry for Primary Industries. (Refer to: mpi.govt.nz/news-and-resources/statistics-and-forecasting/greenhouse-gas-reporting/)

- Pickering A, Gibbs J, Wear S, Fick J, Tomlin H. 2020. Methodology for Calculation of New Zealand's Agricultural Greenhouse Gas Emissions. Version 7. Wellington: Ministry for Primary Industries. Retrieved from www.mpi.govt.nz/dmsdocument/13906-Detailed-methodologies-for-agricultural-greenhouse-gas-emission-calculation (7 January 2022).
- Poyry Management Consulting (NZ) Ltd. Unpublished. Accuracy Assessment of LUCAS 2012 Land Use Map. Contract report prepared for the Ministry for the Environment.
- PricewaterhouseCoopers. Unpublished. LUCAS Data Quality Framework. Contract report prepared for the Ministry for the Environment.
- Robertson KA. 1998. Loss of organic matter and carbon during slash burns in New Zealand exotic forests. *New Zealand Journal of Forestry Science* 28(2): 221–241.
- Saggar S, Giltrap DL, Davison R, Gibson R, de Klein C, Rollo M, Ettema P, Rys G. 2015. Estimating direct N₂O emissions from sheep, beef, and deer grazed pastures in New Zealand hill country: Accounting for the effect of land slope on the N₂O emission factors from urine and dung. *Agriculture Ecosystems & Environment* 205: 70–78.
- Scott NA, Tate KR, Giltrap DJ, Tattersall SC, Wilde RH, Newsome P, Davis MR. 2002. Monitoring land-use change effects on soil carbon in New Zealand: Quantifying baseline soil carbon stocks. *Environmental Pollution* 116: S167–186.
- Shepherd JD, Bunting P, Dymond JR. 2019. Operational large-scale segmentation of imagery based on iterative elimination. *Remote Sensing*. 11(6): 658
- Shepherd JD, Dymond JR. 2003. Correcting satellite imagery for the variance of reflectance and illumination with topography. *International Journal of Remote Sensing* 24: 3503–3514.
- Shepherd JD, Newsome P. Unpublished(a). Establishing New Zealand's Kyoto Land Use and Land-use change and Forestry 1990 Map. Contract report prepared for the Ministry for the Environment.
- Shepherd JD, Newsome P. Unpublished(b). Establishing New Zealand's Kyoto Land Use and Land-use change and Forestry 2008 Map. Contract report prepared for the Ministry for the Environment.
- Sherlock RR, Jewell P, Clough T. 2008. *Review of New Zealand Specific Frac_{GASM} and Frac_{GASF} Emissions Factors*. Report prepared for the Ministry of Agriculture and Forestry by Landcare Research and AgResearch. Wellington: Ministry of Agriculture and Forestry.
- Suttie J. 2012. *Report to the Deer Industry New Zealand: Estimation of Deer Population and Productivity Data 1990 to 2012*. Dunedin: Suttie Consulting Limited.
- Tate KR, Barton JP, Trustrum NA, Baisden WT, Saggar S, Wilde RH, Giltrap DJ, Scott NA. 2003a. Monitoring and modelling soil organic carbon stocks and flows in New Zealand. In: CA Scott-Smith (ed.) *Soil Organic Carbon and Agriculture: Developing Indicators for Policy Analysis*. Proceedings of an OECD Expert Meeting, Ottawa, ON. France: Agriculture and Agri-Food Canada and Organisation for Economic Co-operation and Development.
- Tate KR, Scott NA, Saggar S, Giltrap DJ, Baisden WT, Newsome PF, Trotter CM, Wilde RH. 2003b. Land-use change alters New Zealand's terrestrial carbon budget: Uncertainties associated with estimates of soil carbon change between 1990–2000. *Tellus, Series B: Chemical and Physical Meteorology* 55(2): 364–377.
- Tate KR, Wilde RH, Giltrap DJ, Baisden WT, Saggar S, Trustrum NA, Scott NA. 2004. Current approaches to soil carbon monitoring in New Zealand. In: *SuperSoil 2004: Proceedings of the 3rd Australian New Zealand Soils Conference*, 5–9 December 2004, Sydney, University of Sydney, Australia. Retrieved from www.regional.org.au/au/asssi/supersoil2004 (27 February 2018).
- Tate KR, Wilde RH, Giltrap DJ, Baisden WT, Saggar S, Trustrum NA, Scott NA, Barton JP. 2005. Soil organic carbon stocks and flows in New Zealand: System development, measurement and modelling. *Canadian Journal of Soil Science* 85(4): 481–489.
- Taylor NH, Pohlen IJ. 1962. *Soil Survey Method: A New Zealand Handbook for the Field Study of Soils*. Lower Hutt: New Zealand Soil Bureau.

- Thomas S, Hume E, Fraser T, Curtin, D. 2011. *Factors and Activity Data to Estimate Nitrous Oxide Emissions from Cropping Systems, and Stubble and Tussock Burning*. Report prepared for Ministry of Agriculture and Forestry by Plant and Food Research. Wellington: Ministry of Agriculture and Forestry.
- Thomas S, Wallace D, Beare M. 2014. *Pasture Renewal Activity Data and Factors for New Zealand*. Report prepared for the Ministry for Primary Industries by Plant and Food Research. Wellington: Ministry for Primary Industries.
- Thomas SM, Fraser T, Curtin D, Brown H, Lawrence E. 2008. *Review of Nitrous Oxide Emission Factors and Activity Data for Crops*. Report prepared for the Ministry of Agriculture and Forestry by Plant and Food Research. Wellington: Ministry of Agriculture and Forestry.
- Thompson S, Gruner I, Gapare N. 2004. *New Zealand Land Cover Database Version 2: Illustrated guide to target classes*. Report prepared for the Ministry for the Environment. Wellington: Ministry for the Environment.
- Trotter C, MacKay A. Unpublished. Potential Forest Land. Landcare Research New Zealand Ltd Contract Report: 04/05-0410-L.
- UNFCCC. 2020. FCCC/ARR/2019/NZL. Report of the individual review of the annual submission of New Zealand submitted in 2019. In-country Review.
- van der Weerden T, Cox N, Luo J, Di HJ, Podolyan A, Phillips RL, Saggarr S, de Klein CAM, Ettema P, Rys G. 2016. Refining the New Zealand nitrous oxide emission factor for urea fertiliser and farm dairy effluent. *Agriculture Ecosystems & Environment* 222: 133–137.
- van der Weerden T, Noble A, Giltrap D, Luo S, Saggarr S. 2019. *Meta-analysis of nitrous oxide emission factors for excreta deposited onto pasture: final report*. Report prepared for the Ministry for Primary Industries by AgResearch.
- Wakelin SJ. Unpublished(a). Scientific research underpinning New Zealand’s carbon inventory in planted forests. Objective 9 – UNFCCC planted forest carbon inventory methodology. Milestone 2 – Review of shrubland clearance assumptions in the national carbon inventory. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion) in 2004.
- Wakelin SJ. Unpublished(b). Review of LULUCF Biomass Burning Assumptions in New Zealand’s Greenhouse Gas Inventory. Contract report prepared for the Ministry for the Environment by Ensis in 2006.
- Wakelin SJ. Unpublished(c). Carbon Inventory of New Zealand’s Planted Forests – Calculations revised in October 2008 for New Zealand’s 2007 Greenhouse Gas Inventory. Contract report prepared for the Ministry of Agriculture and Forestry by New Zealand Forest Research Institute Ltd (trading as Scion) in 2008.
- Wakelin SJ. Unpublished(d). Apportioning Wildfire Emissions to Forest Sub-categories in the National Greenhouse Gas Inventory. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion) in 2011.
- Wakelin SJ. Unpublished(e). Controlled Biomass Burning Emissions for the 2011 Greenhouse Gas Inventory. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion) in 2012.
- Wakelin SJ. Unpublished(f). Harvested Wood Products Model Revision. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion) in 2017.
- Wakelin SJ. Unpublished(g). Biomass Burning Activity Data for New Zealand’s 1990–2017 Greenhouse Gas Inventory. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion) in 2018.
- Wakelin SJ. Unpublished(h). Review of Emissions from Prescribed Burning of Standing Vegetation in New Zealand. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion) in 2018.

Wakelin SJ, Beets PN. Unpublished. Emission Factors for managed and unmanaged Grassland with Woody Biomass. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion) in 2013.

Wakelin SJ, Kimberley M. Unpublished. Harvested wood products model for UNFCCC and Kyoto Protocol reporting. Contract report prepared for the Ministry for the Environment by New Zealand Forest Research Institute Ltd (trading as Scion) in 2017.

Wakelin SJ, Searles N, Lawrence D, Paul TSH. 2020. Estimating New Zealand's harvested wood products carbon stocks and stock changes. *Carbon Balance and Management* 15(10).

Welten B, Mercer G, Smith C, Sprosen M, Ledgard S. 2021. *Refining estimates of nitrogen leaching for the New Zealand agricultural greenhouse gas inventory*. Report prepared for the Ministry for Primary Industries by AgResearch.

Wilde HR. 2003. *Manual for National Soils Database*. Palmerston North: Landcare Research New Zealand Ltd.

Wilde HR, Davis M, Tate K, Giltrap DJ. 2004. Testing the representativeness of soil carbon data held in databases underpinning the New Zealand Soil Carbon Monitoring System. *SuperSoil 2004: 3rd Australian New Zealand Soils Conference*, 5–9 December 2004. University of Sydney, Australia.

Wiser S. 2016. Vegetation Classification of All Measurements of the LUCAS Natural Forest Plots. Contract report prepared for the Ministry for the Environment by Manaaki Whenua Landcare Research.

Woollens R. Unpublished. Commentary on Analysis of the 2007–2008 Planted Forest Carbon Monitoring System Inventory Data of Post-1989 Forests. Report 11448 prepared for Ministry for the Environment.

Annex 4: Methodology and data collection for estimating emissions from fossil fuel combustion

New Zealand emission factors are based on gross calorific value. Energy activity data and emission factors in New Zealand are conventionally reported in gross (higher heating value) terms, with some minor exceptions. The convention adopted by New Zealand to convert gross calorific value to net calorific value follows the Organisation for Economic Co-operation and Development and International Energy Agency assumptions:

$$\text{Net calorific value} = 0.95 \times \text{gross calorific value for coal and liquid fuels}$$

$$\text{Net calorific value} = 0.90 \times \text{gross calorific value for gas}$$

$$\text{Net calorific value} = 0.80 \times \text{gross calorific value for wood}$$

Emission factors for gas, coal, biomass and liquid fuels used by New Zealand are shown in tables A4.1–A4.4. Where Intergovernmental Panel on Climate Change (IPCC) default emission factors are used, a net-to-gross factor as above is used to account for New Zealand activity data representing gross energy figures:

$$\text{Gross EF} = \text{Net EF} \times \text{Factor}$$

Table A4.1 Gross carbon dioxide emission factors used for New Zealand's energy sector in 2021

	Emission factor (t CO ₂ /TJ)	Source
Gas		
Weighted average	53.61	
Liquid fuels		
Crude oil	69.67	1
Regular petrol	66.77	2
Petrol – premium	66.95	2
Diesel (10 parts (sulphur) per million)	69.45	2
Jet kerosene	68.33	2
Av gas	65.89	2
LPG	59.27	3
Heavy fuel oil	73.33	2
Light fuel oil	73.02	2
Bitumen (asphalt)	76.43	2
Biomass		
Biogas	49.17	1
Wood (industrial)	89.47	1
Bioethanol	64.20	4
Biodiesel	67.26	3
Wood (residential)	85.80	3
Coal		
All sectors excl. electricity (sub-bituminous)	95.47	4
All sectors (bituminous)	87.78	4
All sectors (lignite)	76.12	4

1. IPCC Guidelines (2006).

2. Refining NZ.

3. *New Zealand Energy Information Handbook* (Eng et al., 2008).

4. *Review of Default Emission Factors in Draft Stationary Energy and Industrial Processes Regulations: Coal* (CRL Energy, 2009).

Table A4.2 Consumption-weighted average emission factors used for New Zealand's sub-bituminous coal-fired electricity generation for 1990 to 2021

Year	Emission factor (t CO ₂ /TJ)
1990	91.20
1991	91.24
1992	91.29
1993	91.33
1994	91.38
1995	91.42
1996	91.47
1997	91.51
1998	91.56
1999	91.60
2000	91.64
2001	91.69
2002	91.73
2003	91.78
2004	91.82
2005	91.87
2006	91.91
2007	92.43
2008	92.31
2009	92.39
2010 onwards	92.20

Table A4.3 Methane emission factors used for New Zealand's energy sector for 1990 to 2021

	Emission factor (t CH ₄ /PJ)	Source
Natural gas		
Electricity industries	0.90	IPCC 2006 (table 2.2)
Commercial	4.50	IPCC 2006 (table 2.4)
Residential	4.50	IPCC 2006 (table 2.5)
Domestic transport (CNG)	82.80	IPCC 2006 (table 3.2.2)
Other stationary (mainly industrial)	0.90	IPCC 2006 (table 2.3)
Liquid fuels		
Stationary sources		
Electricity – residual oil	2.85	IPCC 2006 (table 2.2)
Industrial (including refining) – residual oil	2.85	IPCC 2006 (table 2.3)
Industrial – LPG	0.95	IPCC 2006 (table 2.3)
Commercial – residual oil	9.50	IPCC 2006 (table 2.4)
Commercial – distillate oil	9.50	IPCC 2006 (table 2.4)
Commercial – LPG	4.75	IPCC 2006 (table 2.4)
Residential – distillate oil	9.50	IPCC 2006 (table 2.5)
Residential – LPG	4.75	IPCC 2006 (table 2.5)
Agriculture – stationary	2.85	IPCC 2006 (table 2.5)
Mobile sources		
LPG	58.9	IPCC 2006 (table 3.2.2)
Petrol	28.05	IPCC 2006 (table 3.2.2)
Diesel	3.71	IPCC 2006 (table 3.2.2)
Navigation (fuel oil and diesel)	6.65	IPCC 2006 (table 3.5.3)
Aviation fuel/kerosene	0.48	IPCC 2006 (table 3.6.5)

	Emission factor (t CH ₄ /PJ)	Source
Coal		
Electricity generation	0.95	IPCC 2006 (table 2.2)
Industry	9.50	IPCC 2006 (table 2.3)
Commercial	9.50	IPCC 2006 (table 2.4)
Residential	285.00	IPCC 2006 (table 2.5)
Biomass		
Wood/wood waste	24.00	IPCC 2006 (table 2.3)
Wood – fireplaces	240.00	IPCC 2006 (table 2.5) wood – residential
Bioethanol	18.00	IPCC 2006 (table 3.2.2) – ethanol, cars, Brazil
Biodiesel	18.00	IPCC 2006 (table 3.2.2) – ethanol, cars, Brazil
Gas biomass	0.90	IPCC 2006 (table 2.2)

Table A4.4 Nitrous oxide emission factors used for New Zealand’s energy sector for 1990 to 2021

	Emission factor (t N ₂ O/PJ)	Source
Natural gas		
Electricity generation	0.09	IPCC 2006 (table 2.2)
Commercial	0.09	IPCC 2006 (table 2.4)
Residential	0.09	IPCC 2006 (table 2.5)
Domestic transport (CNG)	2.70	IPCC 2006 (table 3.2.2)
Other stationary (mainly industrial)	0.09	IPCC 2006 (table 2.3)
Liquid fuels		
Stationary sources		
Electricity – residual oil	0.57	IPCC 2006 (table 2.2)
Electricity – distillate oil	0.57	IPCC 2006 (table 2.2)
Industrial (including refining) – residual oil	0.57	IPCC 2006 (table 2.2)
Industrial – distillate oil	0.57	IPCC 2006 (table 2.3)
Commercial – residual oil	0.57	IPCC 2006 (table 2.4)
Commercial – distillate oil	0.57	IPCC 2006 (table 2.4)
Residential (all oil)	0.57	IPCC 2006 (table 2.5)
LPG (all uses)	0.095	IPCC 2006 (tables 2.2–2.5)
Agriculture – stationary	0.38	Tier 2, diesel engines – agriculture
Mobile sources		
LPG	0.19	IPCC 2006 (table 3.2.2)
Petrol	7.60	IPCC 2006 (table 3.2.2)
Diesel	3.71	IPCC 2006 (table 3.2.2)
Fuel oil (ships)	1.90	IPCC 2006 (table 3.5.3)
Aviation fuel/kerosene	1.90	IPCC 2006 (table 3.6.5)
Coal		
Electricity generation	1.43	IPCC 2006 (table 2.2)
Industry	1.43	IPCC 2006 (table 2.3)
Commercial	1.43	IPCC 2006 (table 2.4)
Residential	1.43	IPCC 2006 (table 2.5)
Biomass		
Wood (all uses)	3.20	IPCC 2006 (table 2.5) wood/wood waste
Gas biomass	0.09	IPCC 2006 (table 2.5)

A4.1 Emissions from liquid fuels

A4.1.1 Activity data and uncertainties

The *Delivery of Petroleum Fuels by Industry Survey* is conducted by the Ministry of Business, Innovation and Employment (MBIE). Because it is a census, it has no sampling error. The only possible sources of error are non-sampling errors (such as respondent error and processing error). The 2021 statistical difference for liquid fuels in the balance table of the publication *Energy in New Zealand* was 2.3 per cent (MBIE, 2022). This is used as the activity data uncertainty for liquid fuels in 2021.

A4.1.2 Emission factors and uncertainties

The carbon dioxide (CO₂) emission factors are described in table A4.1. A complete time series of gross calorific values is available online: www.mbie.govt.nz/building-and-energy/energy-and-natural-resources/energy-statistics-and-modelling/energy-statistics/oil-statistics/. Table A4.5 gives a complete time series of carbon content of liquid fuels. This information is supplied by Refining NZ and is used in the calculation of annual emission factors for liquid fuels.

A 2009 consultant report (Hale and Twomey, unpublished) to the Ministry for the Environment estimates the uncertainty of CO₂ emission factors for liquid fuels at ± 0.5 per cent. The uncertainty for methane (CH₄) and nitrous oxide (N₂O) emission factors is ± 50.0 per cent because almost all emission factors are IPCC defaults.

Table A4.6 provides emission factors for European gasoline and diesel vehicles from the COPERT IV model that are used to estimate non-CO₂ emissions from road transport.

Table A4.5 Carbon content (per cent mass) for liquid fuels for 1990 to 2021

	Premium petrol	Regular petrol	Diesel	Jet kerosene	Heavy fuel oil	Light fuel oil	Bitumen (asphalt)
1990	84.87	84.92	86.28	85.92	86.22	86.67	86.57
1991	85.04	85.04	86.33	85.89	86.26	86.30	86.57
1992	85.03	85.13	86.29	85.84	86.25	86.18	86.57
1993	85.25	85.13	86.32	85.94	86.27	86.20	86.56
1994	85.21	85.19	86.30	85.99	86.25	86.13	86.57
1995	85.30	85.13	86.63	86.05	86.25	86.39	86.57
1996	85.66	85.13	86.73	86.16	86.28	86.45	86.57
1997	85.63	85.04	86.64	86.04	86.35	86.55	86.58
1998	85.72	85.17	86.52	86.14	86.22	86.39	86.63
1999	85.65	85.15	86.69	86.10	86.20	86.53	86.63
2000	85.67	85.16	86.64	86.25	86.22	86.58	86.63
2001	85.65	85.09	86.53	86.18	86.21	86.49	86.64
2002	85.68	85.06	86.57	86.10	86.25	86.68	86.66
2003	85.76	85.19	86.58	86.23	86.23	86.76	86.63
2004	85.66	85.22	86.62	86.20	86.24	86.58	86.58
2005	85.58	85.22	86.62	86.12	86.18	86.52	86.57
2006	85.54	85.25	86.57	86.24	86.34	86.93	86.57
2007	85.54	85.23	86.61	86.24	86.30	86.87	86.57
2008	85.63	85.32	86.70	86.32	86.39	86.87	86.57
2009	85.56	85.38	86.72	86.36	86.37	86.83	86.60
2010	85.54	85.40	86.77	86.35	86.31	86.90	86.59
2011	85.55	85.37	86.78	86.32	86.37	86.87	86.64
2012	85.51	85.38	86.84	86.34	86.25	86.89	86.63

	Premium petrol	Regular petrol	Diesel	Jet kerosene	Heavy fuel oil	Light fuel oil	Bitumen (asphalt)
2013	85.49	85.35	86.73	86.22	86.24	86.68	86.65
2014	85.57	85.42	86.74	86.23	86.33	86.87	86.65
2015	85.54	85.40	86.81	86.33	86.30	86.90	86.62
2016	85.66	85.48	86.56	86.11	86.28	86.58	86.60
2017	85.68	85.46	86.60	86.15	86.30	86.89	86.63
2018	85.69	85.49	86.61	86.31	86.04	86.93	86.04
2019	85.66	85.53	86.65	86.19	85.97	86.96	86.04
2020	85.66	85.53	86.65	86.19	85.97	86.96	86.04
2021	85.66	85.53	86.65	86.19	85.97	86.96	86.04

Table A4.6 Emission factors for European gasoline and diesel vehicles – COPERT IV model (European Environment Agency, 2007)

Vehicle type and emission standard	N ₂ O emission factors (mg/km)				CH ₄ emission factors (mg/km)			
	Cold	Urban		Highway	Cold	Urban		Highway
		Hot	Rural			Hot	Rural	
Passenger car								
Gasoline								
pre-Euro	10.0	10.0	6.5	6.5	201.0	131.0	86.0	41.0
Euro 1	18.8	26.5	10.7	5.5	45.0	26.0	16.0	14.0
Euro 2	12.6	12.7	4.9	2.7	94.0	17.0	13.0	11.0
Euro 3	8.3	1.50	0.33	0.23	83.0	3.0	2.0	4.0
Euro 4	5.5	1.95	0.34	0.22	57.0	2.87	2.69	5.08
Euro 5	2.15	2.22	0.19	1.20	57.0	2.87	2.69	5.08
Euro 6	2.15	2.22	0.19	1.20	57.0	2.87	2.69	5.08
Diesel								
pre-Euro	0.0	0.0	0.0	0.0	22.0	28.0	12.0	8.0
Euro 1	0.0	2.0	4.0	4.0	18.0	11.0	9.0	3.0
Euro 2	3.0	4.0	6.0	6.0	6.0	7.0	3.0	2.0
Euro 3	15.0	9.0	4.0	4.0	3.0	3.0	0.0	0.0
Euro 4	15.0	9.0	4.0	4.0	1.1	1.1	0.0	0.0
Euro 5	15.0	9.0	4.0	4.0	1.1	1.1	0.0	0.0
Euro 6	9.0	9.0	4.0	4.0	1.1	1.1	0.0	0.0
LPG								
pre-Euro	0.0	0.0	0.0	0.0	80.0	80.0	35.0	25.0
Euro 1	38.0	21.0	13.0	8.0	80.0	80.0	35.0	25.0
Euro 2	23.0	13.0	3.0	2.0	80.0	80.0	35.0	25.0
Euro 3	9.0	5.0	2.0	1.0	80.0	80.0	35.0	25.0
Euro 4	9.0	5.0	2.0	1.0	80.0	80.0	35.0	25.0
Euro 5	1.8	2.1	0.2	1.0	80.0	80.0	35.0	25.0
Euro 6	1.8	2.1	0.2	1.0	80.0	80.0	35.0	25.0
Light duty vehicles								
Gasoline								
pre-Euro	10.0	10.0	6.5	6.5	201.0	131.0	86.0	41.0
Euro 1	47.3	46.3	27.5	13.8	45.0	26.0	16.0	14.0
Euro 2	83.8	27.7	15.8	12.3	94.0	17.0	13.0	11.0
Euro 3	17.1	8.5	1.5	1.5	83.0	3.0	2.0	4.0
Euro 4	14.1	1.17	0.36	0.36	57.0	2.0	2.0	0.0
Euro 5	2.10	2.22	0.19	1.20	57.0	2.0	2.0	0.0
Euro 6	2.10	2.22	0.19	1.20	57.0	2.0	2.0	0.0

Vehicle type and emission standard	N ₂ O emission factors (mg/km)				CH ₄ emission factors (mg/km)			
	Urban		Rural	Highway	Urban		Rural	Highway
	Cold	Hot			Cold	Hot		
Diesel								
pre-Euro	0.0	0.0	0.0	0.0	22.0	28.0	12.0	8.0
Euro 1	0.0	2.0	4.0	4.0	18.0	11.0	9.0	3.0
Euro 2	3.0	4.0	6.0	6.0	6.0	7.0	3.0	2.0
Euro 3	15.0	9.0	4.0	4.0	3.0	3.0	0.0	0.0
Euro 4	15.0	9.0	4.0	4.0	1.1	1.1	0.0	0.0
Euro 5	15.0	9.0	4.0	4.0	1.1	1.1	0.0	0.0
Euro 6	9.0	9.0	4.0	4.0	1.1	1.1	0.0	0.0
Heavy duty truck and bus								
Gasoline all technologies	6.0	6.0	6.0	6.0	140.0	140.0	110.0	70.0
Diesel								
		GVW≤12t				GVW≤12t		
pre-Euro	30.0	30.0	30.0	30.0	85.0	85.0	23.0	20.0
Euro I	6.0	6.0	5.0	3.0	85.0	85.0	23.0	20.0
Euro II	5.0	5.0	5.0	3.0	54.4	54.4	20.0	18.6
Euro III	3.0	3.0	3.0	2.0	47.6	47.6	21.4	18.2
Euro IV	6.0	6.0	7.2	5.8	2.6	2.6	1.6	1.2
Euro V	15.0	15.0	19.8	17.2	2.6	2.6	1.6	1.2
Euro VI	18.5	18.5	19.0	15.0	2.6	2.6	1.6	1.2
		12t<GVW≤16t				12t<GVW≤16t		
pre-Euro	30.0	30.0	30.0	30.0	85.0	85.0	23.0	20.0
Euro I	11.0	11.0	9.0	7.0	85.0	85.0	23.0	20.0
Euro II	11.0	11.0	9.0	6.0	54.4	54.4	20.0	18.6
Euro III	5.0	5.0	5.0	4.0	47.6	47.6	21.4	18.2
Euro IV	11.2	11.2	13.8	11.4	2.6	2.6	1.6	1.2
Euro V	29.8	29.8	40.2	33.6	2.6	2.6	1.6	1.2
Euro VI	37.0	37.0	39.0	29.0	2.6	2.6	1.6	1.2
		16t<GVW≤28t				16t<GVW≤28t		
pre-Euro	30.0	30.0	30.0	30.0	175.0	175.0	80.0	70.0
Euro I	11.0	11.0	9.0	7.0	175.0	175.0	80.0	70.0
Euro II	11.0	11.0	9.0	6.0	112.0	112.0	69.6	65.1
Euro III	5.0	5.0	5.0	4.0	98.0	98.0	74.4	63.7
Euro IV	11.2	11.2	13.8	11.4	5.3	5.3	5.6	4.2
Euro V	29.8	29.8	40.2	33.6	5.3	5.3	5.6	4.2
Euro VI	37.0	37.0	39.0	29.0	5.3	5.3	5.6	4.2
		28t<GVW≤34t				28t<GVW≤34t		
pre-Euro	30.0	30.0	30.0	30.0	175.0	175.0	80.0	70.0
Euro I	17.0	17.0	14.0	10.0	175.0	175.0	80.0	70.0
Euro II	17.0	17.0	14.0	10.0	112.0	112.0	69.6	65.1
Euro III	8.0	8.0	8.0	6.0	98.0	98.0	74.4	63.7
Euro IV	17.4	17.4	21.4	17.4	5.3	5.3	5.6	4.2
Euro V	45.6	45.6	61.6	51.6	5.3	5.3	5.6	4.2
Euro VI	56.5	56.5	59.5	44.5	5.3	5.3	5.6	4.2
		GVW>34t				GVW>34t		
pre-Euro	30.0	30.0	30.0	30.0	175.0	175.0	80.0	70.0
Euro I	18.0	18.0	15.0	11.0	175.0	175.0	80.0	70.0

Vehicle type and emission standard	N ₂ O emission factors (mg/km)				CH ₄ emission factors (mg/km)			
	Urban		Rural	Highway	Urban		Rural	Highway
	Cold	Hot			Cold	Hot		
Euro II	18.0	18.0	15.0	10.0	112.0	112.0	69.6	65.1
Euro III	9.0	9.0	9.0	7.0	98.0	98.0	74.4	63.7
Euro IV	19.0	19.0	23.4	19.2	5.3	5.3	5.6	4.2
Euro V	49.0	49.0	66.6	55.8	5.3	5.3	5.6	4.2
Euro VI	61.0	61.0	64.0	48.0	5.3	5.3	5.6	4.2
Urban bus or coach		All types				All types		
pre-Euro	30.0	30.0	30.0	30.0	175.0	175.0	80.0	70.0
Euro I	12.0	12.0	9.0	7.0	175.0	175.0	80.0	70.0
Euro II	12.0	12.0	9.0	6.0	113.8	113.8	52.0	45.5
Euro III	6.0	6.0	5.0	4.0	103.3	103.3	47.2	41.3
Euro IV	12.8	12.8	13.8	11.4	5.3	5.3	2.4	2.1
Euro V	33.2	33.2	40.2	33.6	5.3	5.3	2.4	2.1
Euro VI	41.5	41.5	39.0	29.0	5.3	5.3	2.4	2.1
CNG								
pre-Euro					6,800	6,800	6,800	6,800
Euro I					6,800	6,800	6,800	6,800
Euro II					4,500	4,500	4,500	4,500
Euro III					1,280	1,280	1,280	1,280
Euro IV and later					980	980	980	980
Power two wheeler								
Gasoline								
<50 cm ³	1.0	1.0	1.0	1.0	219	219	219	219
>50 cm ³ 2-stroke	2.0	2.0	2.0	2.0	150	150	150	150
>50 cm ³ 4-stroke	2.0	2.0	2.0	2.0	200	200	200	200

A4.2 Emissions from solid fuels

A4.2.1 Activity data and uncertainties

The *New Zealand Quarterly Statistical Return of Coal Production and Sales* conducted by MBIE has near coverage of the sector, meaning that sampling error is small. The only other possible sources of error are non-sample errors (such as respondent error and processing error). The 2021 statistical difference for solid fuels in the balance table of the publication *Energy in New Zealand* was 1.0 per cent (MBIE, 2022). This is used as the activity data uncertainty for solid fuels in 2021.

A4.2.2 Emission factors and uncertainties

The estimated uncertainty in CO₂ emission factors for solid fuels is ±2.2 per cent. This is based on the difference between the range of updated emission factors for the three different ranks of coal used in New Zealand. The uncertainty for CH₄ and N₂O emission factors is ±50.0 per cent because almost all emission factors are IPCC defaults.

A4.3 Emissions from gaseous fuels

A4.3.1 Activity data

Through the various surveys and information it collects, MBIE has full coverage of the natural gas sector. This means that there is no sampling error in natural gas statistics and the only possible sources of error include those such as respondent error and processing error. The 2021 statistical difference for gaseous fuels in the balance table of the publication *Energy in New Zealand* was 1.4 per cent (MBIE, 2022). This is used as the activity data uncertainty for gaseous fuels in 2021.

A4.3.2 Emission factors

The estimated uncertainty in CO₂ emission factors for gaseous fuels is ± 2.4 per cent. This is based on the difference between the range of emission factors for three large gas fields in New Zealand. Together, these gas fields contributed over half of New Zealand's total gas supply in 2020. The uncertainty for CH₄ and N₂O emission factors is ± 50.0 per cent because almost all emission factors are IPCC defaults.

A4.4 Energy balance

Detailed and up-to-date energy balance tables for New Zealand are available online: www.mbie.govt.nz/building-and-energy/energy-and-natural-resources/energy-statistics-and-modelling/energy-statistics/energy-balances.

Further information can be found within the publication *Energy in New Zealand* (MBIE, 2022), which is also available online: www.mbie.govt.nz/building-and-energy/energy-and-natural-resources/energy-statistics-and-modelling/energy-publications-and-technical-papers/energy-in-new-zealand.

Table A4.7 gives a time series of energy use versus non-energy use of natural gas.

Table A4.7 Split of energy use and non-energy use of natural gas in petajoules

	Energy use	Non-energy use
1990	129.5	14.2
1991	143.9	22.1
1992	152.6	18.8
1993	148.0	21.1
1994	137.7	25.8
1995	127.4	36.2
1996	147.7	47.5
1997	170.4	48.9
1998	146.2	46.6
1999	168.5	54.2
2000	173.9	61.8
2001	190.6	55.4
2002	177.1	57.8
2003	151.9	26.1
2004	129.8	32.1
2005	136.4	13.0

	Energy use	Non-energy use
2006	137.2	15.0
2007	148.6	15.4
2008	135.5	18.4
2009	132.5	25.5
2010	147.1	25.6
2011	133.5	24.5
2012	145.6	32.0
2013	148.2	40.3
2014	149.5	60.7
2015	141.4	51.4
2016	133.3	59.1
2017	145.9	53.8
2018	134.8	45.3
2019	140.8	51.3
2020	136.3	46.6
2021	117.4	38.4

A4.5 Carbon dioxide reference approach for the Energy sector

A4.5.1 Estimation of carbon dioxide using the IPCC reference approach

The reference approach uses a country's energy supply data to calculate the CO₂ emissions from the combustion of fossil fuels using the apparent consumption equation. The apparent consumption in the reference approach is derived from production, import and export data. This information is included as a check for combustion-related emissions calculated from the sectoral approach.

The apparent consumption for primary fuels in the reference approach is obtained from 'calculated' energy-use figures (see annex 2 and section A4.4). These are derived as a residual figure from an energy balance equation comprising production, imports, exports, stock change and international transport on the supply side, according to the IPCC Guidelines for National Greenhouse Gas Inventories (IPCC, 2006).

The majority of the CO₂ emission factors for the reference approach are specific to New Zealand. Most emission factors for liquid fuels are based on annual carbon content and the gross calorific value data provided by New Zealand's only oil refinery, The New Zealand Refining Company Ltd. Where these data are not available, an IPCC default is used. The natural gas emission factor is based on a production-derived, weighted average of emission factors from all gas production fields.

Solid fuels in iron and steel manufacture

As mentioned in chapter 3, section 3.2.3, some of the coal production activity data in the reference approach are used in steel production. The Industrial Processes and Product Use sector accounts for the CO₂ emissions from this coal in the sectoral approach, as recommended by the IPCC Guidelines (IPCC, 2006); therefore they are not included in the common reporting format table 1.AA *Fuel combustion* – sectoral approach.


For simplicity, all feedstock carbon is excluded from the reference approach according to the IPCC Guidelines (IPCC, 2006). Without taking into account the use of by-product gases, this can create some discrepancies between the reference and sectoral approaches.

A4.5.2 Comparison of the IPCC reference approach with the New Zealand sectoral methodology

For 2020, CO₂ emissions estimated with the sectoral approach were 6.7 per cent lower than those estimated with the reference approach. Figure A4.1 shows the results for the two approaches for the period 1990 to 2020.

In some years, differences exist between the reference and sectoral approaches. In large part, these differences are due to the statistical differences found in the energy balance tables (MBIE, 2022) that are used as the basis for the reference and sectoral approach. Since 2000, the standard of national energy data has improved significantly, due to increased resources and focus. Before 2008, various energy statistics were collected by Stats NZ or MBIE. In 2008, Stats NZ delegated responsibility for the collection and analysis of national energy data to MBIE. The change resulted in a more consistent and transparent approach to energy data collection because one agency collected data across the supply chain.

Figure A4.1 Reference and sectoral approaches to carbon dioxide by fuel type (kt CO₂)



Sources of differences

- For gaseous fuels, the field-specific emission factors are used for natural gas supplied for industrial processes, while the reference approach uses an average emission factor.
- For liquid fuels, the energy balance is mass balanced but not carbon balanced. The fuel category 'other oil' is an aggregation of several fuel types, and so it is difficult to quantify a reliable carbon emission factor for the reference approach.
- In the sectoral approach, sector- or even plant-specific calorific values are used to calculate energy consumption, whereas in the reference approach, average (country-specific) calorific values are applied.

Annex 4: References

CRL Energy Ltd. 2009. *Reviewing Default Emission Factors in Draft Stationary Energy and Industrial Processes Regulations: Coal*. Contract report prepared for the Ministry for the Environment. Wellington: Ministry for the Environment.

Eng G, Bywater I, Hendtlass C. 2008. *New Zealand Energy Information Handbook*. Christchurch: New Zealand Centre for Advanced Engineering.

European Environment Agency. 2007. *EMEP/CORINAIR Emission Inventory Guidebook – 2007*. Copenhagen: European Environment Agency.

Hale R, Twomey I. Unpublished. Reviewing default emission factors for liquid fossil fuels adopted by the New Zealand Emissions Trading Scheme. Report by consultants Hale & Twomey Limited prepared for the Ministry for the Environment in 2009.

IPCC. 2006. Eggleston HS, Buendia L, Miwa K, Ngara T, Tanabe K (eds). *2006 IPCC Guidelines for National Greenhouse Gas Inventories. Volume 2. Energy*. IPCC National Greenhouse Gas Inventories Programme. Japan: Institute for Global Environmental Strategies for IPCC.

MBIE. 2022. *Energy in New Zealand 2022*. Wellington: Ministry of Business, Innovation and Employment.

Annex 5: Supplementary information for the KP-LULUCF sector

A5.1 Technical corrections to the FMRL

A5.1.1 Introduction

This submission covers the period outside the 2013 to 2020 forest management reference level (FMRL), therefore no technical corrections have been made. New Zealand is yet to set the FMRL for 2021 onwards. Reporting for the second commitment period (CP2) of the Kyoto Protocol (2013-2020) was covered in the previous inventory submission.

Information on New Zealand's 2013 to 2020 FMRL

New Zealand's FMRL was initially set at 11.15 million tonnes carbon dioxide equivalent (Mt CO₂-e) on average per year for the period 2013 to 2020 (New Zealand Government, 2011). This value was constructed using a business-as-usual projection of pre-1990 planted forest growth and harvest for the period 2013 to 2020. It was based on yield tables and statistics on the area in each age class of pre-1990 planted forest from the National Exotic Forest Description (NEFD) as at 2009 (Ministry for Primary Industries, 2010).

The 2011 FMRL included the following assumptions:

- pre-1990 natural forests were in steady state
- no pre-1990 planted forest deforestation would occur between 2013 and 2020 (pre-1990 natural forests were excluded from the analyses; post-1989 forest deforestation is reported under Article 3.3 – *Deforestation*)
- between 2013 and 2020, 2,000 hectares per year would be converted to non-forest land, and the equivalent forest would be planted elsewhere (i.e., 2,000 hectares per year would be reported as carbon equivalent forest (CEF) and be accounted for under *Forest management*)
- while harvest of post-1989 planted forest will increase over the period, pre-1990 planted forests will still make up a substantial proportion of total forest harvest
- all carbon is instantly emitted at the time of harvest (emissions and removals by the *Harvested wood products* pool were not considered)
- no allowance was made for the impacts of potential natural disturbances beyond background levels captured in the carbon stock yield tables.

The FMRL also reflects the following New Zealand legislation (including amendments) and current policies:

- the Forest Act 1949, which regulates the removal of timber from natural indigenous forests
- the South Island Landless Natives Act 1906, which transferred 17,000 hectares of natural indigenous forest to South Island Māori. The harvesting of this forest is also subject to the Resource Management Act 1991
- the Climate Change Response Act 2002, which makes owners of pre-1990 planted forest who deforest liable for the emissions associated with that activity

- the New Zealand’s biofuels policy of the time (under which it was thought most feedstock for biofuel was likely to be derived from non-forest sources).

It was assumed that this legislation and these policies would prevent any significant deforestation of pre-1990 forests, and that the New Zealand Emissions Trading Scheme would encourage harvest in pre-1990 planted forests over post-1989 forest.

The 2011 FMRL was determined by modelling the pre-1990 planted forest estate using a Forestry-Oriented Linear Programming Interpreter (FOLPI). As mentioned above, the model developed in FOLPI was based on an age-class distribution of pre-1990 planted forest as at 2009 from the NEFD, and simulated expected harvesting and replanting of this forest. Some additional modelling of decay of residues from harvest events was also carried out in MS Excel.

Since the 2011 FMRL was submitted, supplementary guidance has been prepared that describes the circumstances that would trigger a technical correction to the FMRL (IPCC, 2014). Changes to policies that affect harvest rate (as listed above) cannot be corrected for, but corrections can be made to reflect changes to the method for reporting against the FMRL and to address recommendations made by United Nations Framework Convention on Climate Change (UNFCCC) expert review teams (ERTs).

A technical assessment of New Zealand’s reference level submission was carried out by an ERT in 2011 (UNFCCC, 2011). The ERT noted a number of items for New Zealand to address by either providing additional data or applying technical corrections. These items included (UNFCCC, 2011, pp 6–10):

- maintaining consistency in the fraction of harvested biomass instantly oxidised when estimating emissions from harvest in the FMRL and in reporting against it (paragraph 21)
- ensuring consistency between the National Inventory Report (NIR) and the FMRL and, therefore, updating the current FMRL when new data or information become available (paragraph 22)
- making efforts to disaggregate gains and losses by biomass pool (paragraph 35)
- providing further information on how forest owners will be able to move from historical and current harvesting practice to the longer rotation length projected in the FOLPI model (paragraph 36)
- explaining in more detail how the difference in both harvested areas and harvesting age as calculated by FOLPI could be achieved (paragraph 36)
- comparing the results provided in its submission with a rerun of the FOLPI model in which the harvesting of over-mature forests (over 32 years of age) is constrained, and modify the reference level accordingly if necessary (paragraph 36)
- if estimates for natural forests are included in future NIR submissions, making a technical adjustment of the FMRL (paragraph 37)
- agreeing that in the future a technical correction should be made to incorporate the *Harvested wood products* pool (paragraph 38).

A5.1.2 Technical corrections required

For the 2016 submission, the following technical corrections were made to meet Intergovernmental Panel on Climate Change (IPCC) guidance and address recommendations by the UNFCCC ERT. These aimed to:

1. ensure consistency between the method used for greenhouse gas reporting of *Forest management* and that used to calculate the FMRL (IPCC, 2014, sections 2.7.5.2 and 2.7.6). This involved making changes to:
 - a. align forest area estimates
 - b. align CEF emissions calculation methods
 - c. include overplanting estimates (pre-1990 natural forest conversions to pre-1990 planted forest)
 - d. include non-CO₂ emissions
2. include an estimate for pre-1990 natural forest emissions following completion of the re-measurement of the pre-1990 natural forest inventory and subsequent analysis
3. address new elements of Decision 2/CMP.7 including:
 - a. accounting for *Harvested wood products* (processed domestically)
 - b. the application of the natural disturbances provision.

For the 2019 submission, an additional technical correction was applied to the FMRL to capture improvements to the *Harvested wood products* estimates on exported, unprocessed logs.

For the 2021 submission, a technical correction was made to address a number of methodological inconsistencies between emissions estimated for *Forest management* and those used to calculate the FMRL, including:

1. aligning pre-1990 natural forest methods
2. aligning the pre-1990 planted forest yield tables
3. correcting a model assumption to align the harvesting projections with the *Harvested wood products* estimates
4. correcting the background disturbance level and aligning this calculation with the FMRL.

For the 2022 submission, the final submission under the 2013 to 2020 commitment period, a further technical correction has been made. The FMRL has been updated to respond to preliminary recommendations the ERT made during the review of the 2021 submission, as well as to align the methods used to calculate emissions from *Forest management* with those used to calculate the FMRL. In summary, the updates that have been applied to the technically corrected FMRL involve:

1. applying a number of changes to pre-1990 planted forest in the calculation of emissions for *Forest management*. Updates to the FMRL have been applied to ensure consistency between the methods used to calculate *Forest management* and the FMRL. They include:
 - a. updating the harvest area by age and forest age profile calculations
 - b. using input data to produce the FMRL based on the method change applied for calculating the forest age profile and harvest age profile
2. updating the planted forest model to ensure that the forest age profile at 2009 is consistent with the NIR
3. removing the projected rate of deforestation and carbon equivalent forests for pre-1990 planted forests and natural forests from 2010 to 2020. The area of both forests is now assumed to remain constant from 2009 onwards
4. removing the previous technical correction applied to include net emissions from overplanting between 2010 and 2013, where pre-1990 natural forest is removed and replaced with planted forest.

5. updating the yield tables used for measuring *Forest management* emissions for pre-1990 planted forest to make it consistent with the approach for reporting *Forest management* emissions for pre-1990 planted forest
6. updating the emission factors and area estimates for pre-1990 natural forest to make them consistent with the approach for reporting *Forest management* emissions for pre-1990 natural forest
7. updating the *Harvested wood products* model input data to make them consistent with the revised above-ground biomass (AGB) harvest removals as part of this technical correction (to align with items 5 and 6 below), and so that the model approach and assumptions are consistent with the approach used to estimate emissions from *Forest management*.

Further detail on each of these updates to the FMRL is provided in the section below.

Technical corrections: Addressing methodological inconsistencies between the 2011 FMRL and *Forest management* reporting

Replicate the FMRL using the inventory reporting system

The first step taken to calculate technical corrections to the FMRL was to replicate the FMRL as submitted in 2011, applying the same policy assumptions, but using the reporting system and historical data that are used to report on *Forest management* in the inventory.

This technical correction addresses two of the findings of the technical assessment (listed above) by:

1. maintaining consistency in the fraction of harvested biomass instantly oxidised
2. ensuring consistency between the emissions reported in the inventory for *Forest management* and the FMRL.

The original 2011 FMRL submissions assumed that 85 per cent of stem carbon is removed on harvest. For the estimate for pre-1990 planted forest, the NIR assumed that 70 per cent of AGB is removed on harvest. These two ratios are roughly equivalent, when converting stem carbon to AGB.

As the technically corrected forest management reference level (FMRL_{corr}) was re-created using the same reporting system as used for *Forest management* in the inventory, a technical correction was applied so that 70 per cent of AGB is assumed to be removed on harvest.

To ensure consistency in historical emissions from planted forest emissions reported in chapter 11 for *Forest management* and the FMRL_{corr}, the same underlying activity data and emission factor data were used for the FMRL_{corr} up to 2009. This results in the following updates to be made to the FMRL:

- adjusting the forest area estimate to be consistent at 2009
- adjusting the planted forest age profile to be consistent at 2009
- adjusting the harvest and deforestation area by age estimates to be consistent from 1990 to 2009.

Further details of these technical corrections to historical activity data are provided below.

Aligning forest area estimates

The total area for pre-1990 planted forest as at 2009 for the 2011 FMRL was not consistent with the area estimate for the *Forest management*.

The 2011 FMRL submission was based on data derived from the 2010 NEFD (Ministry for Primary Industries, 2010). The NEFD is an annual survey of forest owners that represents the 'net stocked area' of the planted production forest estate established with the primary intention of producing wood or fibre. The Land Use and Carbon Analysis System (LUCAS), which is used for reporting emissions for *Forest management* in the inventory, uses complete wall-to-wall mapping to estimate forest area. This means LUCAS maps to a 'gross stocked area' where harvested areas, skid sites, forest roads and unstocked gullies are included in the mapped forest area. This gross stocked area is also the basis for the national sampling system used for deriving emission factors for the *Forest land* use classes.


For modelling emissions for reporting under the UNFCCC, LUCAS has isolated the net stocked area from the mapped gross stocked area so the modelled area is compatible between the two data sources (LUCAS and NEFD). The LUCAS gross stocked area of pre-1990 planted forest area is 1.47 million hectares as at 2009. The LUCAS net stocked area, as at 2009, is estimated to be 1.25 million hectares (a 12.4 per cent difference).

In comparison, 1.14 million hectares was estimated from the NEFD as at 2009 for the 2011 FMRL. Because the 2011 FMRL did not take into account differences in the data sources due to the two purposes for which the data are collected, a technical correction is required to correct the original NEFD-based FMRL to the LUCAS mapped area estimates used for reporting for *Forest management*.

Aligning the planted forest age profile at 2009

A technical correction was applied to the pre-1990 planted forest age profile at 2009, to be consistent with the forest age profile reported for pre-1990 planted forest (*Forest management*) in this submission (see figure A5.1.1). This was achieved by using the same forest area, harvesting and deforestation data up to 2009 as used for reporting *Forest management*. This required an update to the estimated harvest and deforestation area at 2009, which was based on a projection in the original 2011 FMRL.

Figure A5.1.1 Pre-1990 planted forest age profile as at 2009, as reported for the original 2011 FMRL, the FMRL_{corr} and *Forest management*



Harvest data

The annual harvest area estimate from 2010 to 2020 remains the same as the original 2011 FMRL. This is because this estimate of harvest area is considered to be part of the underlying policy assumptions of the original 2011 FMRL, and so would not meet the criteria for a technical correction.

A technical correction was applied to the average harvest age from 2010 to 2020. The 2011 FMRL predicted an average harvest age of between 31 and 33 years from 2010 to 2020. A technical correction was applied to adjust this down to range between 28 and 30 years over this period (see table A5.1.1). This is to address issue 9 that the ERT raised in the technical assessment of the FMRL submitted in 2011 (UNFCCC, 2011): that harvest ages in the projection were older than those observed historically and there were no policies in place that would influence rotation length or change the average harvest ages of planted forests. The updated average harvest ages are now more consistent with the historical time series, while increasing slightly through time to account for the increasing area of older stands that are projected to be available for harvest.

When adjusting the average harvest age down, the same harvest age profile (proportion of harvest area by age) was also included in this FMRL technical correction. This ensures methodological consistency in the approach to estimate harvest area by age with *Forest management*.

Table A5.1.1 Pre-1990 planted forest data used to estimate emissions for the technically corrected FMRL

Year	Pre-1990 planted forest deforestation (kha)	Pre-1990 planted forest harvested (kha)	Pre-1990 planted forest harvest average age (years)
1990	–	19.288	28.9
1991	–	19.801	29.0
1992	–	22.557	29.0
1993	–	23.194	28.9
1994	–	24.919	28.8
1995	–	29.194	28.7
1996	–	31.169	28.3
1997	–	32.093	28.2
1998	–	31.494	27.9
1999	–	33.994	27.6
2000	2.746	34.726	27.2
2001	2.668	38.541	27.3
2002	2.075	45.290	27.4
2003	3.557	39.643	27.6
2004	7.105	33.260	27.7
2005	12.852	27.760	28.1
2006	16.175	27.765	28.4
2007	21.463	23.168	28.5
2008	3.773	37.425	28.6
2009	5.561	37.160	28.8
2010	0	33.086	28.4
2011	0	37.479	28.4
2012	0	41.354	28.6
2013	0	46.112	28.7
2014	0	50.021	28.9
2015	0	49.697	29.2
2016	0	49.724	29.4
2017	0	50.018	29.8

Year	Pre-1990 planted forest deforestation (kha)	Pre-1990 planted forest harvested (kha)	Pre-1990 planted forest harvest average age (years)
2018	0	49.967	30.1
2019	0	45.817	30.2
2020	0	43.817	30.5

Note: kha = kilohectare.

Carbon equivalent forests

Projections for changes in carbon stocks due to CEF were included in the 2011 FMRL. Technical corrections to the FMRL for CEF had previously been applied to ensure consistency with the provisions of Decision 2/CMP.7 (UNFCCC, 2012) and the guidance for reporting (IPCC, 2014). However, following the review of the 2020 submission (see KL.16, 2019) and the 2021 submission, the step of removing CEF from the FMRL entirely was discussed with the ERT, and included in the ERT's preliminary findings of the 2021 submission. This is because the establishment of a CEF is a deviation from the 'business-as-usual' management of forest land. As such, its impact should not be included in the FMRL and the decision to apply the CEF provision does not trigger a technical correction (IPCC, 2014, p.2.101).

Overplanting

Overplanting is where pre-1990 natural forest is converted to planted forest. In previous submissions, a technical correction has been applied to account for the emissions that were projected to occur as a result of this management practice. However, the ERT review of the 2019 submission noted in KL.14, 2019 (UNFCCC, 2019) that this technical correction was not in accordance with the Kyoto Protocol Supplement (IPCC, 2014, section 2.7.5.1) nor with guidance provided in appendix II to Decision 2/CMP.6 because the conversion of natural forest to a forest plantation is considered a change in management practice.

This finding was again noted by the ERT during the review of the 2021 submission. Therefore, the technical correction for overplanting has been removed from this submission. Note that this change for the 2022 submission does not deviate from the original FMRL and corrections for overplanting are no longer included in the FMRL_{corr}. This explanatory text has been included in the annex to explain why a previously applied technical correction has been removed and to demonstrate that the ERT recommendation KL.14, 2019 (UNFCCC, 2019) has been addressed.

Non-carbon dioxide emissions

Non-carbon dioxide (non-CO₂) emissions were not included in the 2011 FMRL submission; therefore, a technical correction is required to include these emissions. Non-CO₂ emissions are estimated based on the average controlled burning from 1990 to 2009, the minimum historical level for wildfire and the area of planted forest under *Forest management* in 2009 to estimate nitrous oxide (N₂O) emissions from drained organic soils associated with *Forest management*.

Controlled burning

Emissions from the burning of pre-1990 planted forest harvest residues are now included. The harvest rate is as per the FMRL, and the proportion burned is that applied to the LULUCF category *Forest land remaining forest land* during the first commitment period of the Kyoto Protocol.

Burning of residues associated with conversions of pre-1990 natural forest to pre-1990 planted forest is included. It is assumed to occur at the same rate as reported during the first commitment period.

Wildfire emissions

Wildfires are hard to predict and are influenced by inter-annual climatic conditions and regional drought. To estimate emissions from wildfire, the default methodology described in section 2.3.9.6 of the Kyoto Protocol Supplement (IPCC, 2014) has been applied:

...the value of the mean [of natural disturbance time series data] plus two times the standard deviation is calculated using the entire time series of data in the calibration period. Any outlier value (ie above mean plus two times the standard deviation) is removed. This process is repeated until there are no outliers.

The default method calibration period has been applied between 1990 and 2009. This approach is taken to be consistent with New Zealand's background level of natural disturbance.

Nitrous oxide emissions

It is assumed that there are no N₂O emissions from fertilisation of forests within the FMRL. These are minor and captured within the Agriculture sector.

In the 2022 submission, New Zealand has reported on N₂O emissions, as a result of oxidation of organic matter, from the drainage of organic soils for *Forest management*. Emissions are estimated following the methodology outlined in the 2006 IPCC Guidelines (IPCC, 2006) and described in chapter 6, section 6.10.2. As the FMRL assumes no deforestation in pre-1990 forest, the area of drained organic soils under *Forest management* is assumed to remain constant through the commitment period at the area reported in 2009.

Natural disturbance

Emissions from natural disturbance events were not originally considered in the calculation of the 2011 FMRL. New Zealand has reported its intention to apply the natural disturbance provision and, for *Forest management*, the background level has been set using the default method described in section 2.3.9.6 of the Kyoto Protocol Supplement (IPCC, 2014). This is included in the estimate of the non-CO₂ emissions as described above.

However, emissions from, and associated with, salvage logging cannot be excluded from accounting during the second commitment period.¹⁰ This means that, when developing the natural disturbance background level, historical emissions from natural disturbances should exclude these emissions. New Zealand has not excluded these emissions from the historical data used to calculate its background level of natural disturbance emissions under its technically corrected FMRL. If New Zealand applies the provision to exclude emissions from natural disturbances from its accounting, the background level will then be adjusted to remove these salvage logging emissions.

Pre-1990 planted forest

A technical correction has been made to ensure the yield tables used for pre-1990 planted forest are consistent with the yield tables used to calculate carbon stock and stock change for *Forest management* in the 2022 submission. This includes the application of two period-specific yield tables, one for stands planted before 1990 and one for stand planted from 1990 onwards.

¹⁰ Paragraph 33(c) of annex to Decision 2/CMP.7 contained in document FCCC/KP/CMP/2011/10/Add.1, p 18.

The area of pre-1990 planted forest from 2010 to 2020 in the FMRL_{corr} is constant and consistent with the area under *Forest management* reported in 2009.

Pre-1990 natural forest

Emissions and removals by pre-1990 natural forest were not included in the 2011 FMRL submission. Because pre-1990 natural forest is now included in New Zealand's reporting of emissions for *Forest management* land, a technical correction is required. A technical correction was made to include net emissions from pre-1990 natural forest, which includes:

1. an emission factor, using an annual rate of carbon stock change that is consistent with that reported for pre-1990 natural forest from 1990 to 2009 in the 2022 submission
2. the area of pre-1990 natural forest in 2010 to 2020 in the FMRL_{corr} that is consistent with the area under *Forest management* reported in 2009.

Harvested wood products

Emissions and removals for the *Harvested wood products* pool were not included in the 2011 FMRL submission. The technical correction for the final submission of the 2013 to 2020 period uses the same spreadsheet model as that used for New Zealand's *Forest management* reporting. This uses the same underlying emission factor and activity data from 1990 to 2009 as used for *Forest management* in this submission. A different set of activity data, described in more detail below, from 2013 to 2020 is determined to represent a business-as-usual projection for the FMRL_{corr}. The technical correction made reflects that no government policies were either in place, or being planned, that would increase wood use and/or domestic production between 2013 and 2020.

To estimate emissions from *Harvested wood products* associated with *Forest management* from 2013 to 2020, two assumptions were made to estimate business-as-usual activity.

1. Domestic processing capacity and production of products would remain constant.
2. Projected increases in harvest volume would result in the excess logs being exported as raw products.

The basis for these assumptions was that any change in domestic processing capacity would reflect a change in *Forest management*. The activity data for *Harvested wood products* in the FMRL_{corr} were then calculated.

Projecting total roundwood production

The annual AGB projected to be removed as merchantable timber on harvest is estimated from the FMRL_{corr} projection for pre-1990 planted forests. The projected AGB removals from 2013 to 2020 were then converted to roundwood volume. The conversion of AGB removed on harvest to roundwood volume is based on the average annual ratio of these statistics from 1990 to 2009. The roundwood volume from 1990 to 2009 is sourced from the Ministry for Primary Industries (Ministry for Primary Industries, 2021), and is consistent with the estimate for *Harvested wood products* emissions for this submission.

The ratio of estimated AGB removals to roundwood volume was used, rather than simply converting carbon to volume based on a known carbon fraction, because the estimated AGB removed on harvest does not consistently match estimated roundwood production from 1990 to 2009 (see figure A3.2.10, appendix A.3.2.5 for more details).

Between 1990 and 2009, the estimated AGB removed from harvesting planted forests tends to exceed the estimated roundwood production (Ministry for Primary Industries, 2021) over

this period (see figure A3.2.10, appendix A.3.2.5). As a result, if AGB losses from projected harvesting in the FMRL_{corr} were converted to roundwood volume based on a known carbon fraction, this would likely overpredict the roundwood volume used to create *Harvested wood products* in the FMRL_{corr}, relative to the business-as-usual activity observed over the reference period (1990 to 2009).

Projecting domestic production of Harvested wood products

The activity data for *Harvested wood products* processed domestically in New Zealand over the 2013 to 2020 period are estimated to be the same as the annual average from 2000 to 2009. This period was considered to be representative of New Zealand’s business-as-usual processing of *Harvested wood products*. The annual average production over this period was calculated for each individual semi-finished wood product category and for total roundwood volume processed domestically.

Projecting export production of Harvested wood products

Export production of *Harvested wood products* is calculated from an estimated volume of export roundwood (as described in chapter 11, section 11.3.6). Export roundwood production for the FMRL_{corr} is calculated as the projected total roundwood production minus the projected roundwood volume processed domestically.

The inclusion of exported *Harvested wood products* is in line with paragraph 27 of Decision 2/CMP.7. It follows the methodology provided in table 12.1, chapter 12, volume 4 of the 2006 IPCC Guidelines (IPCC, 2006).

Harvested wood products originating from natural forests

Harvested wood products from pre-1990 natural forest is not included in the FMRL_{corr}. The volume produced from the harvesting of pre-1990 natural forests is less than 0.1 per cent of New Zealand’s total harvest volume (Ministry for Primary Industries, 2015).

A5.1.3 Technical corrections and their impact

The impact of the technical corrections made in the 2016, 2019, 2021 and 2022 submissions to the original FMRL is summarised in table A5.1.2.

This submission covers the period outside the 2013 to 2020 forest management reference level (FMRL), therefore no technical corrections have been made. New Zealand is yet to set the FMRL for 2021 onwards.

Table A5.1.2 Summary of the technical corrections to the FMRL

	Emissions (Mt CO ₂ -e yr ⁻¹)
FMRL	11.150
Technical corrections	
Pre-1990 planted forest	-11.627
Non- CO ₂ (including natural disturbance)	0.077
Pre-1990 natural forest	-1.442
Harvested wood products	-12.497
Sum of technical corrections	-25.489
FMRL_{corr}	-14.339

Note: FMRL = forest management reference level; FMRL_{corr} = technically corrected forest management reference level.

Figure A5.1.2 provides a breakdown of the various components of the technical corrections over the time series.

Figure A5.1.2 Technical corrections to the FMRL by category





Figure A5.1.3 provides a comparison of recalculated estimates with previous estimates. This illustrates the time-series consistency of the estimates.

Figure A5.1.3 Comparison of the 2011 FMRL and total of technical corrections from 2013 to 2020



Note: FMRL = forest management reference level; FMRL_{corr} = technically corrected forest management reference level.

A5.2 Natural disturbance

New Zealand applied the default method described in section 2.3.9.6 of the Kyoto Protocol Supplement (IPCC, 2014) for calculating its background level of natural disturbances for both *Afforestation and reforestation* and *Forest management*. This method has been applied following ERT recommendation KL.10, 2019 (UNFCCC, 2019).

The following types of natural disturbance occur in New Zealand:

- wildfires
- invertebrate and vertebrate pests and diseases
- extreme weather events
- geological disturbances.

In all cases except fire, New Zealand assumed a zero baseline between 1990 and 2009. While other natural disturbance events occurred throughout the calibration period, assumptions were made for the purposes of calculating the background level.

For planted forests reported under *Afforestation and reforestation* and *Forest management*, salvage logging is considered to take place in all disturbed forests.

In the case of pre-1990 natural forests, the ground plot measurement programme captures emissions from natural disturbances implicitly, and the emissions from natural disturbance events, apart from wildfires, cannot be separated from other disturbance events. The stock change estimates reported for natural forests include background levels of small-scale natural disturbance events.

Only direct oxidation of biomass in wildfires is considered for the purposes of calculating a background level of natural disturbance for both *Afforestation and reforestation* and *Forest management* land, regardless of forest type. The data used are as reported under the UNFCCC for the period 1990 to 2009 (see chapter 6, section 6.10.8).

A5.2.1 Afforestation and reforestation

Due to the nature of *Afforestation and reforestation* accounting and reporting methods, the background level of carbon dioxide emissions from natural disturbance is already captured implicitly within the reported estimates. New Zealand separately estimates and reports the non- CO₂ emissions from natural disturbances. The background level has been calculated using the default method described in section 2.3.9.6 of the Kyoto Protocol Supplement (IPCC, 2014). However, both the post-1989 forest area and the carbon stock increase during the calibration period. To account for the annual change, the background level for the calibration period is calculated as a proportion of the post-1989 forest estate. This proportion is then multiplied by the carbon stock in post-1989 forest for each year in the reporting period (2013 to 2020). This approach provides the background level and corrects for the increasing area and age (and therefore carbon stock exposed to natural disturbance) in post-1989 forests.

The *Afforestation and reforestation* background level for the period 2013-2020 was calculated as 2.54 kilotonnes carbon dioxide equivalent (kt CO₂-e).

Avoiding the expectation of net credits or net debits for the application of the natural disturbance provision: Afforestation and reforestation

The background level is calculated using the default methodology described in section 2.3.9.6 of the Kyoto Protocol Supplement (IPCC, 2014). The proportion from the calibration period is then multiplied by the carbon stock in post-1989 forest for each year in the reporting period (2013 to 2020). This approach is taken for the following reasons:

- A trend is observed in natural disturbance emissions during the calibration period for *Afforestation and reforestation*. Emissions from natural disturbances have been increasing throughout the calibration period as the age of these forests and, therefore, biomass increase through time. This trend has continued during the second commitment period. The calibration period was used to obtain an annual emissions value by proportion of carbon stocks and then used to calculate the background level for the 2013 year onwards, based on the carbon stocks of *Afforestation and reforestation* lands in each year.
- Gross:net accounting applies to *Afforestation and reforestation* activities. Emissions from natural disturbances occurring in any year of the commitment period, which fall below the background level, are not excluded from the accounting. Emissions from natural disturbances that are greater than the background level in any year of the commitment period are able to be excluded from the accounting if a Party chooses.
- If emissions from natural disturbances are greater than the background level, they can be excluded from the accounting and there is no expectation of net debits arising. If emissions are less than the background level in any year of the commitment period, all emissions from natural disturbance will still be accounted for. There is no expectation of net debits in this scenario. Under gross:net accounting for *Afforestation and reforestation* activities, it would not be possible to expect net credits when applying this approach to excluding the emissions from natural disturbances.

A5.2.2 Forest management

The background level of natural disturbance for *Forest management* for the period 2013-2020 was calculated as 9.34 kt CO₂-e.

Avoiding the expectation of net credits or net debits for the application of the natural disturbance provision: Forest management

The background level has been calculated using the default methodology described in section 2.3.9.6 of the Kyoto Protocol Supplement (IPCC, 2014). Using this method, the expectation of net credits or net debits for the application of the natural disturbance provision is avoided for the following reasons.

- There is no observed trend in natural disturbance emissions during the calibration period for *Forest management* and therefore none can be expected during the second commitment period.
- Any emissions from natural disturbances during the commitment period that fall below the background level are not excluded from the accounting. During the commitment period, emissions from natural disturbances that are above the background level are, subject to New Zealand's discretion, able to be excluded from the accounting.
- The accounting for *Forest management* is against a projected business-as-usual FMRL. The background level is included implicitly within the FMRL, and any emissions greater than the background level can be excluded from the accounting.

A5.3 Carbon equivalent forests

Information on CEF is provided in aggregated form in CRF table 4(KP-I)B.1.2. Details of each application that makes up the reported estimates are provided in table A5.3.1.

Table A5.3.1 Breakdown of carbon equivalent forests by domestic scheme application in 2021

Scheme ID	Management type	2021
CEF – 2	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	1.24
CEF – 3	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	1.29
CEF – 4	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.26
CEF – 8	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.26
CEF – 9	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.12
CEF – 11	Newly established (ha)	–
	Harvested and converted (ha)	47.32
	Net change (t C)	–8.47

Scheme ID	Management type	2021
CEF – 12	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.79
CEF – 13	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.51
CEF – 14	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.58
CEF – 15	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.76
CEF – 17	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.03
CEF – 18	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.22
CEF – 19	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.19
CEF – 20	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.05
CEF – 21	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.67
CEF – 24	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.08
CEF – 25	Newly established (ha)	–
	Harvested and converted (ha)	69.27
	Net change (t C)	–18.17
CEF – 27	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.17
CEF – 31	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.02
CEF – 35	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.01
CEF – 36	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.16

Scheme ID	Management type	2021
CEF – 38	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.01
CEF – 39	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.10
CEF – 40	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.04
CEF – 41	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.01
CEF – 42	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.07
CEF – 43	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.04
CEF – 44	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.02
CEF – 45	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.01
CEF – 47	Newly Established (ha)	–
	Harvested and Converted (ha)	–
	Net change (t C)	0.01
CEF – 49	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.02
CEF – 51	Newly established (ha)	–
	Harvested and converted (ha)	–
	Net change (t C)	0.00
TOTAL	Newly established (ha)	–
	Harvested and converted (ha)	116.59
	Net change (t C)	–18.92

Note: CEF = carbon equivalent forest.

Annex 5: References

- IPCC. 2006. Eggleston HS, Buendia L, Miwa K, Ngara T, Tanabe K (eds). *2006 IPCC Guidelines for National Greenhouse Gas Inventories. Volume 4. Agriculture, Forestry and Other Land Use*. IPCC National Greenhouse Gas Inventories Programme. Japan: Institute for Global Environmental Strategies for IPCC.
- IPCC. 2014. Hiraishi T, Krug T, Tanabe K, Srivastava N, Baasansuren J, Fukuda M, Troxler TG (eds). *2013 Revised Supplementary Methods and Good Practice Guidance Arising from the Kyoto Protocol*. Switzerland: IPCC.
- Ministry for Primary Industries. 2010. *National Exotic Forest Description as at 1 April 2009*. Wellington: Ministry for Primary Industries.
- Ministry for Primary Industries. 2015. *Log and Roundwood Removal Statistics*. Retrieved from www.mpi.govt.nz/news-and-resources/statistics-and-forecasting/forestry (July 2015).
- Ministry for the Environment. 2015. *New Zealand's Greenhouse Gas Inventory 1990–2013*. Wellington: Ministry for the Environment.
- Ministry for Primary Industries. 2021. *Wood processing: Quarterly production, stock and roundwood removals, June 2021 quarter production*. Wellington: Ministry for Primary Industries. Retrieved from www.mpi.govt.nz/forestry/new-zealand-forests-forest-industry/forestry/wood-processing (27 September 2021).
- New Zealand Government. 2011. *Forest Management Reference Level Submission*. Retrieved from unfccc.int/files/meetings/ad_hoc_working_groups/kp/application/pdf/newzealand_frml.pdf (22 February 2016).
- UNFCCC. 2011. *Report of the technical assessment of the forest management reference level submission of New Zealand submitted in 2011*. Retrieved from unfccc.int/resource/docs/2011/tar/nzl01.pdf (22 February 2016).
- UNFCCC. 2012. FCCC/KP/CMP/2011/10/Add.1. *Report of the Conference of the Parties serving as the meeting of the Parties to the Kyoto Protocol on its seventh session, held in Durban from 28 November to 11 December 2011: Addendum – Part 2: Action taken by the Conference of the Parties serving as the meeting of the Parties to the Kyoto Protocol at its seventh session*.
- UNFCCC. 2019. FCCC/ARR/2019/NZL. *Report on the individual review of the annual submission of New Zealand submitted in 2019. Note by the expert review team*.

Annex 6: Additional information on the inventory system and inventory completeness

A6.1 Quality assurance and quality control processes


Quality assurance and quality control (QA/QC) processes have a significant role in the preparation of the inventory, to ensure the core principles of transparency, accuracy, completeness, comparability and consistency are achieved. Table A6.1.1 describes the main QA/QC processes used in the preparation of the inventory. These processes are under continual review and improvement, to ensure they are fit for purpose.

Table A6.1.1 Quality assurance and quality control processes used in preparation of the inventory

ID	QA/QC process or activity description
QA file	All external reviews of the whole or part of the inventory are documented in the QA file. Reviews are performed by qualified personnel, and the review records are included in the submission of the inventory to the United Nations Framework Convention on Climate Change. These reviews help identify improvements to the inventory.
QC 1	Planned recalculations and improvements are approved by the Reporting Governance Group that oversees international climate change reporting by the New Zealand Government. The role of this group is further described in chapter 1.
QC 2	If planned improvements affect emission factors, parameters, methodologies or activity data sources, they are peer reviewed before being implemented. Some sectors have a dedicated panel of experts who review improvements.
QC 3	Tier 1 checklist QC sheets are completed to ensure transparency, accuracy, completeness, comparability and consistency principles are met. Examples are included in the submission of the inventory.
QC 4	The chapter text for each sector is peer reviewed and follows the checklist provided, to ensure that the peer review is comprehensive and consistent.
QC 5	Recalculations that exceed a certain threshold (see figure A6.1.1) are analysed and clearly documented. This includes changes resulting from planned improvements, errors, recommendations from the expert review team, and changes to guidelines.
QC 7	All sectors in the inventory are approved by the relevant member of the Reporting Governance Group that oversees all international climate change reporting by the New Zealand Government before being submitted to the National Inventory Compiler.
QC 10	Common reporting format QC tools identify any potential issues with the data and are used to ensure the data integrity standards are met.
Sector submission checks	Sector submissions are checked against the data integrity standards and chapter formatting standards by the inventory agency before sector submission. Any issues must be resolved before submitting. This enables the remainder of the inventory compilation to proceed smoothly because quality is assured.

Figure A6.1.1 shows how these QA/QC processes align with the overall preparation of the inventory.

Figure A6.1.1 How the quality assurance and quality control processes and products align with the preparation of the inventory



A6.2 General assessment of completeness

A6.2.1 Emissions reported as 'NE' (not estimated)

According to the United Nations Framework Convention on Climate Change (UNFCCC) reporting guidelines (UNFCCC, 2013), the notation key 'NE' (not estimated) signifies that emissions and/or removals occur but have not been estimated or reported. 'NE' can be applied for the following reasons:

- if emissions of a gas from a category are insignificant, that is, they should not exceed 0.05 per cent of the national total of greenhouse gas (GHG) emissions, and not exceed 500 kilotonnes carbon dioxide equivalent (kt CO₂-e) (paragraph 37(b) of the UNFCCC reporting guidelines)
- if the total national aggregate of estimated emissions for all gases and categories considered insignificant remains below 0.1 per cent of the national total of GHG emissions (paragraph 37(b) of the UNFCCC reporting guidelines)
- when an activity occurs in the Party but the *2006 IPCC Guidelines for National Greenhouse Gas Inventories* (IPCC, 2006) do not provide methodologies to estimate emissions and removals (footnote 6 of the UNFCCC reporting guidelines (UNFCCC, 2013)). If this is the case, the category is considered to be non-mandatory, providing the emissions from the category have not been reported previously.

The UNFCCC reporting guidelines also state that, once emissions from a specific category have been reported in a previous submission, emissions from this specific category shall be reported in subsequent GHG inventory submissions (UNFCCC, 2013).

New Zealand's total (ie, gross) emissions were 76,824.6 kt CO₂-e in 2021. The threshold of 0.1 per cent for New Zealand's 2023 submission is 76.8 kt CO₂-e and the threshold of 0.05 per cent is 38.4 kt CO₂-e. Both values are below 500 kt CO₂-e.

Table A6.2.1 summarises New Zealand's direct GHG emissions reported as 'NE' in the 2023 submission.

Table A6.2.1 Summary of 'NE' (not estimated) entries in 2023 submission

CRF category code	Category	Gas	Explanation
Energy			
1.B.2.a.5	Distribution of oil products	CO ₂ , CH ₄	According to paragraph 37(b) of the United Nations Framework Convention on Climate Change (UNFCCC) reporting guidelines (Decision 24/CP.19), this category is not mandatory. The 2006 IPCC Guidelines do not provide the default Intergovernmental Panel on Climate Change (IPCC) emission factor for calculating Tier 1 estimates of methane (CH ₄) emissions from the distribution of refined oil products. New Zealand has not reported emission estimates from this category in previous submissions.
1.B.2.b.3	Processing	CO ₂ , CH ₄	Fugitive emissions of carbon dioxide (CO ₂) and CH ₄ have not been formally estimated, although a rough estimate of the likely level of emissions indicates that they are insignificant. While emissions from the Kapuni Gas Treatment Plant may include traces of CH ₄ , the level of these emissions has been determined to be insignificant in comparison with national emissions. A conservative estimate (using default emission factors from the 2006 IPCC Guidelines) gives nearly 1.5 kilotonnes carbon dioxide equivalent (kt CO ₂ -e) per year. CH ₄ : 625 Mm ³ (Kapuni field production) * 9.7e-5 * 25 = 1.5 kt CO ₂ -e.

CRF category code	Category	Gas	Explanation
			<p>The conservative estimated value is below 0.05 per cent of New Zealand's total emissions. This would keep the national total aggregate of estimated emissions for all gases and categories considered insignificant below 0.1 per cent of the national total greenhouse gas emissions, which is in line with paragraph 37(b) of the UNFCCC reporting guidelines.</p> <p>Carbon dioxide from gas processing is mostly associated with direct venting through a stack and, therefore, is reported under 1.B.2.c.1, as recommended in the 2017 assessment review report. However, there is a possibility of the presence of trace amounts of CO₂ from processing due to leakage, which is estimated to be no higher than 0.1 per cent of vented CO₂. A conservative estimate of 0.1 per cent of vented CO₂ from all categories is 0.26 kt, which is below 0.05 per cent of New Zealand's total emissions and thus can be considered insignificant.</p>
1.B.2.c.ii	Venting – Gas	CH ₄	<p>While direct venting of CH₄ does not occur, the CO₂ stream may include traces of CH₄. The level of these emissions has been determined to be insignificant in comparison with national emissions. A conservative estimate (using default emission factors from the 2006 IPCC Guidelines) gives nearly 1.5 kt CO₂e per year.</p> <p>CH₄: 625 Mm³ (Kapuni field production) * 9.7e-5 * 25 = 1.5 kt CO₂-e.</p> <p>The conservative estimated value is below 0.05 per cent of New Zealand's total emissions. This would keep the national total aggregate of estimated emissions for all gases and categories considered insignificant below 0.1 per cent of New Zealand's total emissions, which is in line with paragraph 37(b) of the UNFCCC reporting guidelines.</p>
Agriculture			
3.A.4 (for both New Zealand and Tokelau)	Poultry	CH ₄	<p>According to paragraph 37(b) of the UNFCCC reporting guidelines, this category is not mandatory. The 2006 IPCC Guidelines state (page 10.27, vol 4-2) that the Tier 1 method for estimating CH₄ emissions from enteric fermentation for poultry is not developed. Also, table 10.10 (page 10.28, vol 4-2) indicates that there is insufficient research to establish a CH₄ emission factor for poultry for either developed or developing countries.</p>
3.B.2.5	Indirect N ₂ O emissions	N ₂ O	<p>According to footnote 6 in paragraph 37(b) of the UNFCCC reporting guidelines (Decision 24/CP.19), this category is not mandatory for reporting. The 2006 IPCC Guidelines for determining indirect nitrous oxide (N₂O) emissions do not provide a methodology for estimating emissions from leaching and runoff. In addition, indirect N₂O emissions from leaching and runoff are insignificant in New Zealand, because almost all livestock are kept outdoors all year around on pasture.</p>
3.D.1.2.c	Other organic fertilisers applied to soils	N ₂ O	<p>Emissions from <i>Other organic fertilisers applied to soils</i> are not estimated due to their insignificance, as defined in accordance with the UNFCCC reporting guidelines (Decision 24/CP.19, paragraph 37(b)). Emissions are estimated to be approximately 20 kt CO₂-e (van der Weerden et al., 2014). Emissions are below the threshold of 0.05 per cent of New Zealand's total emissions and do not exceed 500 kt CO₂-e.</p>
3.I	Other carbon-containing fertilisers	CO ₂	<p>According to the UNFCCC reporting guidelines (Decision 24/CP.19, paragraph 37), this category is not mandatory because the 2006 IPCC Guidelines do not provide guidance for reporting on other carbon-containing fertilisers. Other carbon-containing synthetic fertilisers besides limestone, dolomite and urea are not applied to agricultural land in New Zealand.</p>

CRF category code	Category	Gas	Explanation
Land Use, Land-Use Change and Forestry			
4.D.1	Forest land, cropland, grassland and wetlands: Drainage and rewetting and other management of organic and mineral soils	CH ₄ , N ₂ O	No methodology is provided in the 2006 IPCC Guidelines for estimating emissions from this source category. According to footnote 6 in paragraph 37(b) of the UNFCCC reporting guidelines (Decision 24/CP.19), this category is not mandatory for reporting.
4.A.2, 4.C.2	Settlements converted to other land uses	CO ₂ (losses)	No methodology is provided in the 2006 IPCC Guidelines for calculating Tier 1 estimates of carbon stocks in living biomass for <i>Settlements</i> . Therefore, with land-use change from <i>Settlements</i> to other land uses, no carbon stock loss is reported. As New Zealand uses the stock change approach, losses are only reported in years new land is converted to this category, or where there is harvesting of forest in the year of conversion. When neither of these activities occurs, the losses are reported with biomass gains (stock change approach) and IE is reported for biomass losses.
4.A.2, 4.C.2	Other land converted to other land uses	CO ₂ (losses)	No methodology is provided in the 2006 IPCC Guidelines for calculating Tier 1 estimates of carbon stocks in living biomass for <i>Other land</i> . Therefore, with land-use change from <i>Other land</i> to other land uses, no carbon stock loss is reported. As New Zealand uses the stock change approach, losses are only reported in years new land is converted to this category, or where there is harvesting of forest in the year of conversion. When neither of these activities occurs, the losses are reported with biomass gains (stock change approach) and IE is reported for biomass losses.
4.A, 4.B, 4.C, 4.D, 4.F	Forest land, Cropland, Grassland, Wetlands and Other land: rewetting and other management of organic and mineral soils	CO ₂	No methodology is provided in the 2006 IPCC Guidelines for estimating emissions from this source category. According to footnote 6 in paragraph 37(b) of the UNFCCC reporting guidelines (Decision 24/CP.19), this category is not mandatory for reporting.
4.B.1	Land converted to Cropland	CO ₂ (deadwood)	No methodology is provided in the 2006 IPCC Guidelines for calculating Tier 1 estimates of carbon stock change in dead organic matter for land-use changes to this land use category. Any emissions reported are for loss of dead organic matter from the previous land use which occurs in the year of conversion.
4.B.1	Cropland remaining cropland/4(V) Biomass burning/ Wildfires/ Cropland remaining cropland	CH ₄ , N ₂ O	New Zealand does not have sufficient information on biomass burning activities to reliably report on it.
4.B.2	Land converted to cropland/4(V) Biomass burning/ Wildfires/ Land converted to cropland	CH ₄ , N ₂ O	New Zealand does not have sufficient information on biomass burning activities to reliably report on it.
4.C.1,4.C.2	Land remaining grassland, Land converted to grassland	CO ₂ (deadwood)	No methodology is provided in the 2006 IPCC Guidelines for calculating Tier 1 estimates of carbon stock change in dead organic matter for land-use changes to this land use category. Any emissions reported are for loss of dead organic matter from the previous land use which occurs in the year of conversion.
4.D.1	Wetlands remaining wetlands/4(V) Biomass burning/ Wildfires/ Wetlands remaining wetlands	CH ₄ , N ₂ O	According to paragraph 37(b) of the UNFCCC reporting guidelines (Decision 24/CP.19), this category is not mandatory because no methodology is provided in the 2006 IPCC Guidelines for calculating Tier 1 estimates of carbon stock changes in organic soils for this land use category. New Zealand does not have sufficient information on biomass burning activities to reliably report on it.
4.D.1	Wetlands remaining wetlands, Land converted to wetlands (except transitions into and out of vegetated wetlands)	CO ₂	No methodology is provided in the 2006 IPCC Guidelines for calculating Tier 1 estimates of carbon stock change in this land use (2006 IPCC Guidelines, Volume 4, Chapter 7.3.1).

CRF category code	Category	Gas	Explanation
Waste			
5.C.1.2.a	Incineration of municipal solid waste	CO ₂ , CH ₄ and N ₂ O	Around 100–200 rural schools in New Zealand still incinerate their waste production. Estimates indicate this practice emits 0.04 kt CO ₂ -e per year. Emissions are not estimated because New Zealand does not have sufficient information regarding the practice of incinerating waste in schools, and the estimated amount is below the significance threshold. This is in accordance with paragraph 37(b) of the UNFCCC reporting guidelines (Decision 24/CP.19).
5.D.1 and 5.D.2	Domestic wastewater and Industrial wastewater	Amount of CH ₄ flared and for energy recovery	Emissions are not estimated because New Zealand does not have any information regarding the CH ₄ flaring in this source category. The amount of CH ₄ flared does not contribute to New Zealand's total emissions because it produces biogenic CO ₂ (as per the 2006 IPCC Tier 1 methodology provided in table 5D of the common reporting format tables).

The estimate of emissions for all of New Zealand's source categories marked as 'NE' results in 23.3 kt CO₂-e, which is below the 0.1 per cent of the total emissions threshold (76.8 kt CO₂-e).

A6.2.2 Emissions reported as 'IE' (included elsewhere)

According to the UNFCCC reporting guidelines (UNFCCC, 2013), the notation key 'IE' (included elsewhere) signifies that emissions and/or removals for this activity or category are estimated and included in the inventory but not presented separately for this category.

Table A6.2.2 details where the notation key 'IE' has been used in this submission of the inventory.

Table A6.2.2 Emissions reported using the 'IE' (included elsewhere) notation key

CRF category code	Category	Source category reported under	Notation key explanation
1.A.2.a	Iron and steel – liquid fuels	1.A.2.g.viii – Other – Liquid fuels	Disaggregated data for this category do not exist.
1.A.2.a	Iron and steel – solid fuels	2.C.1 – Iron and steel production	All emissions from the use of coal in this category are included in the Industrial Processes and Product Use sector because the primary purpose of the coal is to produce iron.
1.A.2.f	Non-metallic minerals – biomass	1.A.2.g.viii – Other – Biomass	Disaggregated data for this category do not exist.
1.A.3.b.ii–iv	Road transportation (other than 'Cars') – all fuels (other than gasoline and diesel)	1.A.3.b.i – Cars	Disaggregated data do not exist for all years for all fuels.
1.A.4.c.ii–iii	Agriculture/forestry/fishing – Off-road vehicles and other machinery/ Gaseous Fuels Agriculture/forestry/fishing – Off-road vehicles and other machinery/ Liquefied Petroleum Gases (LPG) Fishing/Gaseous Fuels	1.A.4.c.i – Agriculture/forestry/fishing – Stationary	Agriculture/forestry/fishing has not been disaggregated into stationary, mobile and fishing for some fuels because data are not available.

CRF category code	Category	Source category reported under	Notation key explanation
1.B.2.b.1	Natural gas/exploration	1.B.2.a.1 – Oil exploration	In New Zealand, oil exploration is not separated from gas exploration by planning, processes, equipment or resources. The exploratory wells are drilled without distinction of their purpose, that is, whether the expected outcome is oil, gas, both or none, and there is no reliable way to predict which it would be to estimate proportions of mostly oil and mostly gas wells. Disaggregated data for oil and gas exploration, therefore, do not exist. Considering that available emission factors for well drilling and testing also do not distinguish between oil and gas, all emissions from oil and gas exploration are reported in the oil exploration category.
1.B.2.c.1.i	Venting/oil and Venting/oil	1.B.2.c.1.iii – Venting/combined	The fields produce both oil and gas and, therefore, are reported as combined. Disaggregated data do not exist.
1.B.2.c.2.i-ii	Flaring/oil and Flaring/oil Flaring/oil and Flaring/gas	1.B.2.c.1.iii – Flaring/combined	The fields produce both oil and gas and, therefore, are reported as combined. Disaggregated data do not exist.
2.A.3	Glass production	2.A.4.b – Other process uses of carbonates/Other uses of soda ash	Carbon dioxide emissions are reported in 2.A.4.b because this aggregates emissions from glass production with other uses of carbonates, due to confidentiality concerns for both glass and aluminium production. A very small number of firms in New Zealand are involved in these activities and use carbonates.
3.A.4	Enteric fermentation/other/buffalo	3.A.1.A – Dairy cattle	A small herd of around 200 buffalo was brought into New Zealand around 2007 for specialised cheese and dairy production. These buffalo are reported within the dairy herd so the notation key 'IE' is used from 2007 onwards.
3.B.1.4, 3.B.2.4	Manure management/other/buffalo	3.B.1.A – Dairy cattle 3.B.2.A – Dairy cattle	For both nitrous oxide (N ₂ O) and methane (CH ₄) emissions, the notation key 'NO' (not occurring) is used up to 2006 because no buffalo were recorded in New Zealand before 2007. A small herd of around 200 buffalo was brought into New Zealand around 2007 for specialised cheese and dairy production. See notation key explanation for 3.A.4. For more information, see chapter 5, section 5.1.4 ('Minor livestock categories') of this National Inventory Report.
3.B.2.5	N ₂ O emissions per MMS ¹¹	3.D – Agricultural soils	Direct N ₂ O emissions from anaerobic lagoons (dairy and swine) and daily spread (swine) are reported under <i>Agricultural soils</i> .

¹¹ MMS stands for a manure management system (see chapter 5).

CRF category code	Category	Source category reported under	Notation key explanation
3.D.1.2.b	Sewage sludge applied to soils	Included under the Waste sector 5.A.1.a	Direct N ₂ O emissions from sewage sludge are reported under 5.A.1.a in the Waste sector. Sewage sludge activity data are obtained from water treatment industry surveys and do not disaggregate the amount of sludge used for different purposes. Due to the small amount of emissions coming from sewage sludge, further disaggregation of the activity data is considered resource prohibitive. Sewage sludge is a very small source of nitrogen (van der Weerden et al., 2014).
3.E	Prescribed burning of savannas	Biomass burning (table 4(V) of LULUCF), category C Grassland	Prescribed burning of savanna is reported under the Land Use, Land-Use Change and Forestry (LULUCF) sector. See chapter 6, section 6.10.8 (Biomass burning (table 4(V) of LULUCF), category C Grassland).
4.A.1/4(V)	Controlled burning	Forest land remaining forest land	Carbon dioxide (CO ₂) emissions are captured by the general carbon stock change calculation if the fire-damaged area is harvested and replanted. If the stand is allowed to grow on but with a reduced stocking, the CO ₂ emissions are reported at the eventual time of harvest.
4.A.1/4(II) 4.B.1/4(II)	Forest land Cropland	Emissions and removals from drainage and rewetting and other management of organic and mineral soils	These emissions are reported under organic soils.
4.A.1	Forest land remaining forest land	Forest land remaining forest land	As New Zealand uses the stock change approach, losses are only reported in years new land is converted to this category, or where there is harvesting of forest in this year. When neither of these activities occurs, the losses are reported with biomass gains (stock change approach) and IE is reported for biomass losses.
4.A.2	Land converted to forest land	Land converted to forest land	Because New Zealand uses the stock change approach, CO ₂ emissions from biomass losses are only reported in years new land is converted to this category, or where there is harvesting of forest in this year. When neither of these activities occurs, the losses are reported with biomass gains (stock change approach) and IE is reported for biomass losses.
4.A.2	Land converted to forest land	Included under the Agriculture sector	New Zealand does not disaggregate data on nitrogen fertiliser by land use; therefore, all N ₂ O emissions are reported in the Agriculture sector.
4.A.2/4(V)	Controlled burning/Land converted to forest land Wildfires/Land converted to forest land	Land converted to forest land	Carbon dioxide emissions are captured by the general carbon stock change calculation if the fire-damaged area is harvested and replanted. If the stand is allowed to grow on but with a reduced stocking, the CO ₂ emissions are reported on at the eventual time of harvest.

CRF category code	Category	Source category reported under	Notation key explanation
4.B.1 4.B.2 4C.1 4C.2	Cropland remaining cropland, Land converted to cropland, Grassland remaining grassland, Land converted to grassland	Cropland remaining cropland, Land converted to cropland, Grassland remaining grassland, Land converted to grassland	New Zealand uses the stock change approach to estimate biomass emissions; therefore, biomass losses are reported with biomass gains and IE is reported.
4.B.1/4(V)	Controlled burning/Cropland remaining cropland	Included under the Agriculture sector	Carbon dioxide and CH ₄ emissions from burning of crop stubble are reported in the Agriculture sector.
4.B.1/4(V)	Wildfires/Cropland remaining cropland	Cropland remaining cropland	Any CO ₂ emissions from wildfires on non-forest land are assumed to be offset by the subsequent carbon stock gain from the regrowth of biomass, which is also not reported. Alternatively, if the wildfire resulted in land-use change, then any CO ₂ emissions are captured by the general carbon stock change calculation that is performed when land is converted to a new land use.
4.B.2/4(V) 4.C.1/4(V) 4.C.2/4(V) 4.D.2/4(V)	Wildfires/Land converted to cropland Wildfires/Grassland remaining grassland Wildfires/Land converted to grassland Wildfires/Land converted to wetlands	Land converted to cropland Grassland remaining grassland Land converted to grassland Land converted to wetlands	Any CO ₂ emissions from wildfires on non-forest land are assumed to be offset by the subsequent carbon stock gain from the regrowth of biomass, which is also not reported. Alternatively, if the wildfire resulted in land-use change, then any CO ₂ emissions are captured by the general carbon stock change calculation that is performed when land is converted to a new land use.
4.A.1/4(I) 4.D.1/4(I) 4.D.2/4(I) 4.E.1/4(I) 4.E.2/4(I)	Direct N ₂ O emissions from nitrogen (N) inputs to managed soils Inorganic N fertilisers and Direct N ₂ O emissions from N inputs to managed soils Organic N fertilisers In the following categories: <ul style="list-style-type: none"> • Forest land remaining forest land • Wetlands remaining wetlands • Land converted to wetlands • Settlements remaining settlements • Land converted to settlements • Settlements remaining settlements • Land converted to settlements 	Included under the Agriculture sector	New Zealand does not hold disaggregated activity data on nitrogen fertiliser by land use; therefore, all N ₂ O emissions from organic and inorganic fertilisers are reported in the Agriculture sector.
4.B.1/4(V)	Controlled burning/Cropland remaining cropland	Included under the Agriculture sector	All emissions from burning of crop stubble are reported in the Agriculture sector.
4.C.1/4(V) 4.D.1/4(V)	Controlled burning/Grassland remaining grassland Controlled burning/Wetlands remaining wetlands Wildfires/Wetlands remaining wetlands	Grassland remaining grassland Wetlands remaining wetlands Wetlands remaining wetlands	This is not a significant activity in New Zealand due to the country's temperate climate and rainfall distribution, and any CO ₂ emissions from burning on non-forest land are assumed to be offset by the subsequent carbon stock gain from the regrowth of biomass, which is also not reported. Alternatively, if the fire resulted in land-use change, then any CO ₂ emissions are captured by the general carbon stock change calculation that is performed when land is converted to a new land use.

CRF category code	Category	Source category reported under	Notation key explanation
4.C.2/4(V) 4.D.2/4(V) 4.E/4(V) 4.F/4(v)	Controlled burning/Land converted to grassland Controlled burning/Land converted to wetlands Biomass burning/Land converted to settlements Biomass burning/Land converted to other land	Land converted to grassland Land converted to wetlands Land converted to settlements Land converted to other land	Carbon dioxide emissions from the controlled burning of land converted to this category are captured by the general carbon stock change calculation that is performed when land is converted to a new land use.
4.D.2.2	Land converted to flooded land	Land converted to open water	New Zealand has not yet identified what area of land converted to open water meets the definition of <i>Land converted to flooded land</i> . Emissions associated with the loss of biomass with land conversion to flooded land will be captured under <i>Land converted to open water</i> .
5.D.1	Domestic wastewater	5.A Solid waste	Activity data – sludge amounts are included under solid waste disposal because sludge is disposed to landfill.
5.D.2	Industrial wastewater	5.A Solid waste	Activity data – sludge amounts are included under <i>Solid waste disposal</i> because sludge is disposed to landfill.
5.D.2	Industrial wastewater	1.A.2.e Food processing, beverages and tobacco – Biomass	Emissions of CH ₄ and N ₂ O from the combustion of biogas from the Tirau dairy processing plant are reported under 1.A.2.e <i>Food processing, beverages and tobacco – Biomass</i> .
Within the Tokelau sector 6, categories 1.A.3.b.i and 1.A.4.c.iii were reported elsewhere	Road transport/Gasoline and diesel oil	Domestic navigation	Tokelau has a small petrol car fleet. In 2018 there were about 40 cars and 30 motorbikes, with an entire road network less than 10 kilometres. Four registered cars were recorded in the 2001 Census. Aluminium boats are the main means of family transport. There were, on average, about 100 outboard motors travelling both outside and within the large lagoons. Petrol use for road transport is relatively small compared with <i>Domestic navigation</i> totals, and all emissions are included in <i>Domestic navigation</i> .
Within the Tokelau sector 6, category 1.A.4.b is reported elsewhere	Residential (1.A.4.b) liquid fuels	Domestic navigation	Only gas used for cooking is reported here. The amount of liquid fuel use in this category is small compared with <i>Domestic navigation</i> and all emissions are included in <i>Domestic navigation</i> .

Annex 6: References

IPCC. 2006. Eggleston HS, Buendia L, Miwa K, Ngara T, Tanabe K (eds). *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. IPCC National Greenhouse Gas Inventories Programme. Japan: Published for the IPCC by the Institute for Global Environmental Strategies.

UNFCCC. 2013. FCCC/CP/2013/10/Add.3. *Report of the Conference of the Parties on its nineteenth session, held in Warsaw from 11 to 23 November 2013, Addendum; Decision 24/CP.19 Revision of the UNFCCC reporting guidelines on annual inventories for Parties included in Annex I to the Convention*.

Van der Weerden A, de Klein C, Kelliher F, Rollo M. 2014. *Reporting to 2006 IPCC Guidelines for N₂O Emissions from Additional Sources of Organic N: Final Report*. MPI Technical Report. Wellington: Ministry for Primary Industries.

Annex 7: Tokelau

A7.1 Emissions estimate data and relevant supporting information by category for Tokelau¹²

Tokelau CRF Table 1.A.1.a: [1. Energy] [1.AA Fuel Combustion – Sectoral approach] [1.A.1 Energy Industries] [1.A.1.a Public Electricity and Heat Production] (Part 1 of 3)

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.1 Energy Industries][1.A.1.a Public Electricity and Heat Production]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Fuel Consumption	TJ	3.27	3.27	3.27	3.27	3.27	3.27	3.27	3.27	3.27	3.27
Liquid fuels	TJ	3.27	3.27	3.27	3.27	3.27	3.27	3.27	3.27	3.27	3.27
Calorific value	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Liquid fuels	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Method											
CO ₂		T1									
CH ₄		T1									
N ₂ O		T1									
Emission factor information											
CO ₂		D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
CO ₂	kt	0.23	0.23	0.23	0.23	0.23	0.23	0.23	0.23	0.23	0.23
Liquid fuels	kt	0.23	0.23	0.23	0.23	0.23	0.23	0.23	0.23	0.23	0.23
CH ₄	kt	0.0000093	0.0000093	0.0000093	0.0000093	0.0000093	0.0000093	0.0000093	0.0000093	0.0000093	0.0000093
Liquid fuels	kt	0.0000093	0.0000093	0.0000093	0.0000093	0.0000093	0.0000093	0.0000093	0.0000093	0.0000093	0.0000093
N ₂ O	kt	0.0000019	0.0000019	0.0000019	0.0000019	0.0000019	0.0000019	0.0000019	0.0000019	0.0000019	0.0000019
Liquid fuels	kt	0.0000019	0.0000019	0.0000019	0.0000019	0.0000019	0.0000019	0.0000019	0.0000019	0.0000019	0.0000019

¹² The category names and CRF codes for source categories are consistent with New Zealand's CRF tables. Only the tables that include reported emissions (by value, IE or NE) are included. For explanations and methodological issues, please refer to chapter 8.

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.1 Energy Industries][1.A.1.a Public Electricity and Heat Production]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Amount captured											
CO ₂	kt	NO									
Liquid fuels	kt	NO									
Implied emission factor											
CO ₂											
Liquid fuels	t/TJ	70.39	70.39	70.39	70.39	70.39	70.39	70.39	70.39	70.39	70.39
CH ₄											
Liquid fuels	kg/TJ	2.85	2.85	2.85	2.85	2.85	2.85	2.85	2.85	2.85	2.85
N ₂ O											
Liquid fuels	kg/TJ	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57

Tokelau CRF Table 1.A.1.a: [1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.1 Energy Industries][1.A.1.a Public Electricity and Heat Production] (Part 2 of 3)

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.1 Energy Industries][1.A.1.a Public Electricity and Heat Production]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Fuel Consumption											
Liquid fuels	TJ	3.27	3.27	3.27	3.27	9.81	16.34	16.34	16.34	16.34	16.34
Calorific value											
Liquid fuels	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Method											
CO ₂		T1									
CH ₄		T1									
N ₂ O		T1									
Emission factor information											
CO ₂		D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
CO ₂	kt	0.23	0.23	0.23	0.23	0.69	1.15	1.15	1.15	1.15	1.15
Liquid fuels	kt	0.23	0.23	0.23	0.23	0.69	1.15	1.15	1.15	1.15	1.15
CH ₄	kt	0.0000093	0.0000093	0.0000093	0.0000093	0.0000279	0.0000466	0.0000466	0.0000466	0.0000466	0.0000466
Liquid fuels	kt	0.0000093	0.0000093	0.0000093	0.0000093	0.0000279	0.0000466	0.0000466	0.0000466	0.0000466	0.0000466

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.1 Energy Industries][1.A.1.a Public Electricity and Heat Production]												
	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	
N ₂ O	kt	0.0000019	0.0000019	0.0000019	0.0000019	0.0000056	0.0000093	0.0000093	0.0000093	0.0000093	0.0000093	
Liquid fuels	kt	0.0000019	0.0000019	0.0000019	0.0000019	0.0000056	0.0000093	0.0000093	0.0000093	0.0000093	0.0000093	
Amount captured												
CO ₂	kt	NO										
Liquid fuels	kt	NO										
Implied emission factor												
CO ₂												
Liquid fuels	t/TJ	70.39	70.39	70.39	70.39	70.39	70.4	70.4	70.4	70.4	70.4	
CH ₄												
Liquid fuels	kg/TJ	2.85	2.85	2.85	2.85	2.85	2.85	2.85	2.85	2.85	2.85	
N ₂ O												
Liquid fuels	kg/TJ	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	

Tokelau CRF Table 1.A.1.a: [1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.1 Energy Industries][1.A.1.a Public Electricity and Heat Production] (Part 3 of 3)

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.1 Energy Industries][1.A.1.a Public Electricity and Heat Production]													
	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Fuel Consumption	TJ	16.34	16.34	12.97	2.86	2.86	2.86	3.05	3.24	3.42	3.61	3.21	7.42
Liquid fuels	TJ	16.34	16.34	12.97	2.86	2.86	2.86	3.05	3.24	3.42	3.61	3.21	7.42
Calorific value	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Liquid fuels	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Method													
CO ₂		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
CH ₄		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
N ₂ O		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
Emission factor information													
CO ₂		D	D	D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D	D	D
Emissions													

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.1 Energy Industries][1.A.1.a Public Electricity and Heat Production]													
	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
CO ₂	kt	1.15	1.15	0.91	0.2	0.2	0.2	0.21	0.23	0.24	0.25	0.23	0.52
Liquid fuels	kt	1.15	1.15	0.91	0.2	0.2	0.2	0.21	0.23	0.24	0.25	0.23	0.52
CH ₄	kt	0.0000466	0.0000466	0.000037	0.0000082	0.0000082	0.0000082	0.0000087	0.0000092	0.0000098	0.0000103	0.0000091	0.0000212
Liquid fuels	kt	0.0000466	0.0000466	0.000037	0.0000082	0.0000082	0.0000082	0.0000087	0.0000092	0.0000098	0.0000103	0.0000091	0.0000212
N ₂ O	kt	0.0000093	0.0000093	0.0000074	0.0000016	0.0000016	0.0000016	0.0000017	0.0000018	0.000002	0.0000021	0.0000018	0.0000042
Liquid fuels	kt	0.0000093	0.0000093	0.0000074	0.0000016	0.0000016	0.0000016	0.0000017	0.0000018	0.000002	0.0000021	0.0000018	0.0000042
Amount captured													
CO ₂	kt	NO											
Liquid fuels	kt	NO											
Implied emission factor													
CO ₂													
Liquid fuels	t/TJ	70.4	70.4	70.39	70.4	70.4	70.4	70.39	70.39	70.39	70.39	70.4	70.39
CH ₄													
Liquid fuels	kg/TJ	2.85	2.85	2.85	2.85	2.85	2.85	2.85	2.85	2.85	2.85	2.85	2.85
N ₂ O													
Liquid fuels	kg/TJ	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57	0.57

Tokelau CRF Table 1.A.3.b.i: [1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Gasoline] (Part 1 of 3)

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Gasoline]													
	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999		
Fuel consumption	TJ	IE											
Calorific value		GCV											
Method													
CO ₂		T1											
CH ₄		T1											
N ₂ O		T1											

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Gasoline]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Emission factor information											
CO ₂		D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
CO ₂	kt	IE									
CH ₄	kt	IE									
N ₂ O	kt	IE									
Implied emission factor											
CO ₂	t/TJ	NA									
CH ₄	kg/TJ	NA									
N ₂ O	kg/TJ	NA									

Note: This category is included under 1.A.3.d. For explanation please refer to section 8.2.5.

Tokelau CRF Table 1.A.3.b.i: [1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Gasoline] (Part 2 of 3)

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Gasoline]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Fuel consumption	TJ	IE									
Calorific value	GCV										
Method											
CO ₂		T1									
CH ₄		T1									
N ₂ O		T1									
Emission factor information											
CO ₂		D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
CO ₂	kt	IE									
CH ₄	kt	IE									
N ₂ O	kt	IE									

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Gasoline]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Implied emission factor											
CO ₂	t/TJ	NA									
CH ₄	kg/TJ	NA									
N ₂ O	kg/TJ	NA									

Tokelau CRF Table 1.A.3.b.i: [1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Gasoline] (Part 3 of 3)

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Gasoline]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Fuel consumption	TJ	IE											
Calorific value	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Method													
CO ₂		T1											
CH ₄		T1											
N ₂ O		T1											
Emission factor information													
CO ₂		D	D	D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D	D	D
Emissions													
CO ₂	kt	IE											
CH ₄	kt	IE											
N ₂ O	kt	IE											
Implied emission factor													
CO ₂	t/TJ	NA											
CH ₄	kg/TJ	NA											
N ₂ O	kg/TJ	NA											

Tokelau CRF Table 1.A.3.b.i Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Diesel Oil] (Part 1 of 3)

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Diesel Oil]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Fuel consumption	TJ	IE									
Calorific value		IE									
Method											
CO ₂		T1									
CH ₄		T1									
N ₂ O		T1									
Emission factor information											
CO ₂		D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
CO ₂	kt	IE									
CH ₄	kt	IE									
N ₂ O	kt	IE									
Implied emission factor											
CO ₂	t/TJ	NA									
CH ₄	kg/TJ	NA									
N ₂ O	kg/TJ	NA									

Tokelau CRF Table 1.A.3.b.i Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Diesel Oil] (Part 2 of 3)

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Diesel Oil]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Fuel consumption	TJ	IE									
Calorific value		IE									
Method											
CO ₂		T1									
CH ₄		T1									
N ₂ O		T1									

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Diesel Oil]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Emission factor information											
CO ₂		D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
CO ₂	kt	IE									
CH ₄	kt	IE									
N ₂ O	kt	IE									
Implied emission factor											
CO ₂	t/TJ	NA									
CH ₄	kg/TJ	NA									
N ₂ O	kg/TJ	NA									

Tokelau CRF Table 1.A.3.b.i Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.3 Transport] [1.A.3.b Road Transportation][1.A.3.b.i Cars][Diesel Oil] (Part 3 of 3)

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Diesel Oil]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Fuel consumption	TJ	IE											
Calorific value		IE											
Method													
CO ₂		T1											
CH ₄		T1											
N ₂ O		T1											
Emission factor information													
CO ₂		D	D	D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D	D	D
Emissions													
CO ₂	kt	IE											
CH ₄	kt	IE											
N ₂ O	kt	IE											

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.3 Transport][1.A.3.b Road Transportation][1.A.3.b.i Cars][Diesel Oil]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Implied emission factor													
CO ₂	t/TJ	NA											
CH ₄	kg/TJ	NA											
N ₂ O	kg/TJ	NA											

Tokelau CRF Table 1.A.3.d Gas/Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.3 Transport][1.A.3.d Domestic Navigation][Gas/Diesel Oil] (Part 1 of 3)

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.3 Transport][1.A.3.d Domestic Navigation][Gas/Diesel Oil]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Fuel consumption	TJ	12.76	12.98	13.21	13.43	13.66	13.89	14.11	14.34	14.56	14.79
Calorific value	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Method											
CO ₂		T1									
CH ₄		T1									
N ₂ O		T1									
Emission factor information											
CO ₂		D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
CO ₂	kt	0.9	0.91	0.93	0.95	0.96	0.98	0.99	1.01	1.03	1.04
CH ₄	kt	0.0000848	0.0000863	0.0000878	0.0000893	0.0000908	0.0000923	0.0000938	0.0000953	0.0000968	0.0000983
N ₂ O	kt	0.0000242	0.0000247	0.0000251	0.0000255	0.000026	0.0000264	0.0000268	0.0000272	0.0000277	0.0000281
Implied emission factor											
CO ₂	t/TJ	70.39	70.39	70.39	70.39	70.39	70.4	70.4	70.39	70.39	70.4
CH ₄	kg/TJ	6.65	6.65	6.65	6.65	6.65	6.65	6.65	6.65	6.65	6.65
N ₂ O	kg/TJ	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9

Tokelau CRF Table 1.A.3.d Gas/Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.3 Transport][1.A.3.d Domestic Navigation][Gas/Diesel Oil] (Part 2 of 3)

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.3 Transport][1.A.3.d Domestic Navigation][Gas/Diesel Oil]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Fuel consumption	TJ	15.01	15.24	15.47	15.69	15.92	16.14	16.37	16.59	16.82	17.04
Calorific value	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Method											
CO ₂		T1									
CH ₄		T1									
N ₂ O		T1									
Emission factor information											
CO ₂		D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
CO ₂	kt	1.06	1.07	1.09	1.1	1.12	1.14	1.15	1.17	1.18	1.2
CH ₄	kt	0.0000998	0.0001013	0.0001028	0.0001043	0.0001058	0.0001073	0.0001088	0.0001103	0.0001118	0.0001133
N ₂ O	kt	0.0000285	0.000029	0.0000294	0.0000298	0.0000302	0.0000307	0.0000311	0.0000315	0.000032	0.0000324
Implied emission factor											
CO ₂	t/TJ	70.4	70.39	70.4	70.4	70.4	70.39	70.39	70.39	70.39	70.39
CH ₄	kg/TJ	6.65	6.65	6.65	6.65	6.65	6.65	6.65	6.65	6.65	6.65
N ₂ O	kg/TJ	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9

Tokelau CRF Table 1.A.3.d Gas/Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.3 Transport][1.A.3.d Domestic Navigation][Gas/Diesel Oil] (Part 3 of 3)

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.3 Transport][1.A.3.d Domestic Navigation][Gas/Diesel Oil]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Fuel consumption	TJ	17.27	17.5	17.72	17.95	18.17	18.03	18.89	19.88	21.08	30.91	29.17	19.46
Calorific value	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Method													
CO ₂		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
CH ₄		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.3 Transport][1.A.3.d Domestic Navigation][Gas/Diesel Oil]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
N ₂ O	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
Emission factor information													
CO ₂	D	D	D	D	D	D	D	D	D	D	D	D	D
CH ₄	D	D	D	D	D	D	D	D	D	D	D	D	D
N ₂ O	D	D	D	D	D	D	D	D	D	D	D	D	D
Emissions													
CO ₂	kt	1.22	1.23	1.25	1.26	1.28	1.27	1.33	1.4	1.48	2.18	2.05	1.37
CH ₄	kt	0.0001148	0.0001163	0.0001179	0.0001194	0.0001209	0.0001199	0.0001256	0.0001322	0.0001402	0.0002056	0.000194	0.0001294
N ₂ O	kt	0.0000328	0.0000332	0.0000337	0.0000341	0.0000345	0.0000343	0.0000359	0.0000378	0.00004	0.0000587	0.0000554	0.000037
Implied emission factor													
CO ₂	t/TJ	70.4	70.39	70.4	70.39	70.39	70.4	70.39	70.4	70.39	70.39	70.39	70.39
CH ₄	kg/TJ	6.65	6.65	6.65	6.65	6.65	6.65	6.65	6.65	6.65	6.65	6.65	6.65
N ₂ O	kg/TJ	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9	1.9

Tokelau CRF Table 1.A.4.b: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.4 Other Sectors][1.A.4.b Residential] (Part 1 of 3)

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.4 Other Sectors][1.A.4.b Residential]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Fuel Consumption	TJ	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16
Liquid fuels	TJ	IE									
Gaseous fuels	TJ	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16
Calorific value		GCV									
Liquid fuels		GCV									
Gaseous fuels		GCV									
Method											
CO ₂		T1									
CH ₄		T1									
N ₂ O		T1									

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.4 Other Sectors][1.A.4.b Residential]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Emission factor information											
CO ₂		D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
CO ₂	kt	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12
Liquid fuels	kt	IE									
Gaseous fuels	kt	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12
CH ₄	kt	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204
Liquid fuels	kt	IE									
Gaseous fuels	kt	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204
N ₂ O	kt	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004
Liquid fuels	kt	IE									
Gaseous fuels	kt	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004
No _x	kt	NE									
CO	kt	NE									
NM VOC	kt	NE									
SO ₂	kt	NE									
Amount captured											
CO ₂	kt	NO									
Liquid fuels	kt	NO									
Gaseous fuels	kt	NO									
Implied emission factor											
CO ₂											
Liquid fuels	t/TJ	NO									
Gaseous fuels	t/TJ	56.79	56.79	56.79	56.79	56.79	56.79	56.79	56.79	56.79	56.79
CH ₄											
Liquid fuels	kg/TJ	NO									
Gaseous fuels	kg/TJ	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.4 Other Sectors][1.A.4.b Residential]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
N ₂ O											
Liquid fuels	kg/TJ	NO									
Gaseous fuels	kg/TJ	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18

Tokelau CRF Table 1.A.4.b: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.4 Other Sectors][1.A.4.b Residential] (Part 2 of 3)

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.4 Other Sectors][1.A.4.b Residential]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Fuel Consumption	TJ	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16
Liquid fuels	TJ	IE									
Gaseous fuels	TJ	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16
Calorific value	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Liquid fuels	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Gaseous fuels	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Method											
CO ₂		T1									
CH ₄		T1									
N ₂ O		T1									
Emission factor information											
CO ₂		D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
CO ₂	kt	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12
Liquid fuels	kt	IE									
Gaseous fuels	kt	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12
CH ₄	kt	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204
Liquid fuels	kt	IE									
Gaseous fuels	kt	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204

[1. Energy][1.AA Fuel Combustion – Sectoral approach][1.A.4 Other Sectors][1.A.4.b Residential]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
N ₂ O	kt	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004
Liquid fuels	kt	IE									
Gaseous fuels	kt	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004
No _x	kt	NE									
CO	kt	NE									
NMVOG	kt	NE									
SO ₂	kt	NE									
Amount captured											
CO ₂	kt	NO									
Liquid fuels	kt	NO									
Gaseous fuels	kt	NO									
Implied emission factor											
CO ₂											
Liquid fuels	t/TJ	NO									
Gaseous fuels	t/TJ	56.79	56.79	56.79	56.79	56.79	56.79	56.79	56.79	56.79	56.79
CH ₄											
Liquid fuels	kg/TJ	NO									
Gaseous fuels	kg/TJ	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8
N ₂ O											
Liquid fuels	kg/TJ	NO									
Gaseous fuels	kg/TJ	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18

Tokelau CRF Table 1.A.4.b: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.4 Other Sectors][1.A.4.b Residential] (Part 3 of 3)

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.4 Other Sectors][1.A.4.b Residential]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Fuel Consumption	TJ	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16	1.25	1.76	1.66
Liquid fuels	TJ	IE											
Gaseous fuels	TJ	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16	2.16	1.25	1.76	1.66
Calorific value	GCV												

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.4 Other Sectors][1.A.4.b Residential]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Liquid fuels		GCV											
Gaseous fuels		GCV											
Method													
CO ₂		T1											
CH ₄		T1											
N ₂ O		T1											
Emission factor information													
CO ₂		D	D	D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D	D	D
Emissions													
CO ₂	kt	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.07	0.1	0.09
Liquid fuels	kt	IE											
Gaseous fuels	kt	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.12	0.07	0.1	0.09
CH ₄	kt	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0000698	0.0000983	0.0000929
Liquid fuels	kt	IE											
Gaseous fuels	kt	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0001204	0.0000698	0.0000983	0.0000929
N ₂ O	kt	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000002	0.0000003	0.0000003
Liquid fuels	kt	IE											
Gaseous fuels	kt	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000004	0.0000002	0.0000003	0.0000003
No _x	kt	NE											
CO	kt	NE											
NMVOC	kt	NE											
SO ₂	kt	NE											
Amount captured													
CO ₂	kt	NO											
Liquid fuels	kt	NO											
Gaseous fuels	kt	NO											

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.4 Other Sectors][1.A.4.b Residential]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Implied emission factor													
CO ₂													
Liquid fuels	t/TJ	NO											
Gaseous fuels	t/TJ	56.79	56.79	56.79	56.79	56.79	56.79	56.79	56.79	56.79	56.79	56.79	56.79
CH ₄													
Liquid fuels	kg/TJ	NO											
Gaseous fuels	kg/TJ	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8	55.8
N ₂ O													
Liquid fuels	kg/TJ	NO											
Gaseous fuels	kg/TJ	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18

Tokelau CRF Table 1.A.4.c.iii Gas/Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.4 Other Sectors][1.A.4.c Agriculture/Forestry/Fishing][1.A.4.c.iii Fishing] [Gas/Diesel Oil] (Part 1 of 3)

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.4 Other Sectors][1.A.4.c Agriculture/Forestry/Fishing][1.A.4.c.iii Fishing][Gas/Diesel Oil]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Fuel consumption	TJ	IE									
Calorific value		GCV									
Method											
CO ₂		T1									
CH ₄		T1									
N ₂ O		T1									
Emission factor information											
CO ₂		D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.4 Other Sectors][1.A.4.c Agriculture/Forestry/Fishing][1.A.4.c.iii Fishing][Gas/Diesel Oil]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Emissions											
CO ₂	kt	IE									
CH ₄	kt	IE									
N ₂ O	kt	IE									
Implied emission factor											
CO ₂	t/TJ	NA									
CH ₄	kg/TJ	NA									
N ₂ O	kg/TJ	NA									

Tokelau CRF Table 1.A.4.c.iii Gas/Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.4 Other Sectors][1.A.4.c Agriculture/Forestry/Fishing][1.A.4.c.iii Fishing] [Gas/Diesel Oil] (Part 2 of 3)

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.4 Other Sectors][1.A.4.c Agriculture/Forestry/Fishing][1.A.4.c.iii Fishing][Gas/Diesel Oil]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Fuel consumption	TJ	IE									
Calorific value		GCV									
Method											
CO ₂		T1									
CH ₄		T1									
N ₂ O		T1									
Emission factor information											
CO ₂		D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
CO ₂	kt	IE									
CH ₄	kt	IE									
N ₂ O	kt	IE									

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.4 Other Sectors][1.A.4.c Agriculture/Forestry/Fishing][1.A.4.c.iii Fishing][Gas/Diesel Oil]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Implied emission factor											
CO ₂	t/TJ	NA									
CH ₄	kg/TJ	NA									
N ₂ O	kg/TJ	NA									

Tokelau CRF Table 1.A.4.c.iii Gas/Diesel Oil: [1. Energy][1.AA Fuel Combustion – Sectoral approach] [1.A.4 Other Sectors][1.A.4.c Agriculture/Forestry/Fishing][1.A.4.c.iii Fishing] [Gas/Diesel Oil] (Part 3 of 3)

[1. Energy][1.AA Fuel Combustion - Sectoral approach][1.A.4 Other Sectors][1.A.4.c Agriculture/Forestry/Fishing][1.A.4.c.iii Fishing][Gas/Diesel Oil]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Fuel consumption	TJ	IE											
Calorific value	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Method													
CO ₂		T1											
CH ₄		T1											
N ₂ O		T1											
Emission factor information													
CO ₂		D	D	D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D	D	D
Emissions													
CO ₂	kt	IE											
CH ₄	kt	IE											
N ₂ O	kt	IE											
Implied emission factor													
CO ₂	t/TJ	NA											
CH ₄	kg/TJ	NA											
N ₂ O	kg/TJ	NA											

Tokelau CRF Table 1.AB Gasoline: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Gasoline] (Part 1 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Gasoline]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Imports	PJ	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Exports	PJ	NO									
International bunkers	PJ	NO									
Stock change	PJ	NO									
Apparent consumption	PJ	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value		GCV									
Apparent consumption	TJ	9.8	9.8	9.8	9.8	9.8	9.8	9.8	9.8	9.8	9.8
Emission factor											
C	t/TJ	17.95	17.95	17.95	17.95	17.95	17.95	17.95	17.95	17.95	17.95
Carbon content											
C	kt	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18
Carbon stored											
C	kt	NO									
Net carbon emissions											
C	kt	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18
Fraction of carbon oxidized		1	1	1	1	1	1	1	1	1	1
Emissions											
C	kt	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18
CO ₂	kt	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65

Tokelau CRF Table 1.AB Gasoline: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Gasoline] (Part 2 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Gasoline]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Imports	PJ	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Exports	PJ	NO									
International bunkers	PJ	NO									
Stock change	PJ	NO									
Apparent consumption	PJ	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Gasoline]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value		GCV									
Apparent consumption	TJ	9.8	9.8	9.8	9.8	9.8	9.8	9.8	9.8	9.8	9.8
Emission factor											
C	t/TJ	17.95	17.95	17.95	17.95	17.95	17.95	17.95	17.95	17.95	17.95
Carbon content											
C	kt	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18
Carbon stored											
C	kt	NO									
Net carbon emissions											
C	kt	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18
Fraction of carbon oxidized		1	1	1	1	1	1	1	1	1	1
Emissions											
C	kt	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18
CO ₂	kt	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65

Tokelau CRF Table 1.AB Gasoline: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Gasoline] (Part 3 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Gasoline]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Imports	PJ	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Exports	PJ	NO											
International bunkers	PJ	NO											
Stock change	PJ	NO											
Apparent consumption	PJ	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value		GCV											
Apparent consumption	TJ	9.8	9.8	9.8	9.8	9.8	9.8	9.8	9.8	9.8	9.39	11.26	10.69
Emission factor													
C	t/TJ	17.95	17.95	17.95	17.95	17.95	17.95	17.95	17.95	17.95	17.95	17.95	17.95
Carbon content													
C	kt	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.17	0.2	0.19

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Gasoline]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Carbon stored													
C	kt	NO											
Net carbon emissions													
C	kt	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.17	0.2	0.19
Fraction of carbon oxidized		1	1	1	1	1	1	1	1	1	1	1	1
Emissions													
C	kt	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.18	0.17	0.2	0.19
CO ₂	kt	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.65	0.62	0.74	0.7

Tokelau CRF Table 1.AB Gas Diesel Oil: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Gas / Diesel Oil] (Part 1 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Gas / Diesel Oil]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Imports	PJ	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Exports	PJ	NO									
International bunkers	PJ	NO									
Stock change	PJ	NO									
Apparent consumption	PJ	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value		GCV									
Apparent consumption	TJ	5.83	6.06	6.29	6.51	6.74	6.96	7.19	7.41	7.64	7.86
Emission factor											
C	t/TJ	19.19	19.19	19.19	19.19	19.19	19.19	19.19	19.19	19.19	19.19
Carbon content											
C	kt	0.11	0.12	0.12	0.12	0.13	0.13	0.14	0.14	0.15	0.15
Carbon stored											
C	kt	NO									
Net carbon emissions											
C	kt	0.11	0.12	0.12	0.12	0.13	0.13	0.14	0.14	0.15	0.15
Fraction of carbon oxidized		1	1	1	1	1	1	1	1	1	1
Emissions											
C	kt	0.11	0.12	0.12	0.12	0.13	0.13	0.14	0.14	0.15	0.15
CO ₂	kt	0.41	0.43	0.44	0.46	0.47	0.49	0.51	0.52	0.54	0.55

Tokelau CRF Table 1.AB Gas Diesel Oil: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Gas / Diesel Oil] (Part 2 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Gas / Diesel Oil]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Imports	PJ	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.02
Exports	PJ	NO									
International bunkers	PJ	NO									
Stock change	PJ	NO									
Apparent consumption	PJ	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.02
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value		GCV									
Apparent consumption	TJ	8.09	8.32	8.54	8.77	15.53	22.29	22.52	22.74	22.97	23.19
Emission factor											
C	t/TJ	19.19	19.19	19.19	19.19	19.19	19.19	19.19	19.19	19.19	19.19
Carbon content											
C	kt	0.16	0.16	0.16	0.17	0.3	0.43	0.43	0.44	0.44	0.45
Carbon stored											
C	kt	NO									
Net carbon emissions											
C	kt	0.16	0.16	0.16	0.17	0.3	0.43	0.43	0.44	0.44	0.45
Fraction of carbon oxidized		1	1	1	1	1	1	1	1	1	1
Emissions											
C	kt	0.16	0.16	0.16	0.17	0.3	0.43	0.43	0.44	0.44	0.45
CO ₂	kt	0.57	0.59	0.6	0.62	1.09	1.57	1.58	1.6	1.62	1.63

Tokelau CRF Table 1.AB Gas Diesel Oil: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Gas / Diesel Oil] (Part 3 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Gas / Diesel Oil]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Imports	PJ	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02
Exports	PJ	NO											
International bunkers	PJ	NO											
Stock change	PJ	NO											
Apparent consumption	PJ	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Gas / Diesel Oil]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value		GCV											
Apparent consumption	TJ	23.42	23.65	20.5	10.62	10.84	10.7	11.74	12.93	14.31	24.98	20.92	16.1
Emission factor													
C	t/TJ	19.19	19.19	19.19	19.19	19.19	19.19	19.19	19.19	19.19	19.19	19.19	19.19
Carbon content													
C	kt	0.45	0.45	0.39	0.2	0.21	0.21	0.23	0.25	0.27	0.48	0.4	0.31
Carbon stored													
C	kt	NO											
Net carbon emissions													
C	kt	0.45	0.45	0.39	0.2	0.21	0.21	0.23	0.25	0.27	0.48	0.4	0.31
Fraction of carbon oxidized		1	1	1	1	1	1	1	1	1	1	1	1
Emissions													
C	kt	0.45	0.45	0.39	0.2	0.21	0.21	0.23	0.25	0.27	0.48	0.4	0.31
CO ₂	kt	1.65	1.66	1.44	0.75	0.76	0.75	0.83	0.91	1.01	1.76	1.47	1.13

Tokelau CRF Table 1.AB Other Kerosene: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Other Kerosene] (Part 1 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Other Kerosene]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Imports	PJ	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661
Exports	PJ	NO									
International bunkers	PJ	NO									
Stock change	PJ	NO									
Apparent consumption	PJ	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value		GCV									
Apparent consumption	TJ	0.67	0.67	0.67	0.67	0.67	0.67	0.67	0.67	0.67	0.67
Emission factor											
C	t/TJ	18.62	18.62	18.62	18.62	18.62	18.62	18.62	18.62	18.62	18.62

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Other Kerosene]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Carbon content											
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Carbon stored											
C	kt	NO									
Net carbon emissions											
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Fraction of carbon oxidized											
		1	1	1	1	1	1	1	1	1	1
Emissions											
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
CO ₂	kt	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05

Tokelau CRF Table 1.AB Other Kerosene: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Other Kerosene] (Part 2 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Other Kerosene]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Imports	PJ	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661
Exports	PJ	NO									
International bunkers	PJ	NO									
Stock change	PJ	NO									
Apparent consumption	PJ	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value		GCV									
Apparent consumption	TJ	0.67	0.67	0.67	0.67	0.67	0.67	0.67	0.67	0.67	0.67
Emission factor											
C	t/TJ	18.62	18.62	18.62	18.62	18.62	18.62	18.62	18.62	18.62	18.62
Carbon content											
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Carbon stored											
C	kt	NO									
Net carbon emissions											
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01

[1. Energy][1.AB Fuel Combustion - Reference Approach] [Liquid Fuels][Other Kerosene]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Fraction of carbon oxidized		1	1	1	1	1	1	1	1	1	1
Emissions											
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
CO ₂	kt	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05

Tokelau CRF Table 1.AB Other Kerosene: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels][Other Kerosene] (Part 3 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach] [Liquid Fuels] [Other Kerosene]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Imports	PJ	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0002342	0.0000952	0.000205
Exports	PJ	NO	NO										
International bunkers	PJ	NO	NO										
Stock change	PJ	NO	NO										
Apparent consumption	PJ	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0006661	0.0002342	0.0000952	0.000205
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Apparent consumption	TJ	0.67	0.67	0.67	0.67	0.67	0.67	0.67	0.67	0.67	0.23	0.1	0.2
Emission factor													
C	t/TJ	18.62	18.62	18.62	18.62	18.62	18.62	18.62	18.62	18.62	18.62	18.62	18.62
Carbon content													
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0	0	0
Carbon stored													
C	kt	NO	NO										
Net carbon emissions													
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0	0	0
Fraction of carbon oxidized		1	1	1	1	1	1	1	1	1	1	1	1
Emissions													
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0	0	0
CO ₂	kt	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.02	0.01	0.01

Tokelau CRF Table 1.AB LPG: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels] [Liquefied Petroleum Gases (LPG)] (Part 1 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Liquefied Petroleum Gases (LPG)]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Imports	PJ	0	0	0	0	0	0	0	0	0	0
Exports	PJ	NO									
International bunkers	PJ	NO									
Stock change	PJ	NO									
Apparent consumption	PJ	0	0	0	0	0	0	0	0	0	0
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value		GCV									
Apparent consumption	TJ	1.49	1.49	1.49	1.49	1.49	1.49	1.49	1.49	1.49	1.49
Emission factor											
C	t/TJ	15.48	15.48	15.48	15.48	15.48	15.48	15.48	15.48	15.48	15.48
Carbon content											
C	kt	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Carbon stored											
C	kt	NO									
Net carbon emissions											
C	kt	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Fraction of carbon oxidized		1	1	1	1	1	1	1	1	1	1
Emissions											
C	kt	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
CO ₂	kt	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08

Tokelau CRF Table 1.AB LPG: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels] [Liquefied Petroleum Gases (LPG)] (Part 2 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Liquefied Petroleum Gases (LPG)]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Imports	PJ	0	0	0	0	0	0	0	0	0	0
Exports	PJ	NO									
International bunkers	PJ	NO									
Stock change	PJ	NO									
Apparent consumption	PJ	0	0	0	0	0	0	0	0	0	0

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Liquefied Petroleum Gases (LPG)]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value		GCV									
Apparent consumption	TJ	1.49	1.49	1.49	1.49	1.49	1.49	1.49	1.49	1.49	1.49
Emission factor											
C	t/TJ	15.48	15.48	15.48	15.48	15.48	15.48	15.48	15.48	15.48	15.48
Carbon content											
C	kt	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Carbon stored											
C	kt	NO									
Net carbon emissions											
C	kt	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Fraction of carbon oxidized		1	1	1	1	1	1	1	1	1	1
Emissions											
C	kt	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
CO ₂	kt	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08

Tokelau CRF Table 1.AB LPG: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels] [Liquefied Petroleum Gases (LPG)] (Part 3 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Liquefied Petroleum Gases (LPG)]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Imports	PJ	0	0	0	0	0	0	0	0	0	0	0	0
Exports	PJ	NO											
International bunkers	PJ	NO											
Stock change	PJ	NO											
Apparent consumption	PJ	0	0	0	0	0	0	0	0	0	0	0	0
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value		GCV											
Apparent consumption	TJ	1.49	1.49	1.49	1.49	1.49	1.49	1.49	1.49	1.49	1.02	1.67	1.46
Emission factor													
C	t/TJ	15.48	15.48	15.48	15.48	15.48	15.48	15.48	15.48	15.48	15.48	15.48	15.48

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Liquefied Petroleum Gases (LPG)]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Carbon content													
C	kt	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.02
Carbon stored													
C	kt	NO											
Net carbon emissions													
C	kt	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.02
Fraction of carbon oxidized													
		1	1	1	1	1	1	1	1	1	1	1	1
Emissions													
C	kt	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.02
CO ₂	kt	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.06	0.09	0.08

Tokelau CRF Table 1.AB Lubricants: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels] [Lubricants] (Part 1 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Lubricants]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Imports	PJ	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901
Exports	PJ	NO									
International bunkers	PJ	NO									
Stock change	PJ	NO									
Apparent consumption	PJ	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV	GCV
Apparent consumption	TJ	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39
Emission factor											
C	t/TJ	19	19	19	19	19	19	19	19	19	19
Carbon content											
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Carbon stored											
C	kt	NO									

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Lubricants]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Net carbon emissions											
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Fraction of carbon oxidized		1	1	1	1	1	1	1	1	1	1
Emissions											
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
CO ₂	kt	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03

Tokelau CRF Table 1.AB Lubricants: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels] [Lubricants] (Part 2 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Lubricants]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Imports	PJ	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901
Exports	PJ	NO									
International bunkers	PJ	NO									
Stock change	PJ	NO									
Apparent consumption	PJ	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value		GCV									
Apparent consumption	TJ	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39
Emission factor											
C	t/TJ	19	19	19	19	19	19	19	19	19	19
Carbon content											
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Carbon stored											
C	kt	NO									
Net carbon emissions											
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Fraction of carbon oxidized		1	1	1	1	1	1	1	1	1	1
Emissions											
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
CO ₂	kt	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03

Tokelau CRF Table 1.AB Lubricants: [1. Energy][1.AB Fuel Combustion – Reference Approach] [Liquid Fuels] [Lubricants] (Part 3 of 3)

[1. Energy][1.AB Fuel Combustion - Reference Approach][Liquid Fuels][Lubricants]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Imports	PJ	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0001519	0.0002028	0.0001007
Exports	PJ	NO											
International bunkers	PJ	NO											
Stock change	PJ	NO											
Apparent consumption	PJ	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0003901	0.0001519	0.0002028	0.0001007
Conversion factor	TJ/unit	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000	1000
Calorific value		GCV											
Apparent consumption	TJ	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.39	0.15	0.2	0.1
Emission factor													
C	t/TJ	19	19	19	19	19	19	19	19	19	19	19	19
Carbon content													
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0	0	0
Carbon stored													
C	kt	NO											
Net carbon emissions													
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0	0	0
Fraction of carbon oxidized		1	1	1	1	1	1	1	1	1	1	1	1
Emissions													
C	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0	0	0
CO ₂	kt	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.01	0.01	0.01

Tokelau CRF Table 2.F.1.b HFC-134a Product Uses as Substitutes for ODS: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.b Domestic Refrigeration] [HFC-134a] (Part 1 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.b Domestic Refrigeration][HFC-134a]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Amount											
Filled into new manufactured products	t	NO									
In operating systems (average annual stocks)	t	NO	NO	NO	NO	0.02	0.04	0.07	0.09	0.11	0.13
Remaining in products at decommissioning	t	NO									
Emissions	t	NO	NO	NO	NO	0	0.01	0.01	0.01	0.02	0.02
From manufacturing	t	NO									
From stocks	t	NO	NO	NO	NO	0	0.01	0.01	0.01	0.02	0.02
From disposal	t	NO									
Recovery	t	NO									
Implied emission factor											
Product manufacturing factor	%	NO									
Product life factor	%	NO	NO	NO	NO	15	15	15	15	15	15
Disposal loss factor	%	NO									

Tokelau CRF Table 2.F.1.b HFC-134a Product Uses as Substitutes for ODS: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.b Domestic Refrigeration] [HFC-134a] (Part 2 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.b Domestic Refrigeration][HFC-134a]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Amount											
Filled into new manufactured products	t	NO									
In operating systems (average annual stocks)	t	0.14	0.16	0.2	0.25	0.27	0.29	0.32	0.32	0.31	0.31
Remaining in products at decommissioning	t	NO									
Emissions	t	0.02	0.02	0.03	0.04	0.04	0.04	0.05	0.05	0.05	0.05
From manufacturing	t	NO									
From stocks	t	0.02	0.02	0.03	0.04	0.04	0.04	0.05	0.05	0.05	0.05
From disposal	t	NO									
Recovery	t	NO									

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.b Domestic Refrigeration][HFC-134a]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Implied emission factor											
Product manufacturing factor	%	NO									
Product life factor	%	15	15	15	15	15	15	15	15	15	15
Disposal loss factor	%	NO									

Tokelau CRF Table 2.F.1.b HFC-134a Product Uses as Substitutes for ODS: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.b Domestic Refrigeration] [HFC-134a] (Part 3 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.b Domestic Refrigeration][HFC-134a]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Amount													
Filled into new manufactured products	t	NO											
In operating systems (average annual stocks)	t	0.31	0.31	0.29	0.27	0.25	0.23	0.21	0.21	0.21	0.21	0.21	0.21
Remaining in products at decommissioning	t	NO											
Emissions	t	0.05	0.05	0.04	0.04	0.04	0.03	0.03	0.03	0.03	0.03	0.03	0.03
From manufacturing	t	NO											
From stocks	t	0.05	0.05	0.04	0.04	0.04	0.03	0.03	0.03	0.03	0.03	0.03	0.03
From disposal	t	NO											
Recovery	t	NO											
Implied emission factor													
Product manufacturing factor	%	NO											
Product life factor	%	15	15	15	15	15	15	15	15	15	15	15	15
Disposal loss factor	%	NO											

**Tokelau CRF Table 2.F.1.f: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning]
[2.F.1.f Stationary Air Conditioning] (Part 1 of 3)**

[2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Method											
HFCs		NA									
PFCs		NA									
Unspecified mix of HFCs and PFCs		NA									
SF ₆		NA									
NF ₃		NA									
Emission factor information											
HFCs		NA									
Emissions											
HFCs	t CO ₂ -e	NO									
HFC-32	t	NO									
HFC-125	t	NO									
HFC-134a	t	NO									
HFCs and PFCs	t CO ₂ -e	NO									
Recovery											
Aggregate F-gases	t CO ₂ -e	NO									

**Tokelau CRF Table 2.F.1.f: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning]
[2.F.1.f Stationary Air Conditioning] (Part 2 of 3)**

[2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Method											
HFCs		NA	T1a	T1a	T1a						
PFCs		NA									
Unspecified mix of HFCs and PFCs		NA									
SF ₆		NA									
NF ₃		NA									

[2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Emission factor information											
HFCs		NA	D	D	D						
Emissions											
HFCs	t CO ₂ -e	NO	17.06	34.12	51.17						
HFC-32	t	NO	0	0.01	0.01						
HFC-125	t	NO	0	0.01	0.01						
HFC-134a	t	NO	0.000285	0.00057	0.000855						
HFCs and PFCs	t CO ₂ -e	NO	17.06	34.12	51.17						
Recovery											
Aggregate F-gases	t CO ₂ -e	NO	17.06	34.12	51.17						

Tokelau CRF Table 2.F.1.f: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air Conditioning] (Part 3 of 3)

[2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Method													
HFCs		T1a	T1a	T1a	T1a	T1a	T1a	T1a	T1a	T1a	T1a	T1a	T1a
PFCs		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Unspecified mix of HFCs and PFCs		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
SF ₆		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
NF ₃		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Emission factor information													
HFCs		D	D	D	D	D	D	D	D	D	D	D	D
Emissions													
HFCs	t CO ₂ -e	68.23	85.29	102.35	119.41	136.46	153.52	170.58	170.58	170.58	170.58	170.58	170.58
HFC-32	t	0.01	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03	0.03
HFC-125	t	0.02	0.02	0.03	0.03	0.03	0.04	0.04	0.04	0.04	0.04	0.04	0.04
HFC-134a	t	0	0	0	0	0	0	0	0	0	0	0	0
HFCs and PFCs	t CO ₂ -e	68.23	85.29	102.35	119.41	136.46	153.52	170.58	170.58	170.58	170.58	170.58	170.58

[2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Recovery													
Aggregate F-gases	t CO ₂ -e	68.23	85.29	102.35	119.41	136.46	153.52	170.58	170.58	170.58	170.58	170.58	170.58

Tokelau CRF Table 2.F.1.f HFC-32: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-32] (Part 1 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning][HFC-32]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Amount											
Filled into new manufactured products	t	NO									
In operating systems (average annual stocks)	t	NO									
Remaining in products at decommissioning	t	NO									
Emissions	t	NO									
From manufacturing	t	NO									
From stocks	t	NO									
From disposal	t	NO									
Recovery	t	NO									
Implied emission factor											
Product manufacturing factor	%	NO									
Product life factor	%	NO									
Disposal loss factor	%	NO									

Tokelau CRF Table 2.F.1.f HFC-32: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-32] (Part 2 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning][HFC-32]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Amount											
Filled into new manufactured products	t	NO									
In operating systems (average annual stocks)	t	NO	0.02	0.04	0.06						
Remaining in products at decommissioning	t	NO									
Emissions	t	NO	0	0.01	0.01						
From manufacturing	t	NO									
From stocks	t	NO	0	0.01	0.01						
From disposal	t	NO									
Recovery	t	NO									
Implied emission factor											
Product manufacturing factor	%	NO									
Product life factor	%	NO	15	15	15						
Disposal loss factor	%	NO									

Tokelau CRF Table 2.F.1.f HFC-32: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-32] (Part 3 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning][HFC-32]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Amount													
Filled into new manufactured products	t	NO											
In operating systems (average annual stocks)	t	0.08	0.1	0.12	0.14	0.16	0.18	0.2	0.2	0.2	0.2	0.2	0.2
Remaining in products at decommissioning	t	NO											
Emissions	t	0.01	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03	0.03
From manufacturing	t	NO											
From stocks	t	0.01	0.02	0.02	0.02	0.02	0.03	0.03	0.03	0.03	0.03	0.03	0.03
From disposal	t	NO											
Recovery	t	NO											

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning][HFC-32]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Implied emission factor													
Product manufacturing factor	%	NO											
Product life factor	%	15	15	15	15	15	15	15	15	15	15	15	15
Disposal loss factor	%	NO											

Tokelau CRF Table 2.F.1.f HFC-125: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-125] (Part 1 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning][HFC-125]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Amount											
Filled into new manufactured products	t	NO									
In operating systems (average annual stocks)	t	NO									
Remaining in products at decommissioning	t	NO									
Emissions	t	NO									
From manufacturing	t	NO									
From stocks	t	NO									
From disposal	t	NO									
Recovery	t	NO									
Implied emission factor											
Product manufacturing factor	%	NO									
Product life factor	%	NO									
Disposal loss factor	%	NO									

Tokelau CRF Table 2.F.1.f HFC-125: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-125] (Part 2 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning][HFC-125]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Amount											
Filled into new manufactured products	t	NO									
In operating systems (average annual stocks)	t	NO	0.03	0.06	0.08						
Remaining in products at decommissioning	t	NO									
Emissions	t	NO	0	0.01	0.01						
From manufacturing	t	NO									
From stocks	t	NO	0	0.01	0.01						
From disposal	t	NO									
Recovery	t	NO									
Implied emission factor											
Product manufacturing factor	%	NO									
Product life factor	%	NO	15	15	15						
Disposal loss factor	%	NO									

Tokelau CRF Table 2.F.1.f HFC-125: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-125] (Part 3 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning][HFC-125]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Amount													
Filled into new manufactured products	t	NO											
In operating systems (average annual stocks)	t	0.11	0.14	0.17	0.19	0.22	0.25	0.28	0.28	0.28	0.28	0.28	0.28
Remaining in products at decommissioning	t	NO											
Emissions	t	0.02	0.02	0.03	0.03	0.03	0.04	0.04	0.04	0.04	0.04	0.04	0.04
From manufacturing	t	NO											
From stocks	t	0.02	0.02	0.03	0.03	0.03	0.04	0.04	0.04	0.04	0.04	0.04	0.04
From disposal	t	NO											
Recovery	t	NO											

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning][HFC-125]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Implied emission factor													
Product manufacturing factor	%	NO											
Product life factor	%	15	15	15	15	15	15	15	15	15	15	15	15
Disposal loss factor	%	NO											

Tokelau CRF Table 2.F.1.f HFC-134a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-134a] (Part 1 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning][HFC-134a]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Amount											
Filled into new manufactured products	t	NO									
In operating systems (average annual stocks)	t	NO									
Remaining in products at decommissioning	t	NO									
Emissions	t	NO									
From manufacturing	t	NO									
From stocks	t	NO									
From disposal	t	NO									
Recovery	t	NO									
Implied emission factor											
Product manufacturing factor	%	NO									
Product life factor	%	NO									
Disposal loss factor	%	NO									

Tokelau CRF Table 2.F.1.f HFC-134a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-134a] (Part 2 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning][HFC-134a]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Amount											
Filled into new manufactured products	t	NO	NO	NO							
In operating systems (average annual stocks)	t	NO	0	0	0.01						
Remaining in products at decommissioning	t	NO	NO	NO							
Emissions	t	NO	0.000285	0.00057	0.000855						
From manufacturing	t	NO	NO	NO							
From stocks	t	NO	0.000285	0.00057	0.000855						
From disposal	t	NO	NO	NO							
Recovery	t	NO	NO	NO							
Implied emission factor											
Product manufacturing factor	%	NO	NO	NO							
Product life factor	%	NO	15	15	15						
Disposal loss factor	%	NO	NO	NO							

Tokelau CRF Table 2.F.1.f HFC-134a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.1 Refrigeration and Air conditioning] [2.F.1.f Stationary Air conditioning] [HFC-134a] (Part 3 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning][HFC-134a]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Amount													
Filled into new manufactured products	t	NO											
In operating systems (average annual stocks)	t	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Remaining in products at decommissioning	t	NO											
Emissions	t	0	0	0	0	0	0	0	0	0	0	0	0
From manufacturing	t	NO											
From stocks	t	0	0	0	0	0	0	0	0	0	0	0	0
From disposal	t	NO											
Recovery	t	NO											

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.1 Refrigeration and Air conditioning][2.F.1.f Stationary Air-Conditioning][HFC-134a]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Implied emission factor													
Product manufacturing factor	%	NO											
Product life factor	%	15	15	15	15	15	15	15	15	15	15	15	15
Disposal loss factor	%	NO											

Tokelau CRF Table 2.F.4.a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] (Part 1 of 3)

[2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.4 Aerosols][2.F.4.a Metered Dose Inhalers]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Method											
HFCs		NA	NA	NA	NA	NA	T1a	T1a	T1a	T1a	T1a
PFCs		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Unspecified mix of HFCs and PFCs		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
SF ₆		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
NF ₃		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Emission factor information											
HFCs		NA	NA	NA	NA	NA	D	D	D	D	D
PFCs		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Unspecified mix of HFCs and PFCs		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
SF ₆		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
NF ₃		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Emissions	kt CO ₂ -equivalent	NO	NO	NO	NO	NO	0.0001188	0.0006994	0	0	0
HFCs	t CO ₂ -equivalent	NO	NO	NO	NO	NO	0.12	0.7	1.14	1.69	2.44
HFC-134a	t	NO	NO	NO	NO	NO	0.0000831	0.0004891	0.0008003	0	0
HFC-227ea	t	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Aggregate F-gases	t CO ₂ -equivalent	NO	NO	NO	NO	NO	0.12	0.7	1.14	1.69	2.44

Tokelau CRF Table 2.F.4.a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] (Part 2 of 3)

[2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Method											
HFCs		T1a	T1a	T1a	T1a	T1a	T1a	T1a	T1a	T1a	T1a
PFCs		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Unspecified mix of HFCs and PFCs		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
SF ₆		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
NF ₃		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Emission factor information											
HFCs		D	D	D	D	D	D	D	D	D	D
PFCs		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Unspecified mix of HFCs and PFCs		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
SF ₆		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
NF ₃		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Emissions	kt CO ₂ -equivalent	0	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
HFCs	t CO ₂ -equivalent	2.85	6.09	11.54	13.9	13.66	13.66	13.35	13.62	14.18	14.76
HFC-134a	t	0	0	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
HFC-227ea	t	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Aggregate F-gases	t CO ₂ -equivalent	2.85	6.09	11.54	13.9	13.66	13.66	13.35	13.62	14.18	14.76

Tokelau CRF Table 2.F.4.a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] (Part 3 of 3)

[2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.4 Aerosols][2.F.4.a Metered Dose Inhalers]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Method													
HFCs		T1a											
PFCs		NA											
Unspecified mix of HFCs and PFCs		NA											

[2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.4 Aerosols][2.F.4.a Metered Dose Inhalers]													
	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
SF ₆		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
NF ₃		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Emission factor information													
HFCs		D	D	D	D	D	D	D	D	D	D	D	D
PFCs		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Unspecified mix of HFCs and PFCs		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
SF ₆		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
NF ₃		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Emissions	kt CO ₂ -equivalent	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
HFCs	t CO ₂ -equivalent	15.46	17.22	18.93	19.09	19.1	19.4	19.42	19.12	18.69	18.27	18.27	18.27
HFC-134a	t	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
HFC-227ea	t	NO	0.0003553	0.0007195	0.0007207	0.0007209	0.0007322	0.0007135	0.0006669	0.0006159	0.0005628	0.0005628	0.0005628
Aggregate F-gases	t CO ₂ -equivalent	15.46	17.22	18.93	19.09	19.1	19.4	19.42	19.12	18.69	18.27	18.27	18.27

Tokelau CRF Table 2.F.4.a HFC-134a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] [HFC-134a] (Part 1 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.4 Aerosols][2.F.4.a Metered Dose Inhalers][HFC-134a]												
	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	
Amount												
Filled into new manufactured products	t	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	
In operating systems (average annual stocks)	t	NO	NO	NO	NO	NO	0.0000831	0.0004891	0.0008003	0	0	
Emissions	t	NO	NO	NO	NO	NO	0.0000831	0.0004891	0.0008003	0	0	
From manufacturing	t	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	
From stocks	t	NO	NO	NO	NO	NO	0.0000831	0.0004891	0.0008003	0	0	
Recovery	t	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.4 Aerosols][2.F.4.a Metered Dose Inhalers][HFC-134a]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Implied emission factor											
Product manufacturing factor	%	NO									
Product life factor	%	NO	NO	NO	NO	NO	100	100	100	100	100

Tokelau CRF Table 2.F.4.a HFC-134a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] [HFC-134a] (Part 2 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.4 Aerosols][2.F.4.a Metered Dose Inhalers][HFC-134a]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Amount											
Filled into new manufactured products	t	NO									
In operating systems (average annual stocks)	t	0	0	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Emissions											
From manufacturing	t	NO									
From stocks	t	0	0	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Recovery	t	NO									
Implied emission factor											
Product manufacturing factor	%	NO									
Product life factor	%	100	100	100	100	100	100	100	100	100	100

Tokelau CRF Table 2.F.4.a HFC-134a: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] [HFC-134a] (Part 3 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.4 Aerosols][2.F.4.a Metered Dose Inhalers][HFC-134a]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Amount													
Filled into new manufactured products	t	NO											
In operating systems (average annual stocks)	t	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Emissions													
From manufacturing	t	NO											
From stocks	t	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Recovery	t	NO											

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.4 Aerosols][2.F.4.a Metered Dose Inhalers][HFC-134a]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Implied emission factor													
Product manufacturing factor	%	NO											
Product life factor	%	100	100	100	100	100	100	100	100	100	100	100	100

Tokelau CRF Table 2.F.4.a HFC-227ea: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] [HFC-227ea] (Part 1 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.4 Aerosols][2.F.4.a Metered Dose Inhalers][HFC-227ea]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Amount											
Filled into new manufactured products	t	NO									
In operating systems (average annual stocks)	t	NO									
Emissions											
From manufacturing	t	NO									
From stocks	t	NO									
Recovery	t	NO									
Implied emission factor											
Product manufacturing factor	%	NO									
Product life factor	%	NO									

Tokelau CRF Table 2.F.4.a HFC-227ea: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] [HFC-227ea] (Part 2 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.4 Aerosols][2.F.4.a Metered Dose Inhalers][HFC-227ea]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Amount											
Filled into new manufactured products	t	NO									
In operating systems (average annual stocks)	t	NO									
Emissions											
From manufacturing	t	NO									
From stocks	t	NO									
Recovery	t	NO									

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.4 Aerosols][2.F.4.a Metered Dose Inhalers][HFC-227ea]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Implied emission factor											
Product manufacturing factor	%	NO									
Product life factor	%	NO									

Tokelau CRF Table 2.F.4.a HFC-227ea: [2. Industrial Processes and Product Use] [2.F Product Uses as Substitutes for ODS] [2.F.4 Aerosols] [2.F.4.a Metered Dose Inhalers] [HFC-227ea] (Part 3 of 3)

[Sectors/Totals][2. Industrial Processes and Product Use][2.F Product Uses as Substitutes for ODS][2.F.4 Aerosols][2.F.4.a Metered Dose Inhalers][HFC-227ea]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Amount													
Filled into new manufactured products	t	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
In operating systems (average annual stocks)	t	NO	0.0003553	0.0007195	0.0007207	0.0007209	0.0007322	0.0007135	0.0006669	0.0006159	0.0005628	0.0005628	0.0005628
Emissions	t	NO	0.0003553	0.0007195	0.0007207	0.0007209	0.0007322	0.0007135	0.0006669	0.0006159	0.0005628	0.0005628	0.0005628
From manufacturing	t	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
From stocks	t	NO	0.0003553	0.0007195	0.0007207	0.0007209	0.0007322	0.0007135	0.0006669	0.0006159	0.0005628	0.0005628	0.0005628
Recovery	t	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Implied emission factor													
Product manufacturing factor	%	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Product life factor	%	NO	100	100	100	100	100	100	100	100	100	100	100

Tokelau CRF Table 2.G.3.a: [2. Industrial Processes and Product Use] [2.G Other Product Manufacture and Use] [2.G.3 N2O from Product Uses] [2.G.3.a Medical Applications] (Part 1 of 3)

[2. Industrial Processes and Product Use][2.G Other Product Manufacture and Use][2.G.3 N2O from Product Uses][2.G.3.a Medical Applications]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Activity data											
N ₂ O	kt	0.000154	0.0001391	0.0001303	0.0001217	0.0001136	0.0001058	0.0000986	0.000092	0.0000861	0.0000807
Method											
N ₂ O		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1

[2. Industrial Processes and Product Use][2.G Other Product Manufacture and Use][2.G.3 N2O from Product Uses][2.G.3.a Medical Applications]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Emission factor information											
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
N ₂ O	kt	0.000158	0.0001428	0.0001337	0.0001249	0.0001166	0.0001086	0.0001012	0.0000944	0.0000884	0.0000828
Recovery											
N ₂ O	kt	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Implied emission factor											
N ₂ O	t/t	1.03	1.03	1.03	1.03	1.03	1.03	1.03	1.03	1.03	1.03

Tokelau CRF Table 2.G.3.a: [2. Industrial Processes and Product Use] [2.G Other Product Manufacture and Use] [2.G.3 N2O from Product Uses] [2.G.3.a Medical Applications] (Part 2 of 3)

[2. Industrial Processes and Product Use][2.G Other Product Manufacture and Use][2.G.3 N2O from Product Uses][2.G.3.a Medical Applications]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Activity data											
N ₂ O	kt	0.0000756	0.0000705	0.000063	0.0000553	0.0000482	0.0000416	0.0000355	0.0000454	0.0000515	0.0000462
Method											
N ₂ O		T1									
Emission factor information											
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
N ₂ O	kt	0.0000776	0.0000724	0.0000647	0.0000573	0.0000502	0.0000435	0.0000373	0.0000405	0.0000485	0.0000488
Recovery											
N ₂ O	kt	NO									
Implied emission factor											
N ₂ O	t/t	1.03	1.03	1.03	1.04	1.04	1.05	1.05	0.89	0.94	1.06

Tokelau CRF Table 2.G.3.a: [2. Industrial Processes and Product Use] [2.G Other Product Manufacture and Use] [2.G.3 N2O from Product Uses] [2.G.3.a Medical Applications] (Part 3 of 3)

[2. Industrial Processes and Product Use] [2.G Other Product Manufacture and Use] [2.G.3 N2O from Product Uses] [2.G.3.a Medical Applications]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Activity data													
N ₂ O	kt	0.0000519	0.000046	0.0000566	0.0000542	0.0000538	0.0000564	0.0000508	0.0000599	0.000084	0.000063	0.000063	0.000063
Method													
N ₂ O		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
Emission factor information													
N ₂ O		D	D	D	D	D	D	D	D	D	D	D	D
Emissions													
N ₂ O	kt	0.000049	0.000049	0.0000515	0.0000554	0.0000538	0.0000549	0.0000534	0.0000549	0.0000715	0.0000729	0.0000729	0.0000729
Recovery													
N ₂ O	kt	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Implied emission factor													
N ₂ O	t/t	0.94	1.07	0.91	1.02	1	0.97	1.05	0.92	0.85	1.16	1.16	1.16

Tokelau CRF Table 3.A.3 Tokelau Swine: [3. Agriculture] [3.1 Livestock] [3.A Enteric Fermentation] [3.A.3 Swine] [Other (please specify)] [Tokelau_Swine] (Part 1 of 3)

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.A Enteric Fermentation][3.A.3 Swine][Other (please specify)][Pigs]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Population	1000s	2.29	2.5	2.4	2.29	2.19	2.08	1.98	2.11	2.25	2.38
Average gross energy intake	MJ/head/day	NA									
Average CH ₄ conversion rate	%	NA									
Method											
CH ₄		T1									
Emission factor information											
CH ₄		D	D	D	D	D	D	D	D	D	D
Emissions											
CH ₄	kt	0	0	0	0	0	0	0	0	0	0
Implied emission factor											
CH ₄	kg/head/year	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Additional information											

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.A Enteric Fermentation][3.A.3 Swine][Other (please specify)][Pigs]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Weight	kg	80	80	80	80	80	80	80	80	80	80
Feeding situation		Pen									
Milk yield	kg/day	NA									
Work	h/day	NO									
Pregnant	%	NA									
Digestibility of feed	%	NA									
Gross energy	MJ/day	NA									

Tokelau CRF Table 3.A.3 Tokelau Swine: [3. Agriculture] [3.1 Livestock] [3.A Enteric Fermentation] [3.A.3 Swine] [Other (please specify)] [Tokelau_Swine] (Part 2 of 3)

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.A Enteric Fermentation][3.A.3 Swine][Other (please specify)][Pigs]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Population	1000s	2.52	2.65	2.63	2.61	2.59	2.57	2.55	2.51	2.48	2.44
Average gross energy intake	MJ/head/day	NA									
Average CH ₄ conversion rate	%	NA									
Method											
CH ₄		T1									
Emission factor information											
CH ₄		D	D	D	D	D	D	D	D	D	D
Emissions											
CH ₄	kt	0	0	0	0	0	0	0	0	0	0
Implied emission factor											
CH ₄	kg/head/year	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Additional information											
Weight	kg	80	80	80	80	80	80	80	80	80	80
Feeding situation		Pen									
Milk yield	kg/day	NA									
Work	h/day	NO									
Pregnant	%	NA									
Digestibility of feed	%	NA									
Gross energy	MJ/day	NA									

Tokelau CRF Table 3.A.3 Tokelau Swine: [3. Agriculture] [3.1 Livestock] [3.A Enteric Fermentation] [3.A.3 Swine] [Other (please specify)] [Tokelau_Swine] (Part 3 of 3)

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.A Enteric Fermentation][3.A.3 Swine][Other (please specify)][Pigs]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Population	1000s	2.4	2.36	2.22	2.08	1.93	1.79	1.65	1.65	1.65	1.65	1.65	1.65
Average gross energy intake	MJ/head/day	NA											
Average CH ₄ conversion rate	%	NA											
Method													
CH ₄		T1											
Emission factor information													
CH ₄		D	D	D	D	D	D	D	D	D	D	D	D
Emissions													
CH ₄	kt	0	0	0	0	0	0	0	0	0	0	0	0
Implied emission factor													
CH ₄	kg/head/year	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Additional information													
Weight	kg	80	80	80	80	80	80	80	80	80	80	80	80
Feeding situation		Pen											
Milk yield	kg/day	NA											
Work	h/day	NO											
Pregnant	%	NA											
Digestibility of feed	%	NA											
Gross energy	MJ/day	NA											

Tokelau CRF Table 3.A.4 Tokelau Poultry: [3. Agriculture] [3.1 Livestock] [3.A Enteric Fermentation] [3.A.4 Other livestock] [Tokelau_Poultry] (Part 1 of 3)

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.A Enteric Fermentation][3.A.4 Other livestock][Poultry]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Population	1000s	3.44	3.5	3.39	3.29	3.18	3.08	2.97	2.84	2.71	2.58
Average gross energy intake	MJ/head/day	NA									
Average CH ₄ conversion rate	%	NA									
Method											
CH ₄		NA									

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.A Enteric Fermentation][3.A.4 Other livestock][Poultry]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Emission factor information											
CH ₄		NA									
Emissions											
CH ₄	kt	NE									
Implied emission factor											
CH ₄	kg/head/year	NE									
Additional information											
Weight	kg	NA									
Feeding situation		NA									
Milk yield	kg/day	NA									
Work	h/day	NA									
Pregnant	%	NA									
Digestibility of feed	%	NA									
Gross energy	kJ/day	NA									

Tokelau CRF Table 3.A.4 Tokelau Poultry: [3. Agriculture] [3.1 Livestock] [3.A Enteric Fermentation] [3.A.4 Other livestock] [Tokelau_Poultry] (Part 2 of 3)

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.A Enteric Fermentation][3.A.4 Other livestock][Poultry]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Population	1000s	2.45	2.32	2.23	2.14	2.05	1.96	1.87	1.71	1.55	1.39
Average gross energy intake	kJ/head/day	NA									
Average CH ₄ conversion rate	%	NA									
Method											
CH ₄		NA									
Emission factor information											
CH ₄		NA									
Emissions											
CH ₄	kt	NE									
Implied emission factor											
CH ₄	kg/head/year	NE									
Additional information											
Weight	kg	NA									

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.A Enteric Fermentation][3.A.4 Other livestock][Poultry]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Feeding situation		NA									
Milk yield	kg/day	NA									
Work	h/day	NA									
Pregnant	%	NA									
Digestibility of feed	%	NA									
Gross energy	MJ/day	NA									

Tokelau CRF Table 3.A.4 Tokelau Poultry: [3. Agriculture] [3.1 Livestock] [3.A Enteric Fermentation] [3.A.4 Other livestock] [Tokelau_Poultry] (Part 3 of 3)

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.A Enteric Fermentation][3.A.4 Other livestock][Poultry]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Population	1000s	1.23	1.06	0.98	0.89	0.8	0.71	0.62	0.62	0.62	0.62	0.62	0.62
Average gross energy intake	MJ/head/day	NA											
Average CH ₄ conversion rate	%	NA											
Method													
CH ₄		NA											
Emission factor information													
CH ₄		NA											
Emissions													
CH ₄	kt	NE											
Implied emission factor													
CH ₄	kg/head/year	NE											
Additional information													
Weight	kg	NA											
Feeding situation		NA											
Milk yield	kg/day	NA											
Work	h/day	NA											
Pregnant	%	NA											
Digestibility of feed	%	NA											
Gross energy	MJ/day	NA											

Tokelau CRF Table 3.B.1.3 Tokelau Swine: [3. Agriculture] [3.1 Livestock] [3.B Manure Management] [3.B.1 CH4 Emissions] [3.B.1.3 Swine] [Other (please specify)] [Pigs] (Part 1 of 3)

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.B Manure Management][3.B.1 CH4 Emissions][3.B.1.3 Swine][Other (please specify)][Pigs]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Population	1000s	2.29	2.5	2.4	2.29	2.19	2.08	1.98	2.11	2.25	2.38
Allocation by climate region											
Warm	%	100	100	100	100	100	100	100	100	100	100
Typical animal mass (average)	kg	80	80	80	80	80	80	80	80	80	80
VS daily excretion (average)	kg dm/head/day	NA									
CH ₄ producing potential (average)	m ³ /kg VS	NA									
Method											
CH ₄		T1									
Emission factor information											
CH ₄		D	D	D	D	D	D	D	D	D	D
Emissions											
CH ₄	kt	0.04	0.05	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04
Implied emission factor											
CH ₄	kg/head/year	18.5	18.5	18.5	18.5	18.5	18.5	18.5	18.5	18.5	18.5

Tokelau CRF Table 3.B.1.3 Tokelau Swine: [3. Agriculture] [3.1 Livestock] [3.B Manure Management] [3.B.1 CH4 Emissions] [3.B.1.3 Swine] [Other (please specify)] [Pigs] (Part 2 of 3)

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.B Manure Management][3.B.1 CH4 Emissions][3.B.1.3 Swine][Other (please specify)][Pigs]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Population	1000s	2.52	2.65	2.63	2.61	2.59	2.57	2.55	2.51	2.48	2.44
Allocation by climate region											
Warm	%	100	100	100	100	100	100	100	100	100	100
Typical animal mass (average)	kg	80	80	80	80	80	80	80	80	80	80
VS daily excretion (average)	kg dm/head/day	NA									
CH ₄ producing potential (average)	m ³ /kg VS	NA									
Method											
CH ₄		T1									
Emission factor information											
CH ₄		D	D	D	D	D	D	D	D	D	D

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.B Manure Management][3.B.1 CH4 Emissions][3.B.1.3 Swine][Other (please specify)][Pigs]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Emissions											
CH ₄	kt	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05
Implied emission factor											
CH ₄	kg/head/year	18.5	18.5	18.5	18.5	18.5	18.5	18.5	18.5	18.5	18.5

Tokelau CRF Table 3.B.1.3 Tokelau Swine: [3. Agriculture] [3.1 Livestock] [3.B Manure Management] [3.B.1 CH4 Emissions] [3.B.1.3 Swine] [Other (please specify)] [Pigs] (Part 3 of 3)

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.B Manure Management][3.B.1 CH4 Emissions][3.B.1.3 Swine][Other (please specify)][Pigs]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Population	1000s	2.4	2.36	2.22	2.08	1.93	1.79	1.65	1.65	1.65	1.65	1.65	1.65
Allocation by climate region													
Warm	%	100	100	100	100	100	100	100	100	101	102	103	104
Typical animal mass (average)	kg	80	80	80	80	80	80	80	80	80	80	80	80
VS daily excretion (average)	kg dm/head/day	NA											
CH ₄ producing potential (average)	m ³ /kg VS	NA											
Method													
CH ₄		T1											
Emission factor information													
CH ₄		D	D	D	D	D	D	D	D	D	D	D	D
Emissions													
CH ₄	kt	0.04	0.04	0.04	0.04	0.04	0.03	0.03	0.03	0.03	0.03	0.03	0.03
Implied emission factor													
CH ₄	kg/head/year	18.5	18.5	18.5	18.5	18.5	18.5	18.5	18.5	18.5	18.5	18.5	18.5

Tokelau CRF Table 3.B.1.4 Tokelau Poultry: [3. Agriculture] [3.1 Livestock] [3.B Manure Management] [3.B.1 CH4 Emissions] [3.B.1.4 Other livestock] [Tokelau_Poultry] (Part 1 of 3)

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.B Manure Management][3.B.1 CH4 Emissions][3.B.1.4 Other livestock][Poultry]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Population	1000s	3.44	3.5	3.39	3.29	3.18	3.08	2.97	2.84	2.71	2.58
Allocation by climate region											
Cool	%	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Temperate	%	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Warm	%	100	100	100	100	100	100	100	100	100	100
Typical animal mass (average)	kg	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
VS daily excretion (average)	kg dm/head/day	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
CH ₄ producing potential (average)	m ³ /kg VS	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Method											
CH ₄		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
Emission factor information											
CH ₄		D	D	D	D	D	D	D	D	D	D
Emissions											
CH ₄	kt	0.0001032	0.000105	0.0001018	0.0000986	0.0000955	0.0000923	0.0000891	0.0000852	0.0000813	0.0000774
Implied emission factor											
CH ₄	kg/head/year	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03

Tokelau CRF Table 3.B.1.4 Tokelau Poultry: [3. Agriculture] [3.1 Livestock] [3.B Manure Management] [3.B.1 CH4 Emissions] [3.B.1.4 Other livestock] [Tokelau_Poultry] (Part 2 of 3)

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.B Manure Management][3.B.1 CH4 Emissions][3.B.1.4 Other livestock][Poultry]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Population	1000s	2.45	2.32	2.23	2.14	2.05	1.96	1.87	1.71	1.55	1.39
Allocation by climate region											
Cool	%	NO									
Temperate	%	NO									
Warm	%	100	100	100	100	100	100	100	100	100	100
Typical animal mass (average)	kg	NA									
VS daily excretion (average)	kg dm/head/day	NA									

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.B Manure Management][3.B.1 CH4 Emissions][3.B.1.4 Other livestock][Poultry]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
CH ₄ producing potential (average)	m ³ /kg VS	NA									
Method											
CH ₄		T1									
Emission factor information											
CH ₄		D	D	D	D	D	D	D	D	D	D
Emissions											
CH ₄	kt	0.0000735	0.0000695	0.0000669	0.0000642	0.0000615	0.0000589	0.0000562	0.0000514	0.0000465	0.0000416
Implied emission factor											
CH ₄	kg/head/year	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03

Tokelau CRF Table 3.B.1.4 Tokelau Poultry: [3. Agriculture] [3.1 Livestock] [3.B Manure Management] [3.B.1 CH4 Emissions] [3.B.1.4 Other livestock] [Tokelau_Poultry] (Part 3 of 3)

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.B Manure Management][3.B.1 CH4 Emissions][3.B.1.4 Other livestock][Poultry]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Population	1000s	1.23	1.06	0.98	0.89	0.8	0.71	0.62	0.62	0.62	0.62	0.62	0.62
Allocation by climate region													
Cool	%	NO											
Temperate	%	NO											
Warm	%	100	100	100	100	100	100	100	100	100	100	100	100
Typical animal mass (average)	kg	NA											
VS daily excretion (average)	kg dm/head/day	NA											
CH ₄ producing potential (average)	m ³ /kg VS	NA											
Method													
CH ₄		T1											

[Sectors/Totals][3. Agriculture][3.1 Livestock][3.B Manure Management][3.B.1 CH4 Emissions][3.B.1.4 Other livestock][Poultry]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Emission factor information													
CH ₄		D	D	D	D	D	D	D	D	D	D	D	D
Emissions													
CH ₄	kt	0.0000368	0.0000319	0.0000293	0.0000267	0.000024	0.0000214	0.0000187	0.0000187	0.0000187	0.0000187	0.0000187	0.0000187
Implied emission factor													
CH ₄	kg/head/year	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03

Tokelau CRF Table 5.A: [5. Waste] [5.A Solid Waste Disposal] (Part 1 of 3)

[5. Waste][5.A Solid Waste Disposal]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Method											
CO ₂		NA									
CH ₄		T1									
Emission factor information											
CO ₂		NA									
CH ₄		D	D	D	D	D	D	D	D	D	D
Emissions	kt CO ₂ -equivalent	0.39	0.39	0.39	0.39	0.38	0.38	0.38	0.38	0.37	0.37
CO ₂	kt	NA									
CH ₄	kt	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.01	0.01
NO _x	kt	NE									
CO	kt	NE									
NM VOC	kt	NE									

Tokelau CRF Table 5.A: [5. Waste] [5.A Solid Waste Disposal] (Part 2 of 3)

[5. Waste][5.A Solid Waste Disposal]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Method											
CO ₂		NA									
CH ₄		T1									
Emission factor information											
CO ₂		NA									
CH ₄		D	D	D	D	D	D	D	D	D	D
Emissions	kt CO ₂ -equivalent	0.37	0.37	0.36	0.36	0.35	0.35	0.34	0.33	0.32	0.31
CO ₂	kt	NA									
CH ₄	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
NO _x	kt	NE									
CO	kt	NE									
NM VOC	kt	NE									

Tokelau CRF Table 5.A: [5. Waste] [5.A Solid Waste Disposal] (Part 3 of 3)

[5. Waste][5.A Solid Waste Disposal]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Method													
CO ₂		NA											
CH ₄		T1											
Emission factor information													
CO ₂		NA											
CH ₄		D	D	D	D	D	D	D	D	D	D	D	D
Emissions	kt CO ₂ -equivalent	0.31	0.31	0.3	0.3	0.3	0.3	0.3	0.3	0.31	0.31	0.31	0.31
CO ₂	kt	NA											
CH ₄	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
NO _x	kt	NE											
CO	kt	NE											
NM VOC	kt	NE											

Tokelau CRF Table 5.A.3: [5. Waste] [5.A Solid Waste Disposal] [5.A.3 Uncategorized Waste Disposal Sites] (Part 1 of 3)

[5. Waste][5.A Solid Waste Disposal][5.A.3 Uncategorized Waste Disposal Sites]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Annual waste at the SWDS	kt	0.54	0.53	0.53	0.53	0.52	0.52	0.52	0.52	0.51	0.51
MCF		0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
DOCf	%	50	50	50	50	50	50	50	50	50	50
Method											
CO ₂		NA									
CH ₄		T1									
Emission factor information											
CO ₂		NA									
CH ₄		D	D	D	D	D	D	D	D	D	D
Emissions											
CO ₂	kt	NA									
CH ₄											
Emissions	kt	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.01	0.01
Amount of CH ₄ flared	kt	NO									
Amount of CH ₄ for energy recovery	kt	NO									
NO _x	kt	NE									
CO	kt	NE									
NM VOC	kt	NE									
Implied emission factor											
CO ₂	t/t	NA									
CH ₄	t/t	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03

Tokelau CRF Table 5.A.3: [5. Waste] [5.A Solid Waste Disposal] [5.A.3 Uncategorized Waste Disposal Sites] (Part 2 of 3)

[5. Waste][5.A Solid Waste Disposal][5.A.3 Uncategorized Waste Disposal Sites]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Annual waste at the SWDS	kt	0.5	0.5	0.48	0.46	0.44	0.42	0.4	0.4	0.4	0.41
MCF		0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
DOCf	%	50	50	50	50	50	50	50	50	50	50
Method											
CO ₂		NA									
CH ₄		T1									
Emission factor information											
CO ₂		NA									
CH ₄		D	D	D	D	D	D	D	D	D	D
Emissions											
CO ₂	kt	NA									
CH ₄											
Emissions	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Amount of CH ₄ flared	kt	NO									
Amount of CH ₄ for energy recovery	kt	NO									
NO _x	kt	NE									
CO	kt	NE									
NMVOC	kt	NE									
Implied emission factor											
CO ₂	t/t	NA									
CH ₄	t/t	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03

Tokelau CRF Table 5.A.3: [5. Waste] [5.A Solid Waste Disposal] [5.A.3 Uncategorized Waste Disposal Sites] (Part 3 of 3)

[5. Waste][5.A Solid Waste Disposal][5.A.3 Uncategorized Waste Disposal Sites]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Annual waste at the SWDS	kt	0.41	0.42	0.42	0.43	0.43	0.44	0.44	0.44	0.44	0.45	0.45	0.45
MCF		0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
DOCf	%	50	50	50	50	50	50	50	50	50	50	50	50
Method													
CO ₂		NA											
CH ₄		T1											
Emission factor information													
CO ₂		NA											
CH ₄		D	D	D	D	D	D	D	D	D	D	D	D
Emissions													
CO ₂	kt	NA											
CH ₄													
Emissions	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Amount of CH ₄ flared	kt	NO											
Amount of CH ₄ for energy recovery	kt	NO											
NO _x	kt	NE											
CO	kt	NE											
NMVOC	kt	NE											
Implied emission factor													
CO ₂	t/t	NA											
CH ₄	t/t	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03	0.03

Tokelau CRF Table 5.C.2.2.a: [5. Waste] [5.C Incineration and Open Burning of Waste] [5.C.2 Open Burning of Waste] [5.C.2.2 Non-biogenic] [5.C.2.2.a Municipal Solid Waste] (Part 1 of 3)

[5. Waste][5.C Incineration and Open Burning of Waste][5.C.2 Open Burning of Waste][5.C.2.2 Non-biogenic][5.C.2.2.a Municipal Solid Waste]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Amount of wastes incinerated/open burned	kt	0.54	0.53	0.53	0.53	0.52	0.52	0.52	0.52	0.51	0.51
Method											
CO ₂		T1									
CH ₄		T1									
N ₂ O		T1									
Emission factor information											
CO ₂		D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
CO ₂	kt	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.04	0.04	0.04
CH ₄	kt	0	0	0	0	0	0	0	0	0	0
N ₂ O	kt	0.0000455	0.0000446	0.0000445	0.0000443	0.0000441	0.0000439	0.0000438	0.0000434	0.0000431	0.0000428
Implied emission factor											
CO ₂	kg/t	86.73	86.73	86.73	86.73	86.73	86.73	86.73	86.73	86.73	86.73
CH ₄	kg/t	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5
N ₂ O	kg/t	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08

Tokelau CRF Table 5.C.2.2.a: [5. Waste] [5.C Incineration and Open Burning of Waste] [5.C.2 Open Burning of Waste] [5.C.2.2 Non-biogenic] [5.C.2.2.a Municipal Solid Waste] (Part 2 of 3)

[5. Waste][5.C Incineration and Open Burning of Waste][5.C.2 Open Burning of Waste][5.C.2.2 Non-biogenic][5.C.2.2.a Municipal Solid Waste]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Amount of wastes incinerated/open burned	kt	0.5	0.5	0.48	0.46	0.44	0.42	0.4	0.4	0.4	0.41
Method											
CO ₂		T1									
CH ₄		T1									
N ₂ O		T1									

[5. Waste][5.C Incineration and Open Burning of Waste][5.C.2 Open Burning of Waste][5.C.2.2 Non-biogenic][5.C.2.2.a Municipal Solid Waste]												
	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	
Emission factor information												
CO ₂		D	D	D	D	D	D	D	D	D	D	
CH ₄		D	D	D	D	D	D	D	D	D	D	
N ₂ O		D	D	D	D	D	D	D	D	D	D	
Emissions												
CO ₂	kt	0.04	0.04	0.04	0.04	0.04	0.04	0.03	0.03	0.04	0.04	
CH ₄	kt	0	0	0	0	0	0	0	0	0	0	
N ₂ O	kt	0.0000424	0.0000421	0.0000404	0.0000386	0.0000369	0.0000352	0.0000334	0.0000337	0.0000341	0.0000344	
Implied emission factor												
CO ₂	kg/t	86.73	86.73	86.73	86.73	86.73	86.73	86.73	86.73	86.73	86.73	
CH ₄	kg/t	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	
N ₂ O	kg/t	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	

Tokelau CRF Table 5.C.2.2.a: [5. Waste] [5.C Incineration and Open Burning of Waste] [5.C.2 Open Burning of Waste] [5.C.2.2 Non-biogenic] [5.C.2.2.a Municipal Solid Waste] (Part 3 of 3)

[5. Waste][5.C Incineration and Open Burning of Waste][5.C.2 Open Burning of Waste][5.C.2.2 Non-biogenic][5.C.2.2.a Municipal Solid Waste]													
	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Amount of wastes incinerated/open burned	kt	0.41	0.42	0.42	0.43	0.43	0.44	0.44	0.44	0.44	0.45	0.45	0.45
Method													
CO ₂		T1											
CH ₄		T1											
N ₂ O		T1											
Emission factor information													
CO ₂		D	D	D	D	D	D	D	D	D	D	D	D
CH ₄		D	D	D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D	D	D

[5. Waste][5.C Incineration and Open Burning of Waste][5.C.2 Open Burning of Waste][5.C.2.2 Non-biogenic][5.C.2.2.a Municipal Solid Waste]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Emissions													
CO ₂	kt	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04
CH ₄	kt	0	0	0	0	0	0	0	0	0	0	0	0
N ₂ O	kt	0.0000347	0.000035	0.0000355	0.0000359	0.0000364	0.0000369	0.0000373	0.0000374	0.0000374	0.0000376	0.0000376	0.0000376
Implied emission factor													
CO ₂	kg/t	86.73	86.73	86.73	86.73	86.73	86.73	86.73	86.73	86.73	86.73	86.73	86.73
CH ₄	kg/t	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5	6.5
N ₂ O	kg/t	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08

Tokelau CRF Table 5.D:[5. Waste] [5.D Wastewater Treatment and Discharge] (Part 1 of 3)

[5. Waste][5.D Wastewater Treatment and Discharge]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Method											
CH ₄		T1									
N ₂ O		T1									
Emission factor information											
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions	kt CO ₂ -equivalent	0.17	0.17	0.17	0.18	0.18	0.19	0.19	0.21	0.22	0.23
CH ₄	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
N ₂ O	kt	0.0000593	0.0000562	0.0000537	0.0000511	0.0000486	0.0000461	0.0000436	0.0000379	0.0000323	0.0000268
NO _{2x}	kt	NE									
CO	kt	NE									
NMVOC	kt	NE									
Additional information											
Population	1000s	1.57	1.54	1.53	1.52	1.52	1.51	1.51	1.5	1.48	1.47
Protein consumption	kg/person/yr	32.45	32.45	32.45	32.45	32.45	32.45	32.45	32.45	32.45	32.45
Fraction of nitrogen in protein		0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16

[5. Waste][5.D Wastewater Treatment and Discharge]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Factor of non-consumed protein added to the wastewater		1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1
Factor of industrial and commercial co-discharged protein into the sewer system		1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25
Degree of utilization of modern, centralized WWT plants	%	0	0	0	0	0	0	0	0	0	0

Tokelau CRF Table 5.D:[5. Waste] [5.D Wastewater Treatment and Discharge] (Part 2 of 3)

[5. Waste][5.D Wastewater Treatment and Discharge]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Method											
CH ₄		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
N ₂ O		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
Emission factor information											
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions	kt CO ₂ -equivalent	0.24	0.26	0.25	0.24	0.23	0.22	0.21	0.22	0.22	0.22
CH ₄	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
N ₂ O	kt	0.0000214	0.000016	0.0000145	0.0000131	0.0000117	0.0000104	0.0000092	0.0000087	0.0000081	0.0000075
NO _{2x}	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
CO	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
NMVOC	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
Additional information											
Population	1000s	1.46	1.45	1.39	1.33	1.27	1.21	1.15	1.16	1.17	1.18
Protein consumption	kg/person/yr	32.45	32.45	32.45	32.45	32.45	32.45	32.45	32.45	32.45	32.45
Fraction of nitrogen in protein		0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16
Factor of non-consumed protein added to the wastewater		1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1
Factor of industrial and commercial co-discharged protein into the sewer system		1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25
Degree of utilization of modern, centralized WWT plants	%	0	0	0	0	0	0	0	0	0	0

Tokelau CRF Table 5.D:[5. Waste] [5.D Wastewater Treatment and Discharge] (Part 3 of 3)

[5. Waste][5.D Wastewater Treatment and Discharge]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Method													
CH ₄		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
N ₂ O		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
Emission factor information													
CH ₄		D	D	D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D	D	D
Emissions	kt CO ₂ -equivalent	0.23	0.23	0.24	0.24	0.25	0.26						
CH ₄	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
N ₂ O	kt	0.0000069	0.0000063	0.0000051	0.0000039	0.0000026	0.0000013	NO	NO	NO	NO	NO	NO
No ₂ x	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
CO	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
NMVOC	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
Additional information													
Population	1000s	1.19	1.21	1.22	1.24	1.25	1.27	1.28	1.29	1.29	1.29	1.29	1.29
Protein consumption	kg/person/yr	32.45	32.45	32.45	32.45	32.45	32.45	32.45	32.45	32.45	32.45	32.45	32.45
Fraction of nitrogen in protein		0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16	0.16
Factor of non-consumed protein added to the wastewater		1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1
Factor of industrial and commercial co-discharged protein into the sewer system		1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25
Degree of utilization of modern, centralized WWT plants	%	0	0	0	0	0	0	0	0	1	2	3	4

Tokelau CRF Table 5.D.1: [5. Waste] [5.D Wastewater Treatment and Discharge] [5.D.1 Domestic Wastewater] (Part 1 of 3)

[5. Waste][5.D Wastewater Treatment and Discharge][5.D.1 Domestic Wastewater]	Unit	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
Total organic product	kt DC	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04	0.04
Sludge removed	kt DC	NO									
N in effluent	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0	0	0
Method											
CH ₄		T1									
N ₂ O		T1									
Emission factor information											
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
CH₄											
Emissions	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Amount of CH ₄ flared	kt	NO									
Amount of CH ₄ for energy recovery	kt	NO									
N ₂ O	kt	0.0000593	0.0000562	0.0000537	0.0000511	0.0000486	0.0000461	0.0000436	0.0000379	0.0000323	0.0000268
No _x	kt	NE									
CO	kt	NE									
NMVOC	kt	NE									
Implied emission factor											
CH ₄	kg/kg DC	0.14	0.14	0.15	0.16	0.16	0.17	0.18	0.19	0.21	0.22
N ₂ O	kg N ₂ O-N/kg N	0	0	0	0	0	0	0	0	0	0

Tokelau CRF Table 5.D.1: [5. Waste] [5.D Wastewater Treatment and Discharge] [5.D.1 Domestic Wastewater] (Part 2 of 3)

[5. Waste][5.D Wastewater Treatment and Discharge][5.D.1 Domestic Wastewater]	Unit	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Total organic product	kt DC	0.04	0.04	0.04	0.04	0.03	0.03	0.03	0.03	0.03	0.03
Sludge removed	kt DC	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
N in effluent	kt	0	0	0	0	0	0	0	0	0	0.0009546
Method											
CH ₄		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
N ₂ O		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
Emission factor information											
CH ₄		D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D
Emissions											
CH₄											
Emissions	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Amount of CH ₄ flared	kt	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Amount of CH ₄ for energy recovery	kt	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
N ₂ O	kt	0.0000214	0.000016	0.0000145	0.0000131	0.0000117	0.0000104	0.0000092	0.0000087	0.0000081	0.0000075
No _x	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
CO	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
NMVOC	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
Implied emission factor											
CH ₄	kg/kg DC	0.24	0.25	0.26	0.26	0.26	0.26	0.27	0.27	0.27	0.27
N ₂ O	kg N ₂ O-N/kg N	0	0	0	0	0	0	0	0	0	0

Tokelau CRF Table 5.D.1: [5. Waste] [5.D Wastewater Treatment and Discharge] [5.D.1 Domestic Wastewater] (Part 3 of 3)

[5. Waste][5.D Wastewater Treatment and Discharge][5.D.1 Domestic Wastewater]	Unit	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021
Total organic product	kt DC	0.03	0.03	0.03	0.03	0.03	0.03	0.04	0.04	0.04	0.04	0.04	0.04
Sludge removed	kt DC	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
N in effluent	kt	0.0008781	0.0008	0.0006485	0.0004927	0.0003327	0.0001685	NO	NO	NO	NO	NO	NO
Method													
CH ₄		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
N ₂ O		T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1	T1
Emission factor information													
CH ₄		D	D	D	D	D	D	D	D	D	D	D	D
N ₂ O		D	D	D	D	D	D	D	D	D	D	D	D
Emissions													
CH₄													
Emissions	kt	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Amount of CH ₄ flared	kt	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
Amount of CH ₄ for energy recovery	kt	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO	NO
N ₂ O	kt	0.0000069	0.0000063	0.0000051	0.0000039	0.0000026	0.0000013	NO	NO	NO	NO	NO	NO
No _x	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
CO	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
NMVOC	kt	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
Implied emission factor													
CH ₄	kg/kg DC	0.28	0.28	0.28	0.29	0.29	0.3	0.3	0.3	0.3	0.3	0.3	0.3
N ₂ O	kg N ₂ O-N/kg N	0	0	0	0	0	0	NO	NO	NO	NO	NO	NO

Annex 8: Agricultural emissions from fertilisers and by livestock type

A8.1 Agricultural emissions from fertilisers

Fertilisers provide the nutrients to grow and nourish pastures and crops. Nitrogen, phosphate, potassium and sulphur are the four most important nutrients for pasture and crop growth and sustainable food production.

New Zealand's farmers use both organic and synthetic nitrogen fertilisers. The main type of synthetic nitrogen fertiliser used in New Zealand is urea, followed by smaller amounts of diammonium phosphate (DAP) and ammonium sulphate. Urea is mainly applied to dairy pastures to boost pasture growth during the autumn and spring months.

All nitrogen fertilisers provide nitrogen inputs to agricultural soils that result in direct and indirect emissions of nitrous oxide (N₂O) (see figure 5.5.1 in chapter 5). Urea also releases carbon dioxide (CO₂).

Emissions from organic fertilisers come solely from animal manure. Most animal manure in New Zealand is excreted directly onto pasture, but some manure from dairy farms is kept in manure management systems and applied to soils as an organic fertiliser (see table 5.3.2 in chapter 5, for further details). Some manure is also collected but not stored; rather, it is spread directly onto pasture daily (e.g., swine manure and some dairy manure).

Emissions of N₂O from all synthetic (including urea) nitrogen fertilisers are reported in categories 3.D.1.1 and 3.D.1.2 respectively. Emissions of CO₂ from urea are not included under synthetic nitrogen fertilisers and are reported under a dedicated category 3.H.

2021

In 2021, the combined effect of synthetic and organic nitrogen fertilisers totalled 32.2 per cent of emissions from the *Agricultural soils* category and 6.2 per cent of total agricultural emissions (when CO₂-e from urea is included).

Table A8.1.1 shows comparisons of both N₂O and CO₂ emissions from fertilisers to New Zealand's national totals for each gas and New Zealand's gross emissions.

Table A8.1.1 Direct and indirect emissions by fertiliser in 2021

Fertiliser type	Gas/source	Emissions kt CO ₂ -e	Percentage of	
			N ₂ O emissions from Agriculture soils by gas (%)	All emissions from Agriculture (%)
Synthetic nitrogen fertiliser	Direct N ₂ O emissions	1,398.9	19.1	3.7
	Urea N ₂ O	958.7	13.1	2.5
	Other synthetic nitrogen fertilisers	440.2	6.0	1.2
	Indirect N ₂ O emissions from all synthetic nitrogen fertilisers	296.9	4.1	0.8
	All N ₂ O (direct + indirect) from synthetic nitrogen fertilisers	1,695.8	23.2	4.5
	CO ₂ from urea	553.2	NA	1.5

Fertiliser type	Gas/source	Emissions kt CO ₂ -e	Percentage of	
			N ₂ O emissions from Agriculture soils by gas (%)	All emissions from Agriculture (%)
Organic fertiliser	Direct N ₂ O emissions	73.8	1.0	0.2
	Indirect N ₂ O emissions	30.0	0.4	0.1
	All N ₂ O (direct + indirect) from organic fertilisers	103.8	1.4	0.3

Note: NA = not applicable. Columns may not add up due to rounding.

1990–2021

The total amount of fertilisers applied to agricultural soils in New Zealand has significantly increased since 1990. Since 1990, synthetic nitrogen fertiliser applied to agricultural land has increased by 644.1 per cent, while organic fertiliser use has grown by 173.6 per cent (table A8.1.2).

Table A8.1.2 Use of fertilisers in New Zealand in 1990 and 2021

Fertiliser type	1990			2021			Change in the use between 1990 and 2021	
	Application	Percentage of		Application	Percentage of		tonnes (N)	(%)
	tonnes (N)	synthetic nitrogen fertiliser (%)	all fertilisers (%)	tonnes (N)	synthetic nitrogen fertiliser (%)	all fertilisers (%)		
Synthetic nitrogen fertiliser (ammonium phosphates, for example, DAP)	34,679	58.5	46.3	94,000	21.3	19.4	59,321	171.1
Urea	24,586	41.5	32.8	347,000	78.7	71.7	322,414	1,311.4
Total synthetic nitrogen fertilisers (urea + ammonium phosphates)	59,265	100.0	79.1	441,000	100.0	91.2	381,735	644.1
Organic fertilisers (animal manure applied to soils)	15,644	NA	20.9	42,803	NA	8.8	27,159	173.6

Note: DAP = diammonium phosphate; NA = not applicable. Columns may not add up due to rounding.

Between 1990 and 2021, N₂O emissions from synthetic nitrogen fertiliser (both direct and indirect emissions, including urea) have increased by 516.3 per cent, while total emissions from these fertilisers (N₂O and CO₂) have increased by 614.4 per cent. For the same period, total emissions from organic fertilisers increased by 119.7 per cent (see table A8.1.3).

In 1990 and 2021 respectively, 0.8 per cent and 4.5 per cent of total agricultural emissions originated from N₂O from synthetic nitrogen fertiliser. Total emissions from synthetic nitrogen fertiliser (including urea) have increased from 0.9 per cent to 6.0 per cent of total agricultural emissions for 1990 and 2021 respectively (see chapter 5, for further details).

Table A8.1.3 Emissions from fertilisers in 1990 and 2021

			Synthetic nitrogen fertilisers	Organic fertilisers
1990	N ₂ O emissions	kt CO ₂ -e	275.1	47.2
	CO ₂ emissions	kt	39.2	NA
	Total emissions	kt CO₂-e	314.3	47.2
2021	N ₂ O emissions	kt CO ₂ -e	1,695.8	103.8
	CO ₂ emissions	kt	553.2	NA
	Total emissions	kt CO₂-e	2,249	103.8
Change in N ₂ O emissions between 1990 and 2021		kt CO ₂ -e	1,420.6	56.6
Percentage change in N ₂ O emissions between 1990 and 2021		%	516.3	119.7
Change in all emissions between 1990 and 2021		kt CO ₂ -e	1,934.6	56.6
Percentage change in all emissions between 1990 and 2021		%	615.4	119.7

Note: NA = not applicable.

A8.2 Agricultural emissions by livestock type

This section covers distribution of all greenhouse gas emissions from the Agriculture sector by livestock type in 1990, 2020 and 2021, including the changes in emissions. Table A8.2.1 shows total emissions of all greenhouse gases across all categories of the Agriculture sector. For further details on emissions by gas and by category, refer to the common reporting format tables (sector 3 – Agriculture).

Table A8.2.1 Total emissions by livestock type in 1990, 2020 and 2021

Livestock type	1990	2020	2021	1990–2021		2020–2021	
		kt CO ₂ -e		kt CO ₂ -e	(%)	kt CO ₂ -e	(%)
Dairy cattle	7,832.2	17,665.6	17,428.7	9,596.5	122.5	-236.9	-1.3
Beef cattle	6,914.5	6,988.1	7,037.0	122.5	1.8	48.9	0.7
Sheep	16,112.0	9,172.2	9,011.6	-7,100.4	-44.1	-160.6	-1.8
Deer	499.2	554.9	540.7	41.5	8.3	-14.3	-2.6
Swine	102.1	65.9	69.7	-32.4	-31.7	3.8	5.8
Goats	263.2	28.8	34.8	-228.4	-86.8	6.0	21.0
Horses	78.6	32.3	33.7	-44.9	-57.2	1.4	4.2
Alpaca	0.1	2.6	2.1	2.0	1,808.0	-0.5	-19.1
Mules and asses	0.1	0.1	0.1	0.0	0.0	0.0	0.0
Poultry (including all types of poultry)	26.4	59.7	60.8	34.4	130.2	1.1	1.8
Total, all livestock types	31,828.2	34,570.1	34,219.1	2,390.9	7.0	-351.0	-0.1

Note: Columns may not add up due to rounding.